

# CS 474

## Homework 4

Name: Adharsh Kamath

NetID: ak128

---

### Problem 1

**Soln:** A group is defined as a set  $S$ , along with a binary operation  $\cdot : S \times S \rightarrow S$  and axioms:

1.  $\forall a, b, c. (a \cdot b) \cdot c = a \cdot (b \cdot c)$
2.  $\forall a. ((a \cdot e) = a \wedge (e \cdot a) = a)$
3.  $\forall a \exists b. ((b \cdot a) = e \wedge (a \cdot b) = e)$

where  $e$  is the special constant that is the identity element of the group.

Task 1:

$$G : \forall e'. ((\forall a. ((e \cdot a) = a \wedge (e' \cdot a) = a)) \implies (e = e'))$$

The first two axioms are in prenex normal form, and do not contain any existential quantifiers. Skolemizing the third axiom (by replacing  $\exists b$  with a function  $f(a)$ ), we get:

$$\forall a. (a \cdot f(a)) = e \wedge (a \cdot f(a)) = e$$

Simplifying the goal  $G$ , to prenex normal form:

$$\begin{aligned} & \forall e' \forall a. ((a \cdot e') = a \wedge (e' \cdot a) = a) \implies (e = e') \\ & \equiv \forall e' \neg(\forall a. ((a \cdot e') = a \wedge (e' \cdot a) = a)) \vee (e = e') \\ & \equiv \forall e' (\exists a. \neg((a \cdot e') = a \wedge (e' \cdot a) = a)) \vee (e = e') \end{aligned}$$

To show the validity of the goal, we need to show that the negation of the goal is unsatisfiable along with all the axioms.

$$\begin{aligned} \neg G & \equiv \neg(\forall e' (\exists a. \neg((a \cdot e') = a \wedge (e' \cdot a) = a)) \vee (e = e')) \\ & \equiv \exists e' \neg(\exists a. \neg((a \cdot e') = a \wedge (e' \cdot a) = a)) \wedge \neg(e = e') \\ & \equiv \exists e' (\forall a. ((a \cdot e') = a \wedge (e' \cdot a) = a)) \wedge \neg(e = e') \end{aligned}$$

Skolemizing (with a constant  $e''$  since  $\exists e'$  is the outermost existential quantifier), we get:

$$\forall a. ((a \cdot e'') = a \wedge (e'' \cdot a) = a \wedge \neg(e = e''))$$

Combining all the axioms, we have:

$$\begin{aligned} & \forall a, b, c. (a \cdot b) \cdot c = a \cdot (b \cdot c) \\ & \forall a. ((a \cdot e) = a \wedge (e \cdot a) = a) \\ & \forall a. ((a \cdot f(a)) = e \wedge (f(a) \cdot a) = e) \\ & \forall a. ((a \cdot e'') = a \wedge (e'' \cdot a) = a \wedge \neg(e = e'')) \end{aligned}$$

Instantiating with depth-0 terms  $(e, e'')$ , we get a large set (conjunction) of formulae :

$$\begin{aligned}
(e \cdot e) \cdot e &= e \cdot (e \cdot e) \\
(e \cdot e) \cdot e'' &= e \cdot (e \cdot e'') \\
(e \cdot e'') \cdot e &= e \cdot (e'' \cdot e) \\
(e \cdot e'') \cdot e'' &= e \cdot (e'' \cdot e'') \\
(e'' \cdot e) \cdot e &= e'' \cdot (e \cdot e) \\
(e'' \cdot e) \cdot e'' &= e'' \cdot (e \cdot e'') \\
(e'' \cdot e'') \cdot e &= e'' \cdot (e'' \cdot e) \\
(e'' \cdot e'') \cdot e'' &= e'' \cdot (e'' \cdot e'') \\
(e \cdot e) &= e \wedge (e \cdot e) = e \\
(e'' \cdot e) &= e \wedge (e \cdot e'') = e \\
(e \cdot f(e)) &= e \wedge (f(e) \cdot (e)) = e \\
(e'' \cdot f(e'')) &= e \wedge (f(e'') \cdot (e'')) = e \\
(e \cdot e'') &= e \wedge (e'' \cdot e) = e \wedge \neg(e = e'') \\
(e'' \cdot e'') &= e'' \wedge (e'' \cdot e'') = e'' \wedge \neg(e = e'')
\end{aligned}$$

In this list, we can find two conjuncts that are contradictory:

$$\begin{aligned}
(e \cdot e'') &= e'' \wedge (e'' \cdot e) = e'' \\
(e'' \cdot e) &= e \wedge (e \cdot e'') = e \wedge \neg(e = e'')
\end{aligned}$$

Hence, there exists no model that satisfies all the formulae. Therefore, the goal  $G$  is valid.

The corresponding Z3 program is at the following URL: [https://github.com/adharshkamath/logic-hw-4/blob/main/p1\\_1.py](https://github.com/adharshkamath/logic-hw-4/blob/main/p1_1.py)

Task 2: Given the skolemized set of axioms:

$$\begin{aligned}
\forall a, b, c. (a \cdot b) \cdot c &= a \cdot (b \cdot c) \\
\forall a. ((a \cdot e) = a \wedge (e \cdot a) = a) \\
\forall a. ((a \cdot f(a)) = e \wedge (a \cdot f(a)) = e)
\end{aligned}$$

we can see that  $f$  is the inverse function. We can simplify the goal to

$$G : \forall a, b. ((a \cdot b = e) \wedge (b \cdot a = e)) \implies (b = f(a))$$

Negating this goal gives us:

$$\neg G : \exists a, b. ((a \cdot b = e) \wedge (b \cdot a = e)) \wedge \neg(b = f(a))$$

Skolemizing (with constants  $a', b'$ ) gives us:

$$((a' \cdot b' = e) \wedge (b' \cdot a' = e)) \wedge \neg(b' = f(a'))$$

Instantiating the axioms with depth-0 terms  $(a', b', e)$ , we get a long list (conjunction) of formulae:

$$\begin{aligned}
&(a' \cdot a') \cdot a' = a' \cdot (a' \cdot a') \\
&(a' \cdot a') \cdot b' = a' \cdot (a' \cdot b') \\
&(a' \cdot b') \cdot a' = a' \cdot (b' \cdot a') \\
&(a' \cdot b') \cdot b' = a' \cdot (b' \cdot b') \\
&(b' \cdot a') \cdot a' = b' \cdot (a' \cdot a') \\
&(b' \cdot a') \cdot b' = b' \cdot (a' \cdot b') \\
&(b' \cdot b') \cdot a' = b' \cdot (b' \cdot a') \\
&(b' \cdot b') \cdot b' = b' \cdot (b' \cdot b') \\
&(a' \cdot a') \cdot e = a' \cdot (a' \cdot e) \\
&(a' \cdot e) \cdot a' = a' \cdot (e \cdot a') \\
&(a' \cdot e) \cdot e = a' \cdot (e \cdot e) \\
&(e \cdot a') \cdot a' = e \cdot (a' \cdot a') \\
&(e \cdot a') \cdot e = e \cdot (a' \cdot e) \\
&(e \cdot e) \cdot a' = e \cdot (e \cdot a') \\
&(e \cdot e) \cdot e = e \cdot (e \cdot e) \\
&(b' \cdot b') \cdot e = b' \cdot (b' \cdot e) \\
&(b' \cdot e) \cdot b' = b' \cdot (e \cdot b') \\
&(b' \cdot e) \cdot e = b' \cdot (e \cdot e) \\
&(e \cdot b') \cdot b' = e \cdot (b' \cdot b') \\
&(e \cdot b') \cdot e = e \cdot (b' \cdot e) \\
&(e \cdot e) \cdot b' = e \cdot (e \cdot b') \\
&(a' \cdot b') \cdot e = a' \cdot (b' \cdot e) \\
&(a' \cdot e) \cdot b' = a' \cdot (e \cdot b') \\
&(e \cdot a') \cdot b' = e \cdot (a' \cdot b') \\
&(e \cdot b') \cdot a' = e \cdot (b' \cdot a') \\
&(b' \cdot a') \cdot e = b' \cdot (a' \cdot e) \\
&(b' \cdot e) \cdot a' = b' \cdot (e \cdot a') \\
&(a' \cdot e) = a' \wedge (e \cdot a') = a' \\
&(b' \cdot e) = b' \wedge (e \cdot b') = b' \\
&(e \cdot e) = e \wedge (e \cdot e) = e \\
&(a' \cdot f(a')) = e \wedge (f(a') \cdot a') = e \\
&(b' \cdot f(b')) = e \wedge (f(b') \cdot b') = e \\
&(e \cdot f(e)) = e \wedge (f(e) \cdot e) = e
\end{aligned}$$

In this list, we can find two conjuncts that are contradictory:

$$\begin{aligned}
&(a' \cdot f(a')) = e \wedge (f(a') \cdot a') = e \\
&(b' \cdot f(b')) = e \wedge (f(b') \cdot b') = e
\end{aligned}$$

The Z3 program for this task is at the following URL: [https://github.com/adharshkamath/logic-hw-4/blob/main/p1\\_pt2.py](https://github.com/adharshkamath/logic-hw-4/blob/main/p1_pt2.py)

---

## Problem 2

### Soln:

(a)

The given formula  $\varphi$

$$\varphi = y \leq x \wedge x \leq y \wedge f(y) = f(7) \wedge x \leq 5$$

contains terms from  $T_{UIF}$  and  $T_N$  (theory of UIF and theory of natural numbers).

Replacing  $f(7)$  with  $f(w)$ , and adding  $w = 7$  as an additional formula, we get:

$$\varphi' = y \leq x \wedge x \leq y \wedge x \leq 5 \wedge w = 7 \wedge f(y) = f(w)$$

The part of the formula in  $\Sigma_{UIF}$  :

$$F_{UIF} : f(y) = f(w)$$

and the part of the formula in  $\Sigma_N$  :

$$F_N : y \leq x \wedge x \leq y \wedge x \leq 5 \wedge w = 7$$

The conjunct  $F_{UIF} \wedge F_N$  is  $(T_{UIF} \cup T_N)$ -equisatisfiable to  $\varphi$ .

(b)

The shared variables between  $F_{UIF}$  and  $F_N$  are:

$$V = \text{shared}(F_{UIF}, F_N) = \{y, w\}$$

The two possible equivalence relations and their arrangements are:

$$\begin{aligned} E_1 &= \{\{y\}, \{w\}\}, & \alpha_1(E_1, V) : y &= w \\ E_2 &= \{\{y, w\}\}, & \alpha_2(E_2, V) : y &\neq w \end{aligned}$$

Simplifying  $F_N$ , we can write:

$$\begin{aligned} F_N &= y \leq x \wedge x \leq y \wedge x \leq 5 \wedge w = 7 \\ &\equiv x = y \wedge x \leq 5 \wedge w = 7 \end{aligned}$$

We can see that  $F_N$  is satisfiable over natural numbers only under the arrangement  $\alpha_2$  (since  $y \leq 5$  and  $w = 7$ ). A satisfying model for  $F_N$  is:  $\{x = 5, y = 5, w = 7\}$

We can see that  $F_{UIF}$  is satisfiable under this arrangement only if  $f$  maps all arguments to a constant (since we need to satisfy  $f(y) = f(w) \wedge y \neq w$ ). A satisfying model for  $F_{UIF}$  is:  $\{f(y) = c\}$  where  $c$  is any arbitrary constant.

From the above, we can derive a model for the original formula  $\varphi$ . A satisfying model for  $\varphi$  is:  $\{x = 5, y = 5, f(t) = c\}$  where  $t$  is any term in the combined domains.

---

Problem 3

**Soln:**

(a)

Given  $f : 2^N \rightarrow 2^N$  that is defined as

$$f(S) = \{2\} \cup \{y \mid y = 2x, x \in S\}$$

---