



**Weekly Model Test
Based on
IOE Entrance Syllabus**

In collaboration with



AEC | **Acme Engineering College**
(Affiliated to Purbanchal University)

Programs Offered:

Computer Electronics and Communication Engineering
Civil Engineering
Architecture

Set A/B

27th Shrawan, 2080

TRIBHUVAN UNIVERSITY
INSTITUTE OF ENGINEERING
B.E. / B.Arch. Entrance Examination
Set A

Full Marks: 140 Time: 2hrs.

SECTION-I

Select the best alternatives: (60×1 = 60)

1. The domain of the real function $f(x) = \sqrt{x^2 + x - 6} + \sqrt{4 + 3x - x^2}$ is
 a. [2,4] b. [1,4]
 c. [-3,1] d. $(-\infty, -3) \cup [4, \infty)$

for the two components to be meaningful:

- (i) $x^2 + x - 6 \geq 0 \Rightarrow (x + 3)(x - 2) \geq 0 \Rightarrow x \leq -3$ or $2 \leq x \dots \dots (i)$
 (ii) $4 + 3x - x^2 \geq 0 \Rightarrow x^2 - 3x - 4 \leq 0 \Rightarrow (x - 4)(x + 1) \leq 0 \Rightarrow -1 \leq x \leq 4 \dots \dots (ii)$
 Combining (i) and (ii)
 $2 \leq x \leq 4 \Rightarrow x \in [2, 4]$

2. If $f(x) = \log\left(\frac{1+x}{1-x}\right)$ then $f\left(\frac{2x}{1+x^2}\right)$ equals
 a. $f(x)$ b. $2f(x)$
 c. $3f(x)$ d. $4f(x)$

$$f\left(\frac{2x}{1+x^2}\right) = \log\left(\frac{1+\frac{2x}{1+x^2}}{1-\frac{2x}{1+x^2}}\right)$$

$$\log\left(\frac{(1+x)^2}{(1-x)^2}\right) = 2\log\left(\frac{1+x}{1-x}\right) = 2f(x)$$

3. Inequality $-6 \leq x \leq 1$ by using absolute value sign is

- a. $|x + 3| \leq 4$ b. $|3x + 1| \leq 2$
 c. $|2x + 5| \leq 7$ d. $|x - 5| \leq 1$

$$-6 \leq x \leq 1 \Rightarrow \left[\frac{-6+1}{2} = \frac{-5}{2}\right]$$

$$\Rightarrow -6 + \frac{5}{2} \leq x + \frac{5}{2} \leq 1 + \frac{5}{2} \text{ [subtract } -\frac{5}{2} \text{ from all]}$$

$$\Rightarrow -7 \leq 2x + 5 \leq 7 \Rightarrow |2x + 5| \leq 7$$

4. If $\begin{vmatrix} -a^2 & ab & ac \\ ab & -b^2 & bc \\ ac & bc & -c^2 \end{vmatrix} = \lambda a^2 b^2 c^2$ then λ equals
 a. 1 b. 2

c. 3 d. 4

$$abc \begin{vmatrix} -a & b & c \\ a & -b & c \\ a & b & -c \end{vmatrix} =$$

$$\lambda a^2 b^2 c^2 \Rightarrow \begin{vmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{vmatrix} = \lambda \Rightarrow \lambda = 4$$

5. The true statement is
 a. $1-i < 1+i$ b. $1+2i > 1-2i$
 c. $2i > 1$ d. none

there is no natural ordering of the complex numbers. They are either equal or unequal.

6. If the roots of the quadratic $x^2 + ax + c = 0$ differ by 1, then a^2 is always equal to
 a. $4c + 1$ b. $4c - 1$
 c. $4c$ d. $4c \pm 1$

$$\alpha + \beta = -a, \alpha\beta = c, \alpha - \beta = -1$$

$$\therefore \alpha = \frac{1-a}{2} \text{ and } \beta = -\left(\frac{1+a}{2}\right)$$

$$\therefore \left(\frac{1-a}{2}\right)\left(\frac{1+a}{2}\right) = c \Rightarrow a^2 = 1 + 4c$$

7. If ${}^nC_{r-1} = 36$, ${}^nC_r = 84$ and ${}^nC_{r+1} = 126$ then the value of n and r are respectively equal to
 a. 9,3 b. 3,9
 c. 2,7 d. 7,2

${}^nC_{r-1} = 36$, ${}^nC_r = 84$ and ${}^nC_{r+1} = 126$
 put the value of n and r from option and check, only option (a) satisfies the above condition.

8. If the 21st and 22nd terms in the expansion of $(1+x)^{44}$ are equal then the value of x must be
 a. $7/8$ b. $9/8$
 c. $11/8$ d. $13/8$

In expansion of $(a+x)^n$, $t_{r+1} = (r+1)^{\text{th}}$ terms = $C(n,r) a^{n-r} x^r$
 In $(1+x)^{44}$, $t_{20+1} = 21^{\text{st}}$ term = $C(44, 20) x^{20}$
 $t_{21+1} = 22^{\text{nd}}$ term = $C(44, 21) x^{21}$
 $\therefore C(44, 20) x^{20} = C(44, 21) x^{21} \Rightarrow x = \frac{7}{8}$

9. The general values of x in the equation $2\sin 2x = \sin x$ is

a. $n\pi + \frac{\pi}{2}$ b. $\frac{n\pi}{4}$

c. $2n\pi + \frac{\pi}{2}$ d. $n\pi$

$$4\sin x \cos x - \sin x = 0 \Rightarrow \sin x(4\cos x - 1) = 0$$

$$\therefore \sin x = 0 \Rightarrow x = n\pi + (-1)^n \cdot 0$$


$$[\sin x = \sin \alpha \Rightarrow x = n\pi + (-1)^n \cdot \alpha]$$

$$\therefore x = n\pi$$

10. The circumcenter of a right-angled triangle lies

- a. in hypotenuse of triangle
b. inside triangle
c. outside triangle
d. on vertex of triangle

The circumcenter of right-angled triangle is at midpoint of hypotenuse.



11. $\lim_{n \rightarrow \infty} \left(\frac{1}{n^2} + \frac{2}{n^2} + \frac{3}{n^2} + \dots + \frac{n}{n^2} \right) =$

- a. $\frac{1}{3}$ b. $\frac{1}{4}$
c. $\frac{1}{2}$ d. 1

$$\lim_{n \rightarrow \infty} \left(\frac{1}{n^2} + \frac{2}{n^2} + \frac{3}{n^2} + \dots + \frac{n}{n^2} \right)$$

$$= \lim_{n \rightarrow \infty} \frac{1}{n^2} \left(\frac{n(n+1)}{2} \right) = \lim_{n \rightarrow \infty} \frac{n(1 + \frac{1}{n})}{2n} =$$

$$\frac{1+0}{2} = \frac{1}{2}$$

12. If $y = \frac{1 - \tan x}{\sec x}$ then the value of $\frac{dy}{dx} =$

- a. $-\sin x - \cos x$ b. $\frac{2\sec x}{1 - \tan x}$
c. $\sin x + \cos x$ d. $2\sec x (1 + \tan^2 x)$

$$y = \frac{1 - \tan x}{\sec x} = \cos x - \sin x$$

$$\Rightarrow \frac{dy}{dx} = -\sin x - \cos x$$

13. If $y = \sec^{-1} \left(\frac{1}{\sqrt{1-x^2}} \right)$ then $\frac{dy}{dx} =$

- a. $-\frac{2}{\sqrt{1-x^2}}$ b. $\frac{2x}{\sqrt{1+(1-x^2)^2}}$
c. $\frac{2}{1+x}$ d. $\frac{1}{\sqrt{1-x^2}}$

$$y = \cos^{-1} \sqrt{1-x^2}, \text{ put } x = \sin \theta$$

$$y = \theta = \sin^{-1} x \Rightarrow \frac{dy}{dx} = \frac{d}{dx} \sin^{-1} x \Rightarrow \frac{dy}{dx}$$

$$= \frac{1}{\sqrt{1-x^2}}$$

14. Value of $\int \frac{e^{ax}}{e^{ax} + 1} dx =$

- a. $\frac{2(e^{ax} + 1)}{a} + c$ b. $\frac{1}{a} \log(e^{ax} + 1)$
c. $2a(e^{ax} + 1) + c$ d. $a \frac{\log(e^{ax})}{2} + c$

$$\int \frac{e^{ax}}{e^{ax} + 1} dx = \frac{1}{a} \int \frac{a \cdot e^{ax}}{e^{ax} + 1} dx = \frac{1}{a} \log(e^{ax} + 1)$$

15. If \vec{a} and \vec{b} are two unit vectors such that $(\vec{a} + 2\vec{b})$ and $(5\vec{a} - 4\vec{b})$ are perpendicular to each other then the angle between \vec{a} and \vec{b} is

- a. $\frac{\pi}{2}$ b. $\frac{\pi}{3}$
c. $\frac{\pi}{4}$ d. $\frac{\pi}{6}$

$$(\vec{a} + 2\vec{b}) \cdot (5\vec{a} - 4\vec{b}) = 0$$

[Two vectors are \perp]

$$\Rightarrow 5a^2 - 4\vec{a} \cdot \vec{b} + 10\vec{a} \cdot \vec{b} - 8b^2 = 0$$

$$\Rightarrow \vec{a} \cdot \vec{b} = \frac{1}{2} \text{ [}\vec{a} \text{ \& } \vec{b} \text{ are unit vectors]}$$

$$\Rightarrow ab \cos \theta = \frac{1}{2} \Rightarrow \theta = \frac{\pi}{3}$$

16. The projection of $(\mathbf{i} + \mathbf{j} - \mathbf{k})$ on $(2\mathbf{i} + \mathbf{j} + \mathbf{k})$ is

- a. $\frac{1}{\sqrt{3}}$ b. $\frac{2}{\sqrt{3}}$

c. $\frac{2}{\sqrt{6}}$

d. $\frac{1}{\sqrt{6}}$

The projection of \vec{a} on \vec{b} is $|\vec{a}| \cos \theta = \frac{|\vec{a} \cdot \vec{b}|}{|\vec{b}|}$

17. Equation of a plane parallel to xz plane at unit distance is
 a. $x=1$ b. $y=1$
 c. $xz=1$ d. $xz=1$

A plane parallel to the x-z-plane has equation $y = d$, where d = distance of a plane from xz plane.

18. If the direction ratios of the line are $-1, 2, 2$ then its direction cosines are
 a. $-\frac{1}{3}, \frac{2}{3}, \frac{2}{3}$ b. $\frac{2}{3}, \frac{1}{3}, -\frac{2}{3}$
 c. $\frac{1}{3}, -\frac{2}{3}, -\frac{2}{3}$ d. none

If a, b, c are the direction ratio & l, m, n are the direction cosines. Then

$$\frac{l}{a} = \frac{m}{b} = \frac{n}{c} = \frac{1}{\sqrt{a^2 + b^2 + c^2}}$$

Given, $a, b, c = -1, 2, 2$. Then l, m, n
 $= -\frac{1}{3}, \frac{2}{3}, \frac{2}{3}$

19. The equation of the st. line passing through the point $(-1, 3)$ and x-intercept is thrice the y-intercept is
 a. $x + 3y = 8$ b. $3x - y = 4$
 c. $2x + 3y = 7$ d. $x + y = 12$

$$\frac{x}{3a} + \frac{y}{a} = 1 \dots (i) \Rightarrow -\frac{1}{3a} + \frac{3}{a} = 1$$

[passes through $(-1, 3)$]

$$\therefore a = \frac{8}{3} \text{ (put in (i))}$$

$$\therefore x + 3y = 8$$

20. The eccentricity of parabola $(y-k)^2 = 4a(x-h)$ is
 a. $e = a$ b. $e = 1$
 c. $e = \frac{1}{a}$ d. $e = \sqrt{1 + a^2}$

Parabola has eccentricity $e=1$.

21. The phonemic transcription of the word 'piety' is _____
 a. /paɪtɪ/

b. /paɪtɪ/

c. /pɪtɪ/

d. /'paɪəti/

22. The word 'Influential' has its primary stress on the _____ syllable.
 a. 1st b. 2nd
 c. 3rd d. 4th

Primary syllable stress: in-flu-**en**-tial

23. The synonym of the word 'brevity' is _____
 a. **conciseness** b. sharpness
 c. quickness d. confidence
24. The antonym of the word 'serene' is _____.
 a. complicated b. showy
 c. impure d. **agitated**
25. Which is the wrong sentences?
 a. Never does she arrive in time
 b. Black as she is, she is beautiful.
 c. Will she come to school supposing it rains heavily?
 d. **If he had left home early, he would be with us now.**

26. I am considering _____ home tomorrow.
 a. to go b. to have gone
 c. to going d. **going**
27. Like humans, zoo animals must have dentists _____ their teeth.
 a. **fill** b. filled
 c. filling d. to be filled

Here, to fill (Inf. with to) is the right usage.

28. Were you disappointed _____ your examination result?
 a. by b. from
 c. **with** d. at

Disappointed (Verb) agrees with Prep. With
 Here, with is the right usage.

29. The correct passive of "He let them scold him" is _____.
 a. He let himself to be scolded.
 b. **He let himself be scolded.**
 c. They were let to scold him.
 d. He let himself to have been scolded.

30. After eating lunch at The Cheesecake Factory, Tim went to the gym to exercise. It is a _____ sentences.

- a. simple **b. complex**
c. compound d. mixed

31. 'We congratulated him on his success' is based on the pattern_____.

- a. **S + V_t + O_i + prep. + O_D**
b. S + V_t + O_i + prepositional phrase
c. S + V_i + O_D + Prepositional phrase
d. S + V_i + O_i + prep. + O_D

32. He did nothing but _____.

- a. **play** b. to play
c. played d. playing

33. A passenger in a moving bus tosses a coin. If the coin falls behind him the bus must be moving with

- a. **acceleration**
b. deceleration
c. uniform speed
d. uniform velocity

as acceleration → falls behind him.

a deceleration → falls in front of him.

a uniform speed → falls on his hand.


34. Two bodies of masses m_1 & m_2 have equal KE. Their momenta shall in the ratio of

- a) $m_1:m_2$ b) $m_2:m_1$
c) $\sqrt{m_1}:\sqrt{m_2}$ d) $\sqrt{m_2}:\sqrt{m_1}$

$$\frac{1}{2} m_1 v_1^2 = \frac{1}{2} m_2 v_2^2 \Rightarrow \frac{(m_1 v_1)^2}{(m_2 v_2)^2} = \frac{m_1}{m_2} \therefore P_1:P_2 = \sqrt{m_1}:\sqrt{m_2}$$

35. When a drop splits up into number of drops

- a. area decreases b. volume increases
c. energy absorbed d. energy is liberated



36. The function which doesn't represent SHM is

- a. **$x = \sin^2 \omega t$**
b. $x = \cos 2\omega t$
c. $x = \sin \omega t - \cos \omega t$
d. $x = \sin \omega t + 2\cos 2\omega t$

For SHM

$$a \propto -x$$

37. The inverse square law of intensity is valid for

- a. plane wave front
b. cylindrical wave front
c. spherical wave front
d. conical wave front

38. A current I is flowing through a resistance connected to a source of small internal resistance. The resistance is reduced to half, The current through resistance will be

- a. I b. $2I$
c. $<2I$ d. $>2I$

$$I = E / (R + r), I' = E / (R/2 + r) = \frac{2E}{R + 2r}$$

$$= 2 \left(\frac{E}{R + 2r} \right)$$

$$\therefore I' < 2I$$

39. A hollow metal sphere of radius 10cm is charged such that the potential on its surface is 80V. The potential at centre of sphere is

- a. 0 **b. 80V**
c. 800V d. 8V

The potential is same as on its surface.

40. The magnification of plane mirror is

- a. 0 b. 1
c. -1 d. >1

$v=u$ but image is virtual so, $m = -1$

41. Mean free path of gas atom at a temp

- a. directly proportional to square of pressure
b. inversely proportional to square of pressure
c. inversely proportional to pressure
d. directly proportional to pressure.

Average distance travelled by a gas molecule between two successive collisions is mean free path(λ) and is given by :

$$\lambda = \frac{KT}{\sqrt{2} \pi d^2 p} \Rightarrow \lambda \propto \frac{1}{p}$$

42. Transverse elastic waves can propagate

- a) both in a gas and a metal
b) in a gas but not in a metal
c) in a metal but not in a gas
d) neither in a gas nor in a metal

43. Equal charges are given to two surfaces of different radii. The potential will be
- more on smaller sphere
 - more on bigger sphere
 - equal on both the spheres
 - dependent on nature of material of sphere

$$V = \frac{q^2}{4\pi\epsilon_0 r} \Rightarrow V \propto \frac{1}{r}$$

44. As the temperature of a conductor increases, its resistivity and conductivity changes. The ratio of resistivity to conductivity
- increases
 - decreases
 - remain same
 - may increase or decrease depending on actual temperature

Resistivity and conductivity depend only on material of conductor and temperature
 Resistivity (ρ) : Conductivity (σ) = $\rho^2 : 1$
 $T \uparrow, \rho \uparrow$

45. An electric charge in uniform motion produces
- an electric field only
 - a magnetic field only
 - both electric & magnetic field
 - no such field at all

When a charged particle is at rest then it only creates an electric field around it. When a charged particle is moving with a constant velocity creates both electric fields as well as the magnetic field around it.

46. When light waves interfere then the quantity that conserved is
- intensity
 - energy
 - mass
 - amplitude
47. Wavelength of matter wave is independent of
- mass
 - velocity
 - momentum
 - charge

$$\frac{1}{2}mv^2 = \frac{hc}{\lambda}$$

48. When we apply reverse bias to a junction diode, it
- lower the potential barrier

- raises the potential barrier
- increases the majority carrier current
- increases the minority carrier current

During reverse biasing condition majority charge carriers (Holes in p-type and electrons in N-type) are pulled away from junction. So, depletion layer increases and hence the potential barrier increases.

49. The largest wavelength of particle moving with same velocity will be for
- α -particle
 - proton
 - β -particle
 - neutron

$$m_\beta < m_p < m_n < m_\alpha \Rightarrow \lambda_\beta > \lambda_p > \lambda_n > \lambda_\alpha$$

50. The impurity atom which should be doped in pure silicon to make p type of semiconductor is
- phosphorus
 - antimony
 - boron
 - arsenic

BAG (Boron, Aluminium, Gallium) is used to make p type of semiconductor.

51. The equivalent weight of an element is 4. Its chloride has a vapor density 59.25. Then the valency of the element is
- 4
 - 3
 - 2
 - 1

Valency of metal

$$= \frac{\text{Mol. wt. of metal chloride}}{\text{Eq. wt. of metal} + 35.5} = \frac{2 \times 59.25}{4 + 35.5} = 3 \quad [\text{Mol. wt} = 2 \times \text{V.D.}]$$

52. The maximum number of electrons present in a 3d orbital is
- 2
 - 6
 - 10
 - 14

3d orbital \rightarrow 2 and 3d-subshell \rightarrow 10


53. Which of this is non-polar molecule with polar covalent bond?
- SO₂
 - NH₃
 - CO
 - BeCl₂

For a non-polar molecule with polar covalent bond;

- (i) the central atom must not have lone pair of electrons.
(ii) the surrounding atoms must be same.

54. In which of these compounds iron has least oxidation number

- a) $\text{Fe}_2(\text{SO}_4)_3$ b) $[\text{Fe}(\text{CN})_6]^{-3}$
c) **$\text{Fe}(\text{CO})_5$** d) Fe_3O_4



In case of complex compounds having neutral ligands (H_2O , NH_3 , PH_3 , CO , NO) ON is 0 due to neutral ligands.

55. HNO_3 can be stored in vessels made up of

- a) Fe b) **Al**
c) Zn d) Mg

HNO_3 makes a layer with Al and becomes passive

56. Bromine is manufactured from?

- a) Sea-weeds b) Caliche
c) **Carnallite** d) Petroleum

Carnallite, sea water \longrightarrow Bromine

57. Which of this is incorrect match of ore and metal?

- a) Calamine - Zn b) Cinnabar - Hg
c) Siderite - Fe d) **Bauxite - Cu**

Copper \longrightarrow Malachite ($\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2$),
Cuperite (Cu_2O)

Aluminium \longrightarrow Bauxite ($\text{Al}_2\text{O}_3 \cdot 2\text{H}_2\text{O}$)

58. Who synthesized the first organic compound starting from elements

- a) Wohler b) **Kolbe**
c) Berzelius d) Huckel

Wohler \longrightarrow synthesized first organic compound (urea)

Kolbe \longrightarrow Synthesized first organic compound starting from elements (acetic acid)

59. In carbocation, the hybridization of carbon atom is

- a) Sp b) **Sp^2**
c) Sp^3 d) dSp^2

60. The product of electrolysis of potassium acetate is

- a) Ethyne b) Ethene
c) **Ethane** d) Methane

Electrolysis of :

- (i) Pot. acetate \longrightarrow ethane
(ii) Pot. succinate \longrightarrow ethene
(iii) Pot. maleate / fumarate \longrightarrow ethyne

SECTION -II

Select the best alternatives: [40 x 2=80]

Read the passage carefully and put the correct answer on the answer sheet given:

It is to progress in the human sciences that we must look to undo the evils which have resulted from a knowledge of the physical world hastily and superficially acquired by populations unconscious of the changes in themselves that the new knowledge has made imperative. The road to a happier world than any known in the past lies open before us if atavistic destructive passions can be kept in leash while the necessary adaptations are made. Fears are inevitable in our time, but hopes are equally rational and far more likely to bear good fruit. We must learn to think rather less of the dangers to be avoided than of the good that will lie within our grasp if we can believe in it and let it dominate our thoughts. Science, whatever unpleasant consequences it may have by the way, is in its very nature a liberator, a liberator of bondage to physical nature and in to come, a liberator from the weight of destructive passions. We are on the threshold of utter disaster or unprecedently glorious achievement. No previous age has been fought with problems so momentous; and it is to science that we must look to for a happy future.

Questions:

61. What does science liberate us from?

It liberates us from

- a. bondage to physical nature
b. **fears and destructive passions**
c. idealistic hopes of a glorious future
d. slavery to physical nature and from passions

62. To carve out a bright future, a man should

- a. **try to avoid dangers**

72. Value of the integral $\int_{\pi/6}^{\pi/2} \frac{dx}{1 - \cos 2x}$ =

a. $\frac{1}{2}$

b. $\frac{1}{\sqrt{3}}$


c. $-\frac{\sqrt{3}}{2}$

d. $\frac{\sqrt{2}}{3}$

$$\begin{aligned} \int_{\pi/2}^{\pi/6} \frac{dx}{1 - \cos 2x} &= \int_{\pi/2}^{\pi/6} \frac{dx}{2\sin^2 x} = \frac{1}{2} \int_{\pi/2}^{\pi/6} \operatorname{cosec}^2 x \\ &= \frac{1}{2} [-\cot x]_{\pi/2}^{\pi/6} = -\frac{1}{2} [\cot \pi/6 - \cot \pi/2] \\ &= -\frac{\sqrt{3}}{2} \end{aligned}$$

- a. 12.5 miles/hr b. 10 miles/hr
c. 25 miles/hr d. 5miles/hr

$$\text{Then } \frac{10}{6} = \frac{5+x}{x}$$

$$\Rightarrow x = 7.5$$


a. $4p = 9q$ b. $p^2 + q = 4$
c. $pq = 1$ d. $2p^2 = 9q$

If the function is linear then it is one to one and onto for all $\mathbb{R} \rightarrow \mathbb{R}$ and

76. $\int \frac{dx}{(2x+1)\sqrt{4x+3}} =$

a. $\frac{1}{2} \log \frac{\sqrt{4x+3}-1}{\sqrt{4x+3}+1} + c$

b. $\sin^{-1}(2x+1) - \sqrt{4x+3} + c$

c. $\log \left(\frac{\sqrt{4x+3}}{2x+1} \right) + c$

d. $\tan^{-1} \frac{2x+1}{4} - \sqrt{4x+3} + c$

$$\begin{aligned} &\therefore \int \frac{dx}{(2x+1)\sqrt{4x+3}} \\ &= \frac{1}{2} \int \frac{z \cdot dz}{\left[2\left(\frac{z^2-3}{4}\right)+1\right]z} = \int \frac{dz}{z^2-1} \\ &= \frac{1}{2} \log\left(\frac{z-1}{z+1}\right) = \frac{1}{2} \log\left(\frac{\sqrt{4x+3}-1}{\sqrt{4x+3}+1}\right) + c \end{aligned}$$

- $$ax + by + cz + d_2 = 0 \text{ is } = \frac{|d_1 - d_2|}{\sqrt{a^2 + b^2 + c^2}}$$

- a. $\frac{5}{\sqrt{2}}$ b. $\frac{10}{\sqrt{2}}$
c. $\frac{25}{\sqrt{2}}$ d. $\frac{20}{\sqrt{2}}$

$$\left| \frac{ax_1^2 + by_1^2 + 2hx_1y_1}{\sqrt{(a-b)^2 + 4h^2}} \right| \text{ sq. unit.}$$

79. The values of $\sec^2(\tan^{-1}2) + \operatorname{cosec}^2(\cot^{-1}3) =$
- a. 5 b. 10
c. 15 d. 20

$$\begin{aligned} & \sec^2(\tan^{-1}2) + \operatorname{cosec}^2(\cot^{-1}3) \\ &= 1 + \tan^2(\tan^{-1}2) + 1 + \cot^2(\cot^{-1}3) \\ &= 1 + 4 + 1 + 9 = 15 \end{aligned}$$

80. The speed of a man with water current and against water current is 15 km/hr and 5 km/hr then speed of water is
- a. 15km/hr b. 10 km/hr
c. 5 km/hr d. 2.5km/hr

$x \rightarrow$ speed of man & $y \rightarrow$ speed of water
 $x + y = 15$... (i) and $x - y = 5$ (ii)
 Solving equ. we get, $x = 10$ and $y = 5$

81. The height from the surface of earth at which acceleration due to gravity is the same at a depth of 160 km below the surface of earth is, if radius of earth is 6400 km.
- a. 40km b. 60km
c. 80km d. 120km

82. A small sphere which has a small hole in it is immersed in water to a depth of 40 cm before any water is penetrated into it. The radius of hole if the surface tension of water is $73 \times 10^{-3} \text{ Nm}^{-1}$
- a. $1.85 \times 10^{-5} \text{ m}$ b. $3.7 \times 10^{-5} \text{ m}$
c. $5.4 \times 10^{-5} \text{ m}$ d. $7.2 \times 10^{-5} \text{ m}$

83. A rubber ball of mass 10g and volume 15 cc is dipped in water to a depth of 10m. The time taken to reach the surface is
- a. 1sec b. 1.5 sec
c. 2sec d. 3 sec

$$U - mg = ma \Rightarrow a = 0.5g$$

$$\text{time } (t) = \sqrt{\frac{2h}{a}}$$

84. A room is at 20°C is heated by a heater of resistance 20Ω connected to 200 V. Heat is transmitted through window of 1 m^2 and thickness 0.2 cm. The temperature outside the room, if conductivity of glass is $0.2 \text{ cal/m } ^\circ\text{C sec}$
- a. 12.12°C b. 14.5°C

c. 15.24°C d. 18.54°C

$$\frac{v^2}{R} \times \text{time } (t) = \frac{KA(\theta_1 - \theta_2)}{\ell} \times t$$

85. The equation of motion of an object is $x = 1.2 \cos(0.5t + \pi/4)\text{m}$. The maximum acceleration is
- a. 0.05 m/s^2 b. 0.3 m/s^2
c. 2 m/s^2 d. 2.25 m/s^2

$$x = 1.2 \cos(0.5t + \pi/4)$$

Compare this with $x = a \cos(\omega t + \phi)$
 \therefore amplitude, $a = 1.2$, $\omega = 0.5$, $\text{Accel}^n \text{ max}^m$
 $= -\omega^2 a = 0.3$

86. In a single slit experiment the first minimum of red light 660 nm coincides with first maxima for the wavelength
- a. 660 nm b. 550 nm
c. 510 nm d. 440 nm

Single slit experiment \rightarrow diffraction.

$$\frac{n\lambda D}{d} = \frac{(2n+1)\lambda'D}{2d} \Rightarrow \lambda' = 440\text{nm}.$$

(minima) (maxima)

87. A uniform copper wire of mass $2.23 \times 10^{-3} \text{ kg}$ carries a current of 1A. When 1.7 V is applied across it. The length of wire of density $8.92 \times 10^3 \text{ kg/m}^3$ & resistivity $1.7 \times 10^{-8} \Omega\text{m}$ is
- a. 1.5m b. 2.5m
c. 5m d. 7.5m

$$R = \frac{\rho \ell}{A} \Rightarrow R = \frac{\rho \ell}{\text{volume}}$$

$$\therefore \frac{V}{I} = \frac{\rho \ell^2 \times \text{density}}{\text{mass}}$$

88. A circular coil of 500 turns has a radius of 2 m and carries a current of 2 A. The magnetic field at a point on axis of coil at a distance equal to radius of coil from centre is
- a. $1.1 \times 10^{-3} \text{ T}$ b. $1.1 \times 10^{-4} \text{ T}$
c. $1.1 \times 10^{-5} \text{ T}$ d. $1.1 \times 10^{-6} \text{ T}$

$$B = \frac{\mu_0 N I R^2}{2(R^2 + x^2)^{3/2}} \quad [x = R]$$

89. A 20 V, 750 Hz source is connected to a series combination of $R = 100\Omega$, $C = 100\mu\text{F}$ and $L =$

0.1803 H. The time in which resistance of thermal capacity 2 J/°C is heated by 10°C is

- a. 174 sec b. 274 sec
c. **348 sec** d. 478 sec

$$z = \sqrt{R^2 + (wL - 1/wC)^2} = 833.99 \text{ [R = } 100\Omega\text{]}$$

$$I = E/z = 0.024 \text{ \& H = } I^2 R t$$


90. An object is placed 0.4 m from a convex mirror and a plane mirror is placed at a distance of 0.36 m from the object. The image formed in the two mirror coincide. The focal length of convex mirror is

- a. 0.2m b. 0.3m
c. 0.4m d. **0.6m**

$$u = 0.4\text{m}$$

$$v = 0.32\text{ m}$$

$$\text{Now, } 1/f = 1/u + 1/v \Rightarrow f = \frac{uv}{u+v}$$



91. A parallel beam of light is incident normally on a plane surface absorbing 40% of the light and reflecting rest. The incident beam is 10 W, then force exerted will be

- a. $5.33 \times 10^{-7} \text{ N}$ b. **$5.33 \times 10^{-8} \text{ N}$**
c. $1.33 \times 10^{-9} \text{ N}$ d. $1.33 \times 10^{-11} \text{ N}$

$$\text{Change in momentum } (\Delta p) = P - (-0.6P)$$

$$[60\% \text{ of light is reflected}] = 1.6P$$

$$\text{Momentum}(p) = \text{Power} / c$$

$$[\therefore \lambda = \frac{h}{p} \text{ and power } \left(\frac{\text{Energy}}{t(1)} \right) = \frac{hc}{\lambda}]$$

$$\text{Force} = \frac{\Delta p}{t(1\text{sec})} = \frac{1.6P}{c} = \frac{1.6 \times 10}{c}$$

$$= \frac{16}{c} = 5.33 \times 10^{-8} \text{ N}$$

92. The wavelength of 2nd line of Balmer series is 4861 \AA , then wavelength of 1st line of same series is

- a. 4861 \AA b. 5252 \AA
c. 6262 \AA d. **6562 \AA**

$$\frac{1}{\lambda} = R \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right] \Rightarrow \frac{1}{\lambda} = R \left[\frac{1}{2^2} - \frac{1}{4^2} \right]$$

$$\therefore R = \frac{16}{3\lambda} \text{ \& } \frac{1}{\lambda'} = R \left[\frac{1}{2^2} - \frac{1}{3^2} \right]$$

$$\Rightarrow \lambda' = 6562 \text{ \AA}$$

93. A sample of radioactive material has an activity of 9×10^{12} Becquerel. The material has half life of 80 sec. The time after which activity fall to 2×10^{12} Becquerel is

- a. **173.6 sec** b. 133.4 sec
c. 95.8 sec d. 52 sec

$$A = A_0 e^{-\lambda t} \Rightarrow \frac{2 \times 10^{12}}{9 \times 10^{12}} = e^{-\frac{0.693}{80} \times t}$$

$$\log \left(\frac{2}{9} \right) = -\frac{0.693}{80} \times t \Rightarrow t = 173.6 \text{ sec.}$$

94. 2gram of silver were deposited by passing 1A current through AgNO_3 solution for 35minutes. The efficiency of the current is

- a. 78% b) 81%
c) **85%** d) 91%


$$W = zIt \Rightarrow z = \frac{108}{96500} \times \frac{x}{100} \times 1 \times 35 \times 60 \Rightarrow x = 85\%$$

95. 3.55 grams of hydrated metal carbonate $\text{M}_2\text{CO}_3 \cdot x\text{H}_2\text{O}$ was dissolved in 100cc of solution. 10cc of the solution required 8cc of 0.625N alkali for neutralization. The value of x is (At. wt. of M= 23)

- a. 2 b. 5
c. 7 d. **10**

96. The K_{sp} of CaF_2 is 3.95×10^{-11} . The concentration of F^- ion in saturated solution of CaF_2 is

- a. $2.14 \times 10^{-4} \text{ mols/litre}$
b) $6.28 \times 10^{-6} \text{ mols/litre}$
c. **$4.24 \times 10^{-4} \text{ mols/litre}$**
d) $1.25 \times 10^5 \text{ mols/ litre}$



$$\text{For } \text{CaF}_2: K_{sp} = 4S^3 \therefore S = 2.1 \times 10^{-4} \text{ mol/litr}$$

$$[\text{F}^-] = 2 \times S = 4.2 \times 10^{-4} \text{ mols/litre.}$$

- a. 1:1:1 b. 1:2:3
c. 3:2:1 d. 6:3:2

- a. P_2O_5
c. CaCl_2
- b. CaO
d. conc. H_2SO_4



100. An alkene "X" on ozonolysis gave acetone and acetaldehyde. The product formed by reacting X with HBr in presence of peroxide is
- 2-bromo -2 -methyl butane
 - 2 -bromo - 3 - methyl butane
 - 2 - bromo -2 - methyl propane
 - 2- bromo pentane

$$\begin{array}{c} \text{H}_3\text{C} \quad \text{CH}_3 \\ \diagdown \quad \diagup \\ \text{C} = \text{C} \\ \diagup \quad \diagdown \\ \text{H}_3\text{C} \quad \text{H} \end{array}$$
