## EE3900 Gate Assignment - 4

## Adhvik Mani Sai Murarisetty - AI20BTECH11015

Download latex-tikz codes from

https://github.com/adhvik24/EE3900/blob/main/ Gate\_A4/main.tex

Download python codes from

https://github.com/adhvik24/EE3900/blob/main/ Gate\_A4/codes/plot.py


## Problem

(GATE EC-1997 Qn 5.2) If the fourier transform of the deterministic signal x(t) is X(f), then

- 1) The fourier transform of x(t-2) is
  - a)  $X(f)e^{-j4\pi f}$
  - b) X(2f)
  - c) 2X(2f)
  - d) X(f 2)
- 2) The fourier transform of  $x(\frac{t}{2})$  is
  - a)  $X(f)e^{-j4\pi f}$
  - b) X(2f)
  - c) 2X(2f)
  - d) X(f-2)

## Solution

1) By the time shifting property of Fourier transform,



$$x(t-2) \stackrel{\mathcal{F}}{\rightleftharpoons} e^{-j4\pi f} X(f)$$
 (0.0.2)

Let x(t) = rect(t) and  $\text{sinc}(t) = \frac{\sin(t)}{t}$ 

$$X(f) = \operatorname{sinc}(f) \tag{0.0.3}$$

For x(t) = rect(t-2),

$$e^{-j4\pi f}X(f) = e^{-j4\pi f}\operatorname{sinc}(f)$$
 (0.0.4)

The correct option is (a).




Fig. 1: Plot of rect(t-2)




Fig. 2: Fourier transform Simulation v/s Theoretical




Fig. 3: Amplitude and phase v/s frequency plot

2) By the time scaling property of Fourier transform,

$$x(\alpha t) \stackrel{\mathcal{F}}{\rightleftharpoons} \frac{1}{|\alpha|} X\left(\frac{f}{|\alpha|}\right)$$
 (0.0.5)

$$x(\frac{t}{2}) \stackrel{\mathcal{F}}{\rightleftharpoons} 2X(2f)$$
 (0.0.6)

Let x(t) = rect(t) and  $\text{sinc}(t) = \frac{\sin(t)}{t}$ 

$$X(f) = \operatorname{sinc}(f) \tag{0.0.7}$$

For  $x(t) = \text{rect}\left(\frac{t}{2}\right)$ ,

$$2X(2f) = 2\operatorname{sinc}(2f)$$
 (0.0.8)

The correct option is (c).




Fig. 4: Plot of  $rect(\frac{t}{2})$ 




Fig. 5: Fourier transform Simulation v/s Theoretical