

CSCI 347 Data Mining

Graph Data

Graph Data


- Data instances are often not entirely independent
- they can be interconnected through various types of relationships.
- Graph data or networks are a data structure where instances are depicted as **nodes**, and the connections between these instances are represented by **edges**.

Graph Data

$$G = (V, E)$$


$V = \text{set of vertices}$

$E \subseteq V \times V$, is the set of vertices in the graph




Graph Data (Weighted graph)

- $G = (V, E)$
- V = Vertices or Nodes
- E = Unordered pairs of vertices with different weights (w_{ij})




Graph Data (Directed Graph)

- $G = (V, E)$
- V = Vertices or Nodes
- E = **ordered** pairs of vertices.




Graph Data

- $G = (V, E)$
- $V = \text{Vertices or Nodes}$
- $E = \text{Unordered pairs of vertices}$
- Simple graph = Undirected graph without loops
- Edge, $e = (v_i, v_j)$, v_i and v_j are adjacent or neighbors.
- Order: $|V| = n$, Size: $|E| = m$




Graph Data

- $G = (V, E)$
- $V = \text{Vertices or Nodes}$
- $E = \text{Unordered pairs of vertices}$
- **Simple graph** = Undirected graph without loops
- **Edge**, $e = (v_i, v_j)$, v_i and v_j are adjacent or neighbors.
- **Order**: $|V| = n$, **Size**: $|E| = m$
- A graph $H = (V_H, E_H)$ is called a subgraph of $G = (V, E)$, if $V_H \subseteq V$ and $E_H \subseteq E$.



Degree of a node

- The degree of a node $v_i \in V$ is the number of edges incident with it and is denoted as $d(v_i)$ or just d_i .




Degree of a node

- The degree of a node $v_i \in V$ is the number of edges incident with it and is denoted as $d(v_i)$ or just d_i .

What is the degree of x_9 ?

1. 3
2. 1
3. 4
4. 8



Degree Distribution


- Let N_k denote the number of vertices with degree k . The degree frequency distribution of a graph is given as (N_0, N_1, \dots, N_t)
 - t is the maximum degree of a node in the graph.

Degree Distribution

- Let N_k denote the number of vertices with degree k . The degree frequency distribution of a graph is given as (N_0, N_1, \dots, N_t)
 - t is the maximum degree of a node in the graph.

What is the degree distribution of this graph?

1. (4, 3, 4, 3, 4)
2. (3, 3, 4, 4, 4)
3. (0, 0, 0, 2, 3)
4. (2, 3)



Degree Distribution

- The probability that a given node is of degree k is $\frac{N_k}{n}$.
- Suppose you have a random process of picking a node in a graph, and random variable X that assigns the degree of the picked node.


$$P(X = k) = \frac{N_k}{n}$$

Degree Distribution

- The probability that a given node is of degree k is $\frac{N_k}{n}$.
- Suppose you have a random process of picking a node in a graph, and random variable X that assigns the degree of the picked node.


$$P(X = k) = \frac{N_k}{n}$$

- Given the node distribution (0,0,0,2,3) what is the probability that a node is of degree 3?
 1. 0
 2. 2
 3. 3/5
 4. 3
 5. 2/5
 6. None of the above



Walk, Path, shortest path

- A **walk** in a graph G between nodes x and y is an ordered sequence of vertices, starting at x and ending at y .
 $Walk := \langle v_0, v_1, \dots, v_t \rangle, v_0 = x, v_t = y, \forall i \in [0..t-1]: (v_i, v_{i+1}) \text{ exists}$
- The **length of the walk** t , is the number of edges along the walk.
- A **path** is a walk with distinct vertices.
- A path of minimum length between nodes x and y is called a **shortest path**.




Walk, Path, shortest path

- A path of minimum length between nodes x and y is called a **shortest path**.

What is the length of the shortest path between x_2 and x_{10} ?

1. 6
2. 3
3. 4
4. 1
5. 0




Connectedness

- Two nodes v_i and v_j are said to be **connected** if there exists a **path** between them.
- A graph is **connected** if there is a path between all **pairs of vertices**.
- A **connected component**, or just **component**, of a graph is a **maximal connected subgraph**.
 - **maximal** means that the subgraph cannot be extended any further while still maintaining the property of being connected.



Connectedness

- Is this graph connected?
 - Yes
 - No



Connectedness

- Is this graph connected?
 - Yes
 - No



Adjacency Matrix


- A graph $G = (V, E)$, with $|V| = n$ vertices, can be conveniently represented in the form of an $n \times n$, symmetric binary adjacency matrix, A , defined as:

$$A(i, j) = \begin{cases} 1 & \text{if } v_i \text{ is adjacent to } v_j \\ 0 & \text{otherwise} \end{cases}$$

- A weighted graph can be represented by $n \times n$ weighted adjacency matrix.


$$A(i, j) = \begin{cases} w_{ij} & \text{if } v_i \text{ is adjacent to } v_j \\ 0 & \text{otherwise} \end{cases}$$

Adjacency matrix example.



| | x_1 | x_2 | x_3 | x_4 | x_5 |
|-------|-------|-------|-------|-------|-------|
| x_1 | 0 | 1 | 1 | 1 | 1 |
| x_2 | 1 | 0 | 1 | 0 | 1 |
| x_3 | 1 | 1 | 0 | 1 | 1 |
| x_4 | 1 | 0 | 1 | 0 | 1 |
| x_5 | 1 | 1 | 1 | 1 | 0 |


Adjacency matrix (weighted) example.



| | x_1 | x_2 | x_3 | x_4 | x_5 |
|-------|-------|-------|-------|-------|-------|
| x_1 | 0 | 7 | 9 | 1 | 3 |
| x_2 | 7 | 0 | 3 | 0 | 10 |
| x_3 | 9 | 3 | 0 | 3 | 4 |
| x_4 | 1 | 7 | 5 | 0 | 2 |
| x_5 | 3 | 10 | 4 | 2 | 0 |

Adjacency matrix: directed graph


- In a **directed graph** adjacency matrix is **not symmetric**.



| | x_1 | x_2 | x_3 | x_4 | x_5 |
|-------|-------|-------|-------|-------|-------|
| x_1 | 0 | 0 | 1 | 0 | 0 |
| x_2 | 1 | 0 | 1 | 0 | 1 |
| x_3 | 0 | 0 | 0 | 1 | 1 |
| x_4 | 1 | 0 | 0 | 0 | 1 |
| x_5 | 1 | 0 | 0 | 0 | 0 |

Graphs from Data Matrix


- Given a dataset in the form of a matrix, can we create a graph?



But what can we do about the edges?

Graphs from Data Matrix

- Given a dataset in the form of a matrix, can we create a graph?



**How about we using a similarity measure
and then use the similarity measure as the
edge weights?**

How to create a graph from matrix?

- Define a weighted graph $G = (V, E)$.

$$V = \{v_i \mid v_i \text{ represents the entity } x_i\}$$

$$w_{ij} = \frac{\text{sim}(x_i, x_j)}{\text{sim}(x_i, x_j)}$$

represents the similarity between points x_i and x_j

Gaussian similarity

$$w_{ij} = \text{sim}(x_i, x_j) = e^{-\frac{\|x_i - x_j\|^2}{2\sigma^2}}$$

σ is the spread parameter.


Gaussian similarity

- Similarity is defined as being inversely related to the Euclidean distance.
- If two vectors are far apart, then we say it's less similar.
 - Therefore, we can put lower weight between them.
- But why do we use this?
 - We can use something like $\frac{1}{\|x_i - x_j\|}$

Gaussian similarity

- Exponential Decay
 - The similarity measure w_{ij} decays smoothly and asymptotically to 0 as the distance increases, ensuring that distant points contribute very little but not abruptly.
 - It is bounded between 0 and 1, which simplifies interpretation and normalization in algorithms.
- Inverse distance
 - $\frac{1}{\|x_i - x_j\|}$ decays too slowly as the distance increases, leading to non-negligible contributions from distant points.
 - It has an unbounded range $(0, \infty)$, which can create numerical instability and make it harder to interpret.

Graph for $1/|x|$, $e^{-|x|^2/10}$



Feedback

Gaussian similarity

- Handling zero distance:
 - When handling $\|x_i - x_j\| = 0$, w_{ij} simplifies to $e^0 = 1$, but if we use $\frac{1}{\|x_i - x_j\|}$, then mathematically this is not defined.
- Sensitivity control
 - The parameter σ allows you to control the sensitivity to distance.
 - Smaller σ , similarity decays quickly.
 - Larger σ , similarity decays slowly.
 - We can tweak the graph by changing this parameter.
 - $\frac{1}{\|x_i - x_j\|}$ does not have this property.

Gaussian similarity

- Robustness to outliers.
 - Gaussian similarity drops off quickly for large distances, effectively ignoring the outliers.
 - $\frac{1}{\|x_i - x_j\|}$, even if far away, can have disproportionately large effect to slow decay of $\frac{1}{d}$.

Graphs from Data Matrix

- Gaussian similarity with $\sigma = 25$

| | X_1 | X_2 | X_3 |
|-------|-------|-------|-------|
| x_1 | 0.2 | 1 | 12.3 |
| x_2 | 1.3 | 4 | 89.23 |
| x_3 | 5.6 | 5 | 56.1 |
| x_4 | 4.5 | 7 | 47.3 |
| x_5 | 7.3 | 12 | 45.23 |

| | | | | |
|----------|----------|----------|----------|----------|
| 0 | 0.008709 | 0.207862 | 0.359302 | 0.366178 |
| 0.008709 | 0 | 0.409967 | 0.241611 | 0.196716 |
| 0.207862 | 0.409967 | 0 | 0.936019 | 0.87281 |
| 0.359302 | 0.241611 | 0.936019 | 0 | 0.970737 |
| 0.366178 | 0.196716 | 0.87281 | 0.970737 | 0 |

Graphs from Data Matrix

- Gaussian similarity with $\sigma = 25$

| | X_1 | X_2 | X_3 |
|-------|-------|-------|-------|
| x_1 | 0.2 | 1 | 12.3 |
| x_2 | 1.3 | 4 | 89.23 |
| x_3 | 5.6 | 5 | 56.1 |
| x_4 | 4.5 | 7 | 47.3 |
| x_5 | 7.3 | 12 | 45.23 |

| | x1 | x2 | x3 | x4 | x5 |
|----|----------|----------|----------|----------|----------|
| x1 | 0 | 0.008709 | 0.207862 | 0.359302 | 0.366178 |
| x2 | 0.008709 | 0 | 0.409967 | 0.241611 | 0.196716 |
| x3 | 0.207862 | 0.409967 | 0 | 0.936019 | 0.87281 |
| x4 | 0.359302 | 0.241611 | 0.936019 | 0 | 0.970737 |
| x5 | 0.366178 | 0.196716 | 0.87281 | 0.970737 | 0 |

Graphs from Data Matrix

- Gaussian similarity with $\sigma = 50$

| | X_1 | X_2 | X_3 |
|-------|-------|-------|-------|
| x_1 | 0.2 | 1 | 12.3 |
| x_2 | 1.3 | 4 | 89.23 |
| x_3 | 5.6 | 5 | 56.1 |
| x_4 | 4.5 | 7 | 47.3 |
| x_5 | 7.3 | 12 | 45.23 |

| | x1 | x2 | x3 | x4 | x5 |
|----|----------|----------|----------|----------|----------|
| x1 | 0 | 0.30549 | 0.675218 | 0.774221 | 0.777899 |
| x2 | 0.30549 | 0 | 0.800179 | 0.701098 | 0.665978 |
| x3 | 0.675218 | 0.800179 | 0 | 0.983606 | 0.966562 |
| x4 | 0.774221 | 0.701098 | 0.983606 | 0 | 0.992603 |
| x5 | 0.777899 | 0.665978 | 0.966562 | 0.992603 | 0 |


Creating a graph from matrix

$$\tau = 0.94$$

| | X_1 | X_2 | X_3 |
|-------|-------|-------|-------|
| x_1 | 0.2 | 1 | 12.3 |
| x_2 | 1.3 | 4 | 89.23 |
| x_3 | 5.6 | 5 | 56.1 |
| x_4 | 4.5 | 7 | 47.3 |
| x_5 | 7.3 | 12 | 45.23 |

| | x1 | x2 | x3 | x4 | x5 |
|----|----------|----------|----------|----------|----------|
| x1 | 0 | 0.30549 | 0.675218 | 0.774221 | 0.777899 |
| x2 | 0.30549 | 0 | 0.800179 | 0.701098 | 0.665978 |
| x3 | 0.675218 | 0.800179 | 0 | 0.983606 | 0.966562 |
| x4 | 0.774221 | 0.701098 | 0.983606 | 0 | 0.992603 |
| x5 | 0.777899 | 0.665978 | 0.966562 | 0.992603 | 0 |

| | x1 | x2 | x3 | x4 | x5 |
|----|----|----|----|----|----|
| x1 | 0 | 0 | 0 | 0 | 0 |
| x2 | 0 | 0 | 0 | 0 | 0 |
| x3 | 0 | 0 | 0 | 1 | 1 |
| x4 | 0 | 0 | 1 | 0 | 1 |
| x5 | 0 | 0 | 1 | 1 | 0 |



$$\text{sim}(x_i, x_j) = e^{-\frac{\|x_i - x_j\|^2}{2\sigma^2}}$$

$$A(i, j) = \begin{cases} 1 & \text{if } \text{sim}(x_i, x_j) \geq \tau \\ 0 & \text{otherwise} \end{cases}$$

Iris Similarity Graph: Gaussian Similarity

$\sigma = \frac{1}{\sqrt{2}}$, edge exist if and only if $w_{ij} \geq 0.777$
Order: $|V| = n = 150$, size: $|E| = m = 753$




Figure 4.2: Iris Similarity Graph

Topological graph attributes

- **Topological graph attributes** refer to properties of a graph that describe its structure and connectivity without considering specific geometric or spatial embedding.
- We say a graph attribute is:
 - **Local** if they apply only to a single node.
 - **Global** if they refer to the entire graph.

- Degree

- The degree of a node $v_i \in G$ is defined as:

$$d_i = \sum_j A_{ij}$$

- Clearly, this is a local attribute.
 - The corresponding global attribute for the entire graph G is the average degree:

$$\mu_d = \frac{\sum_i d_i}{n}$$

- We can generalize this to weighted and directed graphs as well.
 - The in-degree of a node $v_i \in G$ is defined as:

$$id(v_i) = \sum_j A(j, i)$$
$$od(v_i) = \sum_j A(i, j)$$

- The average indegree and average outdegree can be obtained by summing them up and dividing by n .

$$\mu_{indeg} = \frac{\sum_i id(v_i)}{n}, \mu_{outdeg} = \frac{\sum_i od(v_i)}{n}$$

Average path length

- The average path length, also called the characteristic path length, of a connected graph is given as:

$$\mu_L = \frac{\sum_i \sum_{j>i} d(v_i, v_j)}{\binom{n}{2}} = \frac{2}{n(n-1)} \sum_i \sum_{j>i} d(v_i, v_j)$$

- For directed graphs,

$$\mu_L = \frac{1}{n(n-1)} \sum_i \sum_{j>i} d(v_i, v_j)$$

For disconnected graphs, the average is taken over only the connected pairs of vertices.

Eccentricity


- The eccentricity of a node v_i is the maximum distance from v_i to any other node in the graph:

$$e(v_i) = \max_j \{d(v_i, v_j)\}$$

- The less eccentric a node is more central it is in the graph.


Example:

- What is the eccentricity of x_3 ?
 - A. 2
 - B. 3
 - C. 4
 - D. 5



Example:

- What is the eccentricity of x_6 ?
 - A. 2
 - B. 3
 - C. 4
 - D. 5



Radius and Diameter of a graph G

- The **radius** of a connected graph, denoted $r(G)$, is the minimum eccentricity of any node in the graph.

$$r(G) = \min_i \{e(v_i)\} = \min_i \left\{ \max_j \{d(v_i, v_j)\} \right\}$$


- The **diameter**, denoted $d(G)$, is the maximum eccentricity of any vertex in the graph

$$d(G) = \max_i \{e(v_i)\} = \max_{ij} \{d(v_i, v_j)\}$$

- For a disconnected graph, values are computed over the connected components of the graph.
- The diameter of a graph G is sensitive to outliers.

- What is the **radius** and **diameter** of the following graph?

- 2
- 3
- 4
- 5




- What is the **radius** and **diameter** of the following graph?

1. 2

2. 3

3. 4

4. 5




Clustering coefficient of a node

- The clustering coefficient of a node v_i is a measure of the density of edges in the neighborhood of v_i .
- Let $G_i = (V_i, E_i)$ be the subgraph induced by the neighbors of vertex v_i . Note that $v_i \notin V_i$, since we assume that G is simple. Let $|V_i| = n_i$ be the number of neighbors of v_i , and $|E_i| = m_i$ be the number of edges among the neighbors of v_i . The clustering coefficient of v_i is defined as:

$$C(v_i) = \frac{\text{\# of edges in } G_i}{\text{maximum number of edges in } G_i} = \frac{m_i}{\binom{n_i}{2}} = \frac{2 \cdot m_i}{n_i(n_i - 1)}$$


What is the clustering coefficient of v_3 ?

1. 1
2. $1/3$
3. $1/2$
4. $1/6$
5. $6/10$




What is the clustering coefficient of v_5 ?

1. 1
2. $1/3$
3. $1/2$
4. $1/6$
5. $6/10$



What is the clustering coefficient of v_6 ?

1. 1
2. $1/3$
3. $1/2$
4. $1/6$
5. $6/10$
6. Not defined



Clustering coefficient of a node

- Note: Clustering coefficient of a node is not defined for nodes with degree less than 2.
- Therefore, if we need, we can consider 0 for the clustering coefficient for nodes with degree less than 2.

Clustering coefficient of a node

- How to interpret this value?
 - $C(v_i) = 1 \rightarrow$ All neighbors are connected to each other.
 - $C(v_i) = 0 \rightarrow$ None of the neighbors are connected to each other.
 - A higher clustering coefficient indicates a more tightly knit local community around the node.
- Example:
 - Fraud Detection in Financial Networks
 - In a bank transaction network, accounts are nodes, and transactions form edges.
 - Fraud rings often have a high clustering coefficient since fraudulent accounts transfer money within a small, well-connected group.

Clustering coefficient of a graph


- The clustering coefficient of a graph G is simply the average clustering coefficient over all the nodes, given as:

$$C(G) = \frac{1}{n} \cdot \sum_i C(v_i)$$

Clustering coefficient of a graph


- The clustering coefficient of a graph G is simply the average clustering coefficient over all the nodes, given as:

$$C(G) = \frac{1}{n} \cdot \sum_i C(v_i)$$



What is the clustering coefficient of this graph?

1. $3/8$
2. $3/16$
3. $5/16$
4. $1/2$




$$C(G) = \frac{1}{8} \cdot \left(\frac{3}{\binom{4}{2}} + \frac{1}{\binom{3}{2}} + 1 + \frac{2}{\binom{4}{2}} + \frac{2}{\binom{4}{2}} + 0 + 0 + 0 \right) = \frac{2.5}{8} = \frac{5}{16}$$

Clustering coefficient of a graph

- The clustering coefficient of a graph G is simply the average clustering coefficient over all the nodes, given as:

$$C(G) = \frac{1}{n} \cdot \sum_i C(v_i)$$



What is the clustering coefficient of this graph?

1. $3/8$
2. $3/16$
3. $5/16$
4. $1/2$

Efficiency

- The efficiency for a pair of nodes v_i and v_j is defined as $\frac{1}{d(v_i, v_j)}$
- If v_i and v_j are not connected, then efficiency between these two vertices is $\frac{1}{\infty} \approx 0$.
- Smaller the distance between two nodes, these nodes are efficient in communicating between them.
- The efficiency of a graph is defined as G is defined as the average efficiency of over all pairs of nodes.

$$\frac{2}{n \cdot (n - 1)} \cdot \sum_i \sum_{j>i} \frac{1}{d(v_i, v_j)}$$

Measure of centrality

- The notion of centrality is used to rank the vertices of a graph in terms of how “**central**” or important they are.

- A centrality can be formally defined as a function:

$$c: V \rightarrow \mathbb{R}$$


that induces a total order on V

- We say that v_i is at least as central as v_j if $c(v_i) \geq c(v_j)$.
- A set has a total order if it has a partial order and every pair of elements in the set are comparable.

Degree centrality

- The simplest notion of centrality is the degree d_i of a vertex v_i .
- the higher the degree the more important or central the vertex.
- For directed graphs, one may further consider the indegree centrality and outdegree centrality of a vertex.


Degree centrality



$$d(v_i) = \sum_j^n A(i, j)$$

$$d(x_3) = \sum_1^n A(3, j) = 4$$


Degree centrality



$$d(v_i) = \sum_j^n A(i, j)$$

$$d(x_5) = \sum_1^n A(5, j) = 1$$

Degree centrality



$$d(x_3) = \sum_{j=1}^n A(3, j) = 4$$

$$d(x_5) = \sum_{j=1}^n A(5, j) = 1$$


x_3 has higher importance than x_5 according to degree centrality.

Eccentricity Centrality

- Eccentricity centrality is thus defined as follows

$$c(v_i) = \frac{1}{e(v_i)} = \frac{1}{\max_j \{d(v_i, v_j)\}}$$

- The less eccentric a node is the more central it is.



What is the eccentric centrality of x_3 ?

- 1. $\frac{1}{2}$
- 2. $\frac{1}{3}$
- 3. 4
- 4. 2
- 5. 3

Closedness centrality

- The closeness centrality uses the sum of all the distances to rank how central a node is.

$$c(v_i) = \frac{1}{\sum_j d(v_i, v_j)}$$


A node v_i with the smallest total distance, $\sum_j d(v_i, v_j)$ is called the ***median node***.

- Why is this important?

Closedness centrality

- Closedness centrality tries to minimize the total distance over all the other nodes, and thus, a median node, which has the highest closedness centrality, is the optimal one to, say, locate a facility that minimizes the distance to all other points.


Closedness centrality



What is the **closedness centrality** of x_3 ?

1. $1/8$
2. $1/3$
3. 3
4. 4
5. 8

Closedness centrality



What is the **eccentricity centrality** of x_3 ?

- 1. $\frac{1}{2}$
- 2. $\frac{1}{3}$
- 3. 4
- 4. 2
- 5. 3

Betweenness centrality

- For a given vertex v_i the betweenness centrality measures how many shortest paths between all pairs of vertices include v_i .
- This gives an indication as to the central “**monitoring**” role played by v_i for various pairs of nodes.


Betweenness centrality

- Let η_{jk} denote the number of shortest paths between vertices v_j and v_k , and let
- $\eta_{jk}(v_i)$ denote the number of such paths that include or contain v_i , then the fraction of paths through v_i is denoted as


$$\gamma_{jk}(v_i) = \frac{\eta_{jk}(v_i)}{\eta_{jk}}.$$

- If the two vertices v_j and v_k are not connected, we assume $\gamma_{jk} = 0$.


$$c(v_i) = \sum_{j \neq i} \sum_{\substack{k \neq i \\ k > j}} \gamma_{jk}$$



What is the betweenness centrality of v_5 ?




| | Shortest paths | Shortest paths through v_5 $\eta_{jk}(v_5)$ |
|-------------|----------------|--|
| η_{12} | 1 | 0 |
| η_{13} | 1 | 0 |
| η_{14} | 1 | 0 |
| η_{16} | 1 | 0 |
| η_{17} | 1 | 0 |
| η_{18} | 1 | 1 |
| η_{23} | 1 | 0 |
| η_{24} | 2 | 1 |
| η_{26} | 1 | 0 |
| η_{27} | 3 | 2 |
| η_{28} | 1 | 1 |
| η_{34} | 1 | 0 |
| ... | | |



What is the betweenness centrality of v_5 ?

$$c(v_5) = \gamma_{18} + \gamma_{24} + \gamma_{27} + \gamma_{28} + \gamma_{38} + \gamma_{46} + \gamma_{48} + \gamma_{67} + \gamma_{68}$$

$$c(v_5) = 1 + \frac{1}{2} + \frac{2}{3} + 1 + \frac{2}{3} + \frac{1}{2} + \frac{1}{2} + \frac{2}{3} = 6.5$$



| Centrality | v_1 | v_2 | v_3 | v_4 | v_5 | v_6 | v_7 | v_8 |
|-----------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|
| Degree | 4 | 3 | 2 | 4 | 4 | 1 | 2 | 2 |
| Eccentricity $e(v_i)$ | 0.5 | 0.33 | 0.33 | 0.33 | 0.5 | 0.25 | 0.25 | 0.33 |
| | 2 | 3 | 3 | 3 | 2 | 4 | 4 | 3 |
| Closeness $\sum_j d(v_i, v_j)$ | 0.100 | 0.083 | 0.071 | 0.091 | 0.100 | 0.056 | 0.067 | 0.071 |
| | 10 | 12 | 14 | 11 | 10 | 18 | 15 | 14 |
| Betweenness | 4.5 | 6 | 0 | 5 | 6.5 | 0 | 0.83 | 1.17 |