

Session C3: VSEPR, Molecular Shapes, Bond Angles & Polarity

Duration: 45 minutes (strict)

Continues from: Session C2, *Drawing Lewis Structures, Covalent Molecules & Polyatomic Ions*. C2 ended with this exact next step: a Lewis structure shows **which** electrons are shared, but not the **3-D shape**. Today uses the Lewis structure to predict the shape, and then the shape to decide whether the molecule is **polar**.

Class worksheet (9/30): an 8-column table: *Chemical Formula · Valence e⁻ · Lewis · Electron Geometry · Molecular Geometry & Bond Angle · Molecular Geometry Redraw · Bond Polarity? · Molecular Polarity?* The rows are PBr₃, HCO₂⁻, NOCl, NO₂⁻ and CF₂Cl₂. Dhairya finished most of the PBr₃ and HCO₂⁻ rows in class. **Expect the test to use this same table**, so every worked example below is laid out column by column.

Grade-9 foundation: NCERT Class 9 Ch9 *Atomic Foundations of Matter*, §9.4.1 *Bonding by sharing of electrons: Covalent Bond* (valence electrons, shared pairs, the octet).

Grade-11 depth: NCERT Class 11 Ch4 *Chemical Bonding and Molecular Structure*: §4.3.6 *Polarity of Bonds* (dipole moment, the crossed arrow, bond dipoles adding as vectors, Table 4.5, pp. 110–112) and §4.4 *The VSEPR Theory* (postulates, repulsion order, Tables 4.6–4.8, pp. 112–116). Electronegativities are the Pauling values from NCERT Class 11 Ch3, the same ones used in C1.

Interactive tools (study-tools site, Chemistry): *VSEPR Shape Explorer* ([vsepr-shapes/](#)), *Worksheet Trainer* ([vsepr-table/](#)), *VSEPR & Polarity flashcards* ([vsepr-flashcards/](#)) and *VSEPR & Polarity Practice Test* ([vsepr-test/](#)).

Lesson Plan

Objectives. By the end of this session Dhairya should be able to fill in every column of the worksheet table for any molecule or ion with up to 4 electron domains:

1. Count the **electron domains** on the central atom from its Lewis structure, and name the **electron geometry**.
2. Name the **molecular geometry** and give the **bond angle**, including how lone pairs squeeze the angle.
3. **Redraw** the molecule in 3-D with plain lines, wedges and dashes.
4. Decide whether each bond is **polar** from ΔEN , and whether the whole **molecule** is polar from its shape.

Time	Segment
0–3 min	Warm-up: 3 quick recall questions
3–13 min	Part 1: VSEPR, electron domains and electron geometry (show the <i>Shape Explorer</i>)
13–20 min	Part 2: molecular geometry, bond angles and the lone-pair squeeze
20–23 min	Part 3: the 3-D redraw with wedges and dashes
23–31 min	Part 4: bond polarity and molecular polarity
31–42 min	Part 5: the worksheet: check rows 1–2 together, then he solves rows 3–5 alone
42–45 min	Exit ticket; homework is the <i>Worksheet Trainer</i> plus the practice test

Pacing note. Part 5 is the priority, since it is exactly what the test looks like. If you run behind, skip the derivations (the 109.5° cube proof and the vector sums). State the results and leave the proofs to the notes. Do not skip the “electron geometry vs molecular geometry” distinction in Part 2. It is the most common test mistake.

The row routine (one column at a time)

Column	How to fill it
Valence e⁻	Add each atom's valence electrons (the last digit of its group). Add 1 for each negative charge; subtract 1 for each positive charge.
Lewis	The C2 4-step method. For an ion, use brackets and write the charge outside.
Electron geometry	Count the electron domains on the central atom: each bonded atom counts 1 (single, double or triple bond), and each lone pair counts 1. Then 2 = linear, 3 = trigonal planar, 4 = tetrahedral.
Molecular geometry & bond angle	Use the atoms only (the lone pairs are invisible) and read the shape from the chart in Part 2.
Redraw	Draw that shape in 3-D: plain lines lie in the page, a wedge comes toward you, a dash goes away. Draw the lone pairs on the central atom.
Bond polarity	For each <i>kind</i> of bond, $\Delta EN = \text{larger EN} - \text{smaller EN}$. Below 0.5 it is nonpolar; from 0.5 to 1.7 it is polar.
Molecular polarity	If there are no polar bonds, the molecule is nonpolar. If there are polar bonds, it is nonpolar only if the shape is symmetric: no lone pairs on the central atom and identical outer atoms. Otherwise it is polar.

Warm-up (3 min)

1. *How many valence electrons does NOCl have?* $\rightarrow N\ 5 + O\ 6 + Cl\ 7 = \mathbf{18}$.
2. *In NOCl, which atom is central, and why?* $\rightarrow \mathbf{N}$. It forms 3 bonds, O forms 2 and Cl forms 1. The atom that forms the most bonds goes in the middle (C2, step 2).
3. *C1 recall: is a C-H bond polar?* $\rightarrow \Delta EN = 2.5 - 2.1 = 0.4$, which is below 0.5, so it is **nonpolar**. (This is the "H: nonpolar" note on the worksheet.)

Bridge to today: "A Lewis structure is flat. Real molecules are 3-D, and their shape decides things like whether water is polar. Today we get the shape from the Lewis structure you already know how to draw."

Part 1 — VSEPR: Electron Domains & Electron Geometry (10 min)

The idea

VSEPR stands for **V**alence **S**hell **E**lectron **P**air **R**epulsion. The main postulates, from NCERT §4.4:


- The shape of a molecule depends on the number of valence-shell electron pairs, **bonded or non-bonded**, around the **central atom**.
- Electron pairs **repel** each other, because their clouds are negatively charged.
- The pairs therefore sit **as far apart as possible**, which minimizes repulsion.
- A **multiple bond is treated as if it were a single electron pair**. The 2 or 3 pairs of a double or triple bond act as one "super pair".
- When a molecule has **resonance structures**, VSEPR applies to any one of them. They all give the same shape.

Counting electron domains


An **electron domain** is one region of electron density around the central atom:

- each **bonded atom** = 1 domain, whether the bond is single, double or triple;
- each **lone pair on the central atom** = 1 domain.


Do not count the lone pairs on the outer atoms. They affect their own atom, not the central atom's shape.




CO₂: 2 domains
2 double bonds, each counts once




CH₂O: 3 domains
1 double + 2 single bonds



SO₂: 3 domains
2 bonds + 1 lone pair




NH₃: 4 domains
3 bonds + 1 lone pair



H₂O: 4 domains
2 bonds + 2 lone pairs

Electron geometry

Electron geometry is the arrangement of **all** the domains, lone pairs included. It depends only on how many domains there are (NCERT Fig. 4.6 and Table 4.6):



Domains	Electron geometry	Angle between domains	Why that angle
2	linear	180°	Two domains get farthest apart on opposite sides.
3	trigonal planar	120°	Three domains spread around a circle: $360^\circ \div 3 = 120^\circ$, all in one plane.
4	tetrahedral	109.5°	In 3-D, four domains point to the corners of a tetrahedron (derivation below).
5, 6	trigonal bipyramidal, octahedral	90°/120°, 90°	Needs an expanded octet (period 3 and below). Not on this test.

Why 109.5°, not 90°? (Derivation.) On paper, 4 bonds look as if they would sit at 90° in a “+” shape. In 3-D they can get farther apart. Put the central atom at the centre of a cube, and the 4 domains toward 4 alternate corners:

$$a = (1, 1, 1), b = (1, -1, -1), c = (-1, 1, -1), d = (-1, -1, 1).$$

The angle θ between any two of them comes from the dot product:

$$\cos \theta = (a \cdot b) / (|a| |b|) = (1 \cdot 1 + 1 \cdot (-1) + 1 \cdot (-1)) / (\sqrt{3} \cdot \sqrt{3}) = (1 - 1 - 1) / 3 = -1/3$$

$$\theta = \cos^{-1}(-1/3) = 109.47^\circ \approx 109.5^\circ.$$






That is wider than 90°, so the tetrahedron is the arrangement that keeps the pairs farthest apart. (Each pair of corners gives the same $-1/3$, so all six angles are equal.)

Part 2 — Molecular Geometry, Bond Angles & the Lone-Pair Squeeze (7 min)

Electron geometry vs molecular geometry

- **Electron geometry** includes every domain, lone pairs too.
- **Molecular geometry** (the “shape”) describes **only where the atoms are**. Imagine the lone pairs are invisible.

When the central atom has **no lone pairs**, the two geometries are the **same**. When it has lone pairs, the molecular geometry gets a **different name**. This is NCERT’s split into “no lone pair” (Table 4.6) and “one or more lone pairs” (Table 4.7).

Electron domains (electron geometry)	0 lone pairs	1 lone pair	2 lone pairs
2 linear	 linear $AX_2 \cdot 180^\circ$ CO_2, HCN, OCS	—	—
3 trigonal planar	 trigonal planar $AX_3 \cdot 120^\circ$ BF_3, SO_3, HCO_2^-	 bent $AX_2E \cdot <120^\circ$ $SO_2, NO_2^-, NOCl$	—
4 tetrahedral	 tetrahedral $AX_4 \cdot 109.5^\circ$ CH_4, CCl_4, CF_2Cl_2	 trigonal pyramidal $AX_3E \cdot \approx 107^\circ$ NH_3, PBr_3, H_3O^+	 bent $AX_2E_2 \cdot \approx 104.5^\circ$ H_2O, OF_2, ClO_2^-

Domains	Bonded atoms	Lone pairs	Electron geometry	Molecular geometry	Bond angle	Examples
2	2	0	linear	linear	180°	CO_2, HCN
3	3	0	trigonal planar	trigonal planar	120°	BF_3, HCO_2^-
3	2	1	trigonal planar	bent	slightly $< 120^\circ$	SO_2 (119.5°), NO_2^- , $NOCl$
4	4	0	tetrahedral	tetrahedral	109.5°	CH_4, CF_2Cl_2
4	3	1	tetrahedral	trigonal pyramidal	$\approx 107^\circ$ ($< 109.5^\circ$)	NH_3, PBr_3
4	2	2	tetrahedral	bent	$\approx 104.5^\circ$ ($< 109.5^\circ$)	H_2O

(The AXE labels in the chart come from NCERT Table 4.7: A = central atom, X = bonded atom, E = lone pair. NH_3 is AX_3E .)


Watch out: “bent” appears **twice**. With 3 domains it is under 120° , and with 4 domains it is about 104.5° . On the test, write the electron geometry too, so the grader can see which bent you mean.

Lone pairs squeeze the bond angle


NCERT §4.4 gives the order of repulsion:

lone pair-lone pair > lone pair-bond pair > bond pair-bond pair


Why: a bonding pair is shared between two nuclei, so it is pulled out into a narrow region. A lone pair belongs to the central atom alone, so it spreads out and takes **more space**. It pushes the bonding pairs closer together, which makes the bond angle smaller than the ideal one (NCERT Table 4.8):



CH₄: 109.5°
4 bonding pairs, 0 lone pairs



NH₃: 107°
1 lone pair pushes the bonds closer



H₂O: 104.5°
2 lone pairs push even harder

- **CH₄**: 4 bond pairs, no squeeze, so **109.5°**.
- **NH₃**: 1 lone pair pushes on 3 bond pairs, so the angle drops to **107°**.
- **H₂O**: 2 lone pairs push harder still (lp-lp is the strongest), so the angle drops to **104.5°**.
- **SO₂**: 3 domains with 1 lone pair, so the angle drops from 120° to **119.5°**.

What to write on the test: trigonal pyramidal “ $\approx 107^\circ$ (less than 109.5°)”; bent from 4 domains “ $\approx 104.5^\circ$ (less than 109.5°)”; bent from 3 domains “less than 120°”. *Beyond the test:* real angles vary with the atoms. PBr₃ is about 101°, because big Br atoms on a large P squeeze further. The ideal-based answers above are what a VSEPR test expects.

Part 3 — The 3-D Redraw: Wedges & Dashes (3 min)

A page is flat, so chemists use three kinds of line to show depth:




Plain line
in the plane of the page



Solid wedge
comes out toward you



Hashed wedge (dash)
goes back, away from you



CH₄ drawn in 3D
2 in plane, 1 toward you, 1 away

How to draw each shape:

- **Linear, trigonal planar, bent:** everything lies in one plane, so use plain lines only.
- **Tetrahedral:** 2 bonds in the plane of the page (plain lines), 1 wedge (toward you) and 1 dash (away).
- **Trigonal pyramidal:** the lone pair points up; below it go 1 plain line, 1 wedge and 1 dash.
- **Always** draw the central atom's lone pairs in the redraw (as a lobe or two dots). They are why the shape is what it is. Outer atoms' lone pairs can be left out of the redraw.

Part 4 — Bond Polarity & Molecular Polarity (8 min)

Bond polarity (C1 recap, now with arrows)

When two different atoms share a pair, the **more electronegative** atom pulls the pair toward itself. It becomes slightly negative (δ^-) and the other atom slightly positive (δ^+). NCERT §4.3.6 calls this a **polar covalent bond**; with identical atoms (H₂, Cl₂) the pair sits in the middle and the bond is **nonpolar**.

The C1 rule for bond polarity, using $\Delta\text{EN} = \text{larger EN} - \text{smaller EN}$:

ΔEN	Bond
less than 0.5	nonpolar covalent
0.5 to 1.7	polar covalent
more than 1.7	mostly ionic (between two non-metals, call it <i>very polar</i> covalent, like H-F in C1)


Electronegativities you'll need (Pauling, NCERT Class 11 Ch3; the same values as C1):

Dipole moment (NCERT §4.3.6). A polar bond has a **dipole moment**, which measures how strongly the charge is separated:

$$\mu = Q \times r$$

where Q is the size of the separated charge and r is the distance between the centres of positive and negative charge. The unit is the **debye**: $1 \text{ D} = 3.33564 \times 10^{-30} \text{ C m}$. It is a **vector**: it has a size and a direction.

The crossed arrow (NCERT's chemistry convention): draw an arrow along the bond with a **cross (+)** at the $\delta+$ end and the **arrowhead** at the $\delta-$ end. The arrow points the way the electrons are pulled, toward the more electronegative atom.




Molecular polarity: add the arrows


For a molecule with more than one bond, NCERT §4.3.6 says the dipole moment of the molecule is the **vector sum** of the bond dipoles. So the **shape** decides the answer:

- If the arrows **cancel**, the molecule is **nonpolar**, even when every bond is polar.
- If they **don't cancel**, there is a **net dipole** and the molecule is **polar**.

Symmetric: bond dipoles cancel → nonpolar molecule



Unsymmetric: bond dipoles add up → polar molecule



Derivations (vector sums):


- **CO₂ (linear)**: the two C=O dipoles are equal and point in opposite directions: $\mu + (-\mu) = 0$. So CO₂ is **nonpolar**, although each C=O bond is polar ($\Delta \text{EN} = 1.0$). NCERT Table 4.5: CO₂ = 0 D.
- **BF₃ (trigonal planar)**: the three equal B-F dipoles point at 90°, 210° and 330°. Adding their components:
 - $x: \mu(\cos 90^\circ + \cos 210^\circ + \cos 330^\circ) = \mu(0 - 0.866 + 0.866) = 0$
 - $y: \mu(\sin 90^\circ + \sin 210^\circ + \sin 330^\circ) = \mu(1 - 0.5 - 0.5) = 0$

The net is **0**, so BF_3 is **nonpolar** (Table 4.5: 0 D). As NCERT puts it, “the resultant of any two is equal and opposite to the third”.

- **CCl_4 (tetrahedral):** using the cube corners from Part 1, the four equal dipoles add to $(1,1,1) + (1,-1,-1) + (-1,1,-1) + (-1,-1,1) = (0, 0, 0)$. CCl_4 is **nonpolar** (Table 4.5: 0 D).
- **H_2O (bent, 104.5°):** two equal O-H dipoles at an angle θ . Their components across the bisector cancel, and their components along the bisector add. Each contributes $\mu \cos(\theta/2)$, so the net is $2\mu \cos(\theta/2)$. NCERT gives the net as **1.85 D**. So $1.85 = 2\mu \cos 52.25^\circ$, which gives $\mu(\text{O-H}) = 1.85 \div (2 \times 0.612) \approx \mathbf{1.51 \text{ D}}$ for each bond. The net is not zero, so water is **polar**.
- **CHCl_3 vs CCl_4 :** swap one Cl for an H and the symmetry is broken. $\text{CHCl}_3 = \mathbf{1.04 \text{ D}}$ (polar), while $\text{CCl}_4 = 0 \text{ D}$ (Table 4.5). This is exactly the CF_2Cl_2 idea on the worksheet.

Lone pairs have their own pull too (NCERT’s NH_3 vs NF_3 example). In NH_3 the lone pair points the same way as the N-H bond dipoles, so the net is large (1.47 D). In NF_3 it points the opposite way to the N-F dipoles, so the net is small (0.23 D). Both are still **polar**, because neither is symmetric.

The quick test for molecular polarity



Symmetric shapes that can be nonpolar: linear, trigonal planar and tetrahedral, with **no lone pairs on the central atom** and **all outer atoms the same**. Any lone pair on the central atom (bent, trigonal pyramidal) or any mix of outer atoms (CHCl_3 , CF_2Cl_2 , HCN , HCO_2^-) makes the molecule **polar**. The one exception is a molecule with no polar bonds at all, like CH_4 or CS_2 : there is nothing to add up, so it is nonpolar.

Ions on the worksheet (HCO_2^- , NO_2^-): an ion carries an overall charge, so “polar” here means the same thing it does for a molecule: whether the bond dipoles cancel. Use the same flowchart.


Part 5 — The 9/30 Worksheet, Row by Row (11 min)

Plan: spend 2 minutes checking rows 1-2 with him (he did most of them in class), then he completes rows 3-5 alone with the row routine. Mark each column together.

Feedback on his rows 1-2


- **PBr_3 :** everything he wrote is right: $26 e^-$, the Lewis structure (lone pair on P, 3 on each Br), tetrahedral with 4 domains, trigonal pyramidal 107° , the wedge/dash redraw, and polar bonds. **Missing:** molecular polarity = **Polar** (it has a lone pair, so it is not symmetric).
- **HCO_2^- :** right so far: $18 e^-$, brackets and the $-$ charge, trigonal planar with $3 e^-$ domains, trigonal planar 120° , and the redraw. **Missing:** bond polarity: C-H is **nonpolar** ($\Delta\text{EN} 0.4$; his “H: nonpolar” note is right) and C-O is **polar** ($\Delta\text{EN} 1.0$). Molecular polarity is **Polar**, because the outer atoms are not all the same (1 H and 2 O).
 - Also mention: HCO_2^- has **2 resonance structures** (the double bond can sit on either O), just like NO_2^- in C2. The shape is the same in both.
- **Spelling on the test:** “trigonal planar” (not “planer”) and “pyramidal”.

Row 1 — PBr₃




1. **Valence e⁻:** P = 5, Br = $3 \times 7 = 21$, so the total is **26**.
2. **Lewis:** P is central (it forms 3 bonds; Br forms only 1). 3 P-Br bonds use 6, leaving 20. Each Br gets 3 lone pairs (18), leaving 2, which go on P as 1 lone pair. P has 3 bonds + 1 lone pair = 8 ✓, and each Br has 8 ✓. All formal charges are 0.
3. **Electron geometry:** P has 3 bonded atoms + 1 lone pair = **4 domains**, so **tetrahedral**.
4. **Molecular geometry & angle:** 3 atoms + 1 lone pair gives **trigonal pyramidal**, $\approx 107^\circ$ (less than 109.5° , squeezed by the lone pair).
5. **Redraw:** lone pair on top of P; below it, 1 plain line, 1 wedge and 1 dash to the three Br.
6. **Bond polarity:** P-Br: $\Delta EN = 2.8 - 2.1 = 0.7$, which is **polar** (Br is δ^-).
7. **Molecular polarity: polar.** There are polar bonds and a lone pair on P, so the three P-Br dipoles (all pointing down toward the Br atoms) add up instead of cancelling.

Row 2 — HCO₂⁻ (formate)



1. **Valence e⁻:** H 1 + C 4 + O 2 $\times$ 6 + 1 for the - charge = **18**.
2. **Lewis:** C is central (it forms 4 bonds; H is never central). The skeleton H-C bonded to two O uses 3 bonds = 6, leaving 12. Each O gets 3 lone pairs (12), leaving 0. C has only 3 bonds = 6 ✗. Move one O lone pair into a **C=O** double bond, and C now has 8 ✓. Draw brackets with the - outside.
 - Formal charges: C = $4 - 0 - 4 = 0$; the O in C=O = $6 - 4 - 2 = 0$; the O in C-O = $6 - 6 - 1 = -1$. The sum is -1 ✓.
 - **Resonance:** the double bond can go to either O, so there are 2 equivalent structures, joined by $\leftrightarrow$.
3. **Electron geometry:** C has 3 bonded atoms and 0 lone pairs = **3 domains** (the double bond counts once), so **trigonal planar**.
4. **Molecular geometry & angle:** no lone pairs, so it is also **trigonal planar**, **120°** .
5. **Redraw:** all in one plane: H up, the two O atoms at lower left and lower right, with one C=O drawn double.
6. **Bond polarity:** C-H: $2.5 - 2.1 = 0.4$, **nonpolar**. C-O: $3.5 - 2.5 = 1.0$, **polar** (O is δ^-).
7. **Molecular polarity: polar.** The shape has no lone pair, but the outer atoms are **not identical** (H vs O), so the two strong C-O dipoles are not balanced by the weak C-H. The net points toward the O side.


Row 3 — NOCl (nitrosyl chloride)



- Valence e⁻:** N 5 + O 6 + Cl 7 = **18**.
- Lewis:** N is central (it forms 3 bonds; O forms 2, Cl forms 1). The skeleton Cl-N-O uses 2 bonds = 4, leaving 14. Cl gets 3 lone pairs (6) and O gets 3 lone pairs (6), leaving 2, which go on N as a lone pair. N has 2 bonds + 1 lone pair = 6 $\times$. Move one lone pair **from O** into the N-O bond, giving **N=O**. N has 8 $\checkmark$, O has 2 bonds + 2 lone pairs = 8 $\checkmark$, and Cl has 8 $\checkmark$.
 - Why O and not Cl for the double bond?** With N=O every formal charge is 0: N = 5 - 2 - 3 = 0, O = 6 - 4 - 2 = 0, Cl = 7 - 6 - 1 = 0. A Cl=N double bond would put +1 on Cl and -1 on O, which is worse (C2's formal-charge rule).
 - Result:** Cl-N=O with 1 lone pair on N, 2 on O and 3 on Cl.
- Electron geometry:** N has 2 bonded atoms + 1 lone pair = **3 domains**, so **trigonal planar**.
- Molecular geometry & angle:** 2 atoms + 1 lone pair gives **bent, less than 120°** (the real angle is about 113°).
- Redraw:** in one plane: the lone pair on top of N, and Cl and O below at each side, with N=O drawn double.
- Bond polarity:** N-Cl: 3.0 - 3.0 = **0.0, nonpolar**. N=O: 3.5 - 3.0 = **0.5, polar** (right on the line; O is δ^-).
- Molecular polarity: polar.** The shape is bent (a lone pair on N) and the outer atoms are different, so the N=O dipole is not cancelled.

Note on the numbers: some periodic tables print more decimals (N 3.04, O 3.44, Cl 3.16). Then N=O is 0.40 and N-Cl is 0.12, so some answer keys call both bonds only slightly polar. The **molecular** answer is **polar** either way. Use the EN table the teacher gives on the test.


Row 4 — NO₂⁻ (nitrite)




- Valence e⁻:** N 5 + O 2 × 6 + **1** = **18**.
- Lewis** (from C2): the skeleton O-N-O uses 4, leaving 14. Each O gets 3 lone pairs (12), leaving 2, which go on N as a lone pair. N has 2 bonds + 1 lone pair = 6 $\times$. Move one O lone pair to make **N=O**. Now N, both O atoms and the total are all right: 8 $\checkmark$, 8 $\checkmark$, 18 $\checkmark$. Brackets and - .
 - Formal charges: N = 5 - 2 - 3 = 0; O in N=O = 0; O in N-O = **-1**. The sum is -1 $\checkmark$. There are **2 resonance structures**.
- Electron geometry:** N has 2 bonded atoms + 1 lone pair = **3 domains**, so **trigonal planar**.
- Molecular geometry & angle:** bent, **less than 120°** (about 115°).
- Redraw:** in one plane: the lone pair on top of N, and the two O atoms below at each side. One N=O is drawn double, but because of resonance the two bonds are really identical.
- Bond polarity:** N-O: 3.5 - 3.0 = **0.5, polar** (O is δ^-).
- Molecular polarity: polar.** It is bent, so the two N-O dipoles point partly the same way (down, toward the O atoms) and add.

Compare with row 2: HCO₂⁻ and NO₂⁻ both have 18 electrons and 2 resonance forms, yet HCO₂⁻ is **trigonal planar** and NO₂⁻ is **bent**. The difference is the **lone pair on N**. Count the domains every time; never guess the shape from the formula.

Row 5 — CF₂Cl₂ (Freon-12)



Lewis structure
all formal charges 0



3D redraw with bond dipoles
red: bond dipoles · blue: net

- Valence e⁻:** C 4 + F 2 × 7 + Cl 2 × 7 = **32**.
- Lewis:** C is central (it forms 4 bonds and is the least electronegative; F is never central). 4 single bonds use 8, leaving 24. Each of the 4 halogens gets 3 lone pairs (24), leaving 0. C has 4 bonds = 8 ✓, and every halogen has 8 ✓. All formal charges are 0. There is **no lone pair on C**.
- Electron geometry:** C has 4 bonded atoms + 0 lone pairs = **4 domains**, so **tetrahedral**.
- Molecular geometry & angle:** **tetrahedral, 109.5°** (approximately, since the atoms differ).
- Redraw:** 2 plain lines in the page (the 2 Cl), 1 wedge and 1 dash (the 2 F). In a tetrahedron all 4 corners are equivalent, so any placement is the same molecule.
- Bond polarity:** C-F: 4.0 – 2.5 = **1.5, polar**. C-Cl: 3.0 – 2.5 = **0.5, polar**. F and Cl are both δ⁻.
- Molecular polarity:** **polar**. The shape is symmetric, but the outer atoms are **not identical**: the C-F dipoles (ΔEN 1.5) are much stronger than the C-Cl dipoles (0.5), so they can't cancel.

Proof by vector sum. Put F on the cube corners (1,1,1) and (1,-1,-1), and Cl on (-1,1,-1) and (-1,-1,1). Weight each by its ΔEN:

$$1.5[(1,1,1) + (1,-1,-1)] + 0.5[(-1,1,-1) + (-1,-1,1)] = 1.5(2,0,0) + 0.5(-2,0,0) = \mathbf{(2, 0, 0) \neq 0}$$

The net points toward the side with the two F atoms. If all four were Cl (CCl₄), the same sum would be (0, 0, 0).

The completed worksheet

Chemical formula	Valence e ⁻	Lewis	Electron geometry	Molecular geometry & bond angle	Molecular geometry redraw	Bond polarity?	Molecular polarity?
PBr ₃	26		tetrahedral 4 domains	trigonal pyramidal ≈107°		P-Br: polar ΔEN 0.7	Polar
HCO ₂ ⁻	18		trigonal planar 3 domains	trigonal planar 120°		C-H: nonpolar ΔEN 0.4 C=O: polar ΔEN 1.0 C-O: polar ΔEN 1.0	Polar
NOCl	18		trigonal planar 3 domains	bent <120°		N-Cl: nonpolar ΔEN 0.0 N=O: polar ΔEN 0.5	Polar
NO ₂ ⁻	18		trigonal planar 3 domains	bent <120°		N=O: polar ΔEN 0.5 N-O: polar ΔEN 0.5	Polar
CF ₂ Cl ₂	32		tetrahedral 4 domains	tetrahedral 109.5°		C-Cl: polar ΔEN 0.5 C-F: polar ΔEN 1.5	Polar

Exit Ticket (3 min)

- How many electron domains are around C in CO₂? → **2**. Each double bond counts once.
- The electron geometry is tetrahedral and the central atom has 1 lone pair. What is the molecular geometry and bond

angle? → **Trigonal pyramidal, $\approx 107^\circ$.**

3. CCl_4 has four polar bonds. Why is it nonpolar? → It is **tetrahedral with 4 identical atoms**, so the 4 equal dipoles cancel.
 4. Which is polar, BF_3 or NF_3 ? → **NF_3** . It has a lone pair on N, so it is trigonal pyramidal and its dipoles don't cancel. BF_3 is trigonal planar and symmetric, so it is nonpolar.
-

Homework — Complete the Row (answers below)

For each one, fill in all 8 worksheet columns: valence e^- , Lewis, electron geometry (with the number of domains), molecular geometry & bond angle, the 3-D redraw, bond polarity (with ΔEN) and molecular polarity. Then do one round of the *Worksheet Trainer* and take the *VSEPR & Polarity Practice Test*.

1. CH_2Cl_2
 2. SO_3^{2-} (sulfite)
 3. COCl_2 (C is central, bonded to O and both Cl)
 4. ClO_2^- (chlorite; Cl is central)
 5. OCS (C is central)
 6. NH_4^+
-

Answer Key — Full Solutions

1. CH_2Cl_2

- Valence e^- : $4 + 2(1) + 2(7) = \mathbf{20}$.
- Lewis: C is central with 2 H and 2 Cl. 4 bonds use 8, leaving 12, which is 3 lone pairs on each Cl. C has 8 ✓; the formal charges are all 0.
- Electron geometry: 4 atoms + 0 lone pairs = 4 domains, so **tetrahedral**.
- Molecular geometry: **tetrahedral, 109.5°** .
- Bond polarity: C-H 0.4 is **nonpolar**; C-Cl 0.5 is **polar**.
- Molecular polarity: **polar**. The outer atoms aren't identical, so the C-Cl dipoles aren't cancelled.

2. SO_3^{2-}

- Valence e^- : $6 + 3(6) + \mathbf{2} = \mathbf{26}$.
- Lewis (from C2): S is central with 3 single bonds (6), each O has 3 lone pairs (18), and the last 2 electrons go on S as a lone pair. The formal charges are S +1 and each O -1, which sum to -2 ✓. Brackets and 2-.
- Electron geometry: 3 atoms + 1 lone pair = 4 domains, so **tetrahedral**.
- Molecular geometry: **trigonal pyramidal, $\approx 107^\circ$** .
- Bond polarity: S-O: $3.5 - 2.5 = 1.0$, **polar**.
- Molecular polarity: **polar**. The lone pair on S makes it unsymmetric.

3. COCl_2 (phosgene)

- Valence e^- : $4 + 6 + 2(7) = \mathbf{24}$.
- Lewis: C is central with O and 2 Cl. 3 bonds use 6, leaving 18. That gives 3 lone pairs on each of O, Cl and Cl, leaving 0. C has only 6 ✗, so move an O lone pair to make **$\text{C}=\text{O}$** . The formal charges are all 0.
- Electron geometry: 3 atoms + 0 lone pairs = 3 domains, so **trigonal planar**.
- Molecular geometry: **trigonal planar, 120°** .
- Bond polarity: $\text{C}=\text{O}$ 1.0 is **polar**; C-Cl 0.5 is **polar**.
- Molecular polarity: **polar**. It is symmetric in shape, but $\text{O} \neq \text{Cl}$, and the $\text{C}=\text{O}$ dipole is bigger than the two C-Cl dipoles.

4. ClO_2^- (chlorite)

- Valence e^- : $7 + 2(6) + \mathbf{1} = \mathbf{20}$.
- Lewis: the skeleton O-Cl-O uses 4, leaving 16. Each O gets 3 lone pairs (12), leaving 4, which go on Cl as **2 lone pairs**. Cl has 2 bonds + 2 lone pairs = 8 ✓, so no step 4 is needed. The formal charges are $\text{Cl} = 7 - 4 - 2 = +1$ and each O = -1, which sum to -1 ✓. Brackets and -.
- Electron geometry: 2 atoms + 2 lone pairs = 4 domains, so **tetrahedral**.
- Molecular geometry: **bent, $\approx 104.5^\circ$** (less than 109.5°), the same pattern as water.
- Bond polarity: Cl-O: $3.5 - 3.0 = 0.5$, **polar**.
- Molecular polarity: **polar** (bent).


5. OCS

- Valence e^- : $6 + 4 + 6 = \mathbf{16}$.

- Lewis: exactly like CO₂: **O=C=S**, with 2 lone pairs on O and 2 on S. The formal charges are all 0.
- Electron geometry: 2 atoms + 0 lone pairs = 2 domains, so **linear**.
- Molecular geometry: **linear, 180°**.
- Bond polarity: C=O 1.0 is **polar**; C=S: 2.5 – 2.5 = 0, **nonpolar**.
- Molecular polarity: **polar**. It is linear like CO₂, but the two ends are different, so only the C=O dipole is there and nothing cancels it.

6. NH₄⁺

- Valence e⁻: 5 + 4(1) – 1 = **8**.
- Lewis: N with 4 N-H bonds (8 electrons), no lone pairs. The formal charge on N is 5 – 0 – 4 = +1 ✓. Brackets and +.
- Electron geometry: 4 atoms + 0 lone pairs = 4 domains, so **tetrahedral**.
- Molecular geometry: **tetrahedral, 109.5°**.
- Bond polarity: N-H: 3.0 – 2.1 = 0.9, **polar**.
- Molecular polarity: **nonpolar**. It is tetrahedral with 4 identical atoms, so the four equal dipoles cancel. (It still carries its + charge; “nonpolar” is about the dipoles.)



Notes for next session

VSEPR shows **where** the atoms go but not **how** the orbitals make that possible. NCERT §4.5 (valence bond theory) and §4.6 (**hybridisation**: sp , sp^2 , sp^3) answer that: tetrahedral CH_4 is sp^3 , trigonal planar BF_3 is sp^2 , and linear $BeCl_2$ is sp . Polarity also leads into **intermolecular forces** (NCERT §4.9 starts with hydrogen bonding): polar molecules attract each other (dipole-dipole forces), which is why water (polar) boils far higher than methane (nonpolar), and why “like dissolves like”. Check with Dhairya which of these his class takes next before planning C4.