

M20550 Calculus III Tutorial
Worksheet 6

1. Evaluate the double integral $\iint_R (4 - 2y) dA$, for $R = [0, 1] \times [0, 1]$, by identifying it as the volume of a solid.
2. Evaluate the iterated integral.

(a) $\int_0^2 \int_0^\pi r \sin^2 \theta \, d\theta dr$

(b) $\iint_R y e^{-xy} dA$ on $R = [0, 2] \times [0, 3]$

3. Find the volume of the solid in the first octant bounded by the cylinder $z = 16 - x^2$ and the plane $y = 5$.
4. Use polar coordinates to show that

$$\int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} e^{-(x^2+y^2)} dA = \pi$$

and deduce that $\int_{-\infty}^{+\infty} e^{-x^2} dx = \sqrt{\pi}$.

5. Evaluate the given integral.

$$\iint_R \arctan\left(\frac{y}{x}\right) dA$$

where $R = \{(x, y) : 1 \leq x^2 + y^2 \leq 4, 0 \leq y \leq x\}$.

6. Find the volume of the solid enclosed by the paraboloid $z = x^2 + y^2$ and the plane $z = 1$.
7. Set up, but do not solve, the integral that gives the volume of the solid region bounded by the paraboloid $z = 3x^2 + 3y^2$ and the cone $z = 4 - \sqrt{x^2 + y^2}$.
8. (Optional) Find the maximum value of the function $f(x, y, z) = x + 2y$ on the curve of intersection of the plane $x + y + z = 1$ and the cylinder $y^2 + z^2 = 4$.
9. (Optional) The plane $x + y + 2z = 2$ intersects the paraboloid $z = x^2 + y^2$ in an ellipse. Find the points on the ellipse that are nearest and farthest from the origin.