


SMART INDIA HACKATHON 2026



TADIT (तड़ित)

AI nowcasting of thunderstorms & lightning, 0–6 hours ahead

- **Problem Statement ID** – [SIH26072](#)
- **Problem Statement Title**- [AIML Based Nowcasting of Thunderstorm and Lightning](#)
- **Theme**- [Disaster Management](#)
- **PS Category**- [Software](#)
- **Team ID**-
- **Team Name** (Registered on portal)



Your
Team
Name

TADIT: AI LIGHTNING NOWCASTING



Tadit (तड़ित = lightning) fuses **radar, satellite, lightning and weather-model** data into one AI that forecasts **where lightning will strike**, in 10-min steps to 3 h and hourly to 6 h.

[Live prototype: tadit.pages.dev](https://tadit.pages.dev)

Proposed Solution



Multimodal AI fusion: one model reads Doppler radar, INSAT / GK2A satellite imagery, lightning strikes and GFS weather-model fields together.



Seamless 0–6 h: Tier 1 deep-learning nowcast in 10-min steps for 0–3 h, then Tier 2 AI-corrected NWP hourly for hours 4–6, switching at the measured crossover hour.




What it delivers: lightning probability per 16 km cell, radar-like storm intensity at 2 km, and LOW / MODERATE / HIGH risk with plain advice for each city.



Built for India's gaps: keeps working without radar or a lightning feed; runs today on INSAT-3DR/3DS or GK2A satellite + GFS.



Working prototype: the Tadit web app shows an animated India map and hourly city risk, live at tadit.pages.dev.



Working prototype: Bhubaneswar lightning outbreak, 2 Sep 2023, forecast issued 13:30 IST (satellite + GFS, no radar)



Observe
radar, satellite,
lightning, NWP



Fuse
one multimodal AI
model



Forecast
10-min steps to 3 h,
hourly to 6 h



Alert
city risk + advice

0.613

lightning CSI at hour 1, 1.8x
persistence

6 / 6

lead hours beating raw NWP
lightning

India

beats persistence vs FY-4A satellite
lightning

How It Addresses the Problem



3 h → 6 h

IMD's nowcasts look 3 h ahead. Tadit forecasts **6 h**: 10-min steps to 3 h, then hourly.



No radar? Still works

Large parts of India lack radar. Satellite + NWP alone reaches **0.461** CSI at hour 1.



From maps to action

City-level hourly risk with advice ("go indoors now") for officials and the public.



Innovation & Uniqueness

1

Mid-fusion U-Net with NWP cross-attention + FiLM: all 18 lead times in one pass.

2

Modality dropout: robust to any missing sensor, with free drop-a-sensor tests.

3

Measured tier crossover stitches nowcast and NWP into one 0–6 h product.

4

India-adapted in training (INSAT / GFS-like inputs) with leak-free lightning labels.

Your
Team
Name

TECHNICAL APPROACH



System architecture: from raw observations to a 0–6 h lightning forecast

1 · DATA SOURCES



Radar

NEXRAD VIL (training); IMD DWR reader built, not yet on live data



Satellite

INSAT-3DR/3DS · GK2A · GOES IR



Lightning

GLM (training); ILDN partnership offered



Weather model

GFS 0.25° · HRRR

2 · PREPROCESSING



384 km tiles on one map grid



Radar / satellite at 2 km, lightning at 8 km



Leak-free lightning labels (past 5 min only)



NWP only from runs issued before the forecast



50% of samples degraded to INSAT & GFS quality

3 · AI MODELS

TIER 1 · Fusion nowcaster (0–3 h, 10-min steps)

60M params · modality dropout

Per-source encoders

each sensor at native res.

Gated fusion

+ ‘missing’ embeddings

U-Net + attention

NWP cross-attn · FiLM

18 lead times

VIL 2 km · lightning 8 km

TIER 2 · AI-corrected NWP (lead hours 1–6)

GFS / HRRR forecast fields

Compact U-Net

Lightning + VIL, 16 km

4 · SEAMLESS OUTPUT

Switch at hour 3

Tier 1 for hours 1–3, Tier 2 for 4–6: the crossover measured on held-out storms.



Probability maps



JSON: hourly risk + sources used



Tadit web app & alerts

Methodology



Data: 3,000 real US storm events (SEVIR) + HRRR; split by date, so test storms are unseen.



Train: Intensity-weighted radar loss + focal lightning loss; modality dropout; one night on an RTX 5060 Ti.



Evaluate: CSI / POD / FAR per lead hour vs persistence and raw NWP; drop-a-sensor tests.




India: Run on INSAT / GK2A + GFS; checked against FY-4A satellite lightning.




Serve: Maps + JSON exported to a static web app on Cloudflare Pages.

Forecasts vs Reality

Held-out US storm: forecast lightning chance (shading) vs observed flashes (o)
Oklahoma City, 26 Aug 2019: CSI **0.61 / 0.54 / 0.55** for hours 1–3 vs persistence 0.52 / 0.33 / 0.18



India, Bhubaneswar 2 Sep 2023: hours 1–3 from satellite + GFS only (★ city)



Technology Stack

AI / ML

Python

PyTorch

NumPy

bf16

DATA

xarray

h5py

s3fs

Herbie

pyproj

cfgrib

WEB

React

TypeScript

Vite

MapLibre GL

INFRA

AWS Open Data

Cloudflare Pages

RTX 5060 Ti (16 GB)

Your
Team
Name

FEASIBILITY AND VIABILITY



Technical Feasibility

Built, trained and tested today



Working two-tier AI: 60M-parameter fusion U-Net for 0–3 h plus an NWP post-processor for 3–6 h, trained on 3,000 real storm events.



Fast: a full 0–6 h forecast takes **a few minutes per city**, mostly data download; the model itself runs in seconds.



Light: trained overnight on **one 16 GB consumer GPU**; the deployed model file is 120 MB.



Reliable build: open-source Python / PyTorch stack, covered by **38 automated tests**.



Operational Feasibility

Runs on India's own data



Indian data readers: INSAT, IMD radar, GFS and strike feeds; **INSAT and GFS verified on real files**.



INSAT cross-checked: our INSAT reader matches the GK2A satellite on the same scene ($r = 0.87$).



Works with gaps: runs on whichever sources are live, and every output lists what it used.



Easy to plug in: standard map + JSON outputs for IMD's nowcast portal and alert apps; the web viewer needs no backend.



Economic Viability

Low cost, scalable, policy-aligned



No new hardware: reuses radar, satellite and NWP data IMD already collects; open GK2A / GFS as backup.



Near-zero serving cost: forecasts are static files served from a free CDN (Cloudflare Pages).



Scales simply: ≈ 30 tiles of 384 km cover India, and one GPU can refresh them all.



Policy fit: supports MoES's **Mission Mausam** (₹2,000 cr) goals of more frequent nowcasts and no undetected weather events.



Potential Challenges & Risks

01

Limited access to data: Indian lightning archives (ILLN / ILDN) and radar volumes are not openly available, so early findings may be misleading.

02

Trained on US storms, used in India: monsoon storms, terrain and INSAT's slower scans differ from the US training data.

03

Sparse radar and sensor outages: radar does not cover all of India, and feeds drop out during storms.

04

Calibration and false alarms: probabilities are tuned to US storm frequency, not Indian climate.

05

Adoption into IMD operations: a new tool must fit forecasters' workflows and existing alert channels.



Strategies to Overcome Them



Improves with Indian data: ILDN (Tripura University) has offered to partner and share lightning data for SIH-2026; **as the model gets validated data, it recalibrates and gets more accurate.**

01



India-mode training: half of all samples are degraded to INSAT (4 km, 15–30 min scans) and GFS (~24 km) quality; fine-tune on Indian cases as they come in.

02



Modality dropout: the model runs on whatever is live. Without radar, hour-1 CSI only dips from **0.613 to 0.592**; satellite + NWP alone still scores **0.461**.

03



Regional recalibration with IMD, and alerts issued as risk levels, not raw numbers. On a dry Delhi day the forecast stayed at **0.04**.

04




Phased rollout: pilot at one regional centre (e.g. Bhubaneswar) side by side with current nowcasts, then scale state by state.

05

CSI = critical success index for lightning (higher is better), on held-out 2019 US storms. Trained on US GLM lightning, adapted to INSAT/GFS, not yet verified against Indian lightning observations. Tile count is an estimate from India's land area.

Your
Team
Name

IMPACT AND BENEFITS




⚡ **2,558 lives**

lost to lightning in India in 2023: 39.7% of all deaths from forces of nature, the largest single share (NCRB). Today's operational nowcasts look only 3 hours ahead.

PS SIH26072 · MoES / IMD
Theme: Disaster Management

Potential Impact on Target Audience



Benefits of the Solution

- Social**
Saves lives where lightning strikes hardest: MP, Bihar, Odisha, UP and Jharkhand account for 59% of India's lightning deaths. Timely alerts build public trust in IMD warnings.
- Economic**
Cuts losses to crops, livestock, flights and power grids. Built on existing and open data (IMD radar, INSAT, GFS): no new sensors are needed, and the model runs on a single GPU.
- Environmental**
Flags lightning-prone zones for forest-fire watch, supports climate adaptation as severe convective weather grows, and helps solar and grid operators plan for storms.
- Strategic**
Atmanirbhar: runs on India's own INSAT-3DR/3DS satellites and GFS, and keeps forecasting when radar or lightning feeds drop out, so it covers districts beyond radar range.

Measured, not claimed

Held-out 2019 storms scored against real satellite lightning (GLM).

1.8×

better than persistence at hour 1

Lightning CSI 0.613 vs 0.341; the gap grows to 5.5× by hour 3

+70%

over raw NWP lightning at hour 6

CSI 0.297 vs 0.175, and ahead of NWP at every hour from 1 to 6

0.461

hour-1 CSI with no radar


Satellite + NWP only: the usual case outside Indian radar coverage

0.71 vs 0.04

India: storm vs dry day

Storm probability 1 h ahead: Bhubaneswar outbreak (2 Sep 2023) vs Delhi dry day

Lightning CSI by lead hour (higher is better)



“From a 3-hour to a 6-hour lightning warning: one seamless AI nowcast, built on India's own satellites.”



CSI = critical success index for lightning in a 16 km cell during each lead hour; Tier 1 is used for hours 1–3 and Tier 2 for hours 4–6. India probabilities from GK2A satellite + GFS input, no radar. Trained on US GLM lightning, adapted to INSAT/GFS, not yet verified against Indian lightning observations. Death figures: NCRB, Accidental Deaths & Suicides in India 2023.

Data Sources

- TRAIN

SEVIR storm events: radar VIL, GOES-16 IR, GLM lightning. Veillette et al., NeurIPS 2020 · [registry.opendata.aws/sevir](#)
- TRAIN

HRRR US weather model (NWP input + baseline), NOAA · [registry.opendata.aws/noaa-hrrr-pds](#)
- INDIA

GFS 0.25° global weather model, NOAA · [registry.opendata.aws/noaa-gfs-bdp-pds](#)
- INDIA

INSAT-3DR Imager L1B (3RIMG_L1B_STD), ISRO / SAC · [mosdac.gov.in](#)
- INDIA

GK2A AMI L1B satellite, KMA via NOAA, no login · [registry.opendata.aws/noaa-gk2a-pds](#)
- VERIFY

FY-4A LMI corrected lightning events, Zhang et al., ESSD 18, 5773–5791 (2026), CC-BY 4.0 · [doi:10.5194/essd-18-5773-2026](#) · [data](#)
- TESTED

ISRO Cherrapunji DWR (RSCHR_L2B_STD), tested, not used · [mosdac.gov.in](#)
- MAPS

Survey of India boundaries (OVSF/1M/7) · [onlinemaps.surveyofindia.gov.in](#)
- PARTNER

ILDN Indian Lightning Detection Network, CeLTS, Tripura University: partnership offered (Sep 2026) · [ildn.in](#)

Methods & Research Papers

- MODEL DESIGN
 - Ronneberger et al., **U-Net**, MICCAI 2015 · [arXiv:1505.04597](#)
 - Perez et al., **FILM** conditioning, AAAI 2018 · [arXiv:1709.07871](#)
 - Leinonen et al., **Thunderstorm nowcasting with deep learning: multi-hazard data fusion**, GRL 2023 (closest prior work) · [doi](#)
- DEEP-LEARNING NOWCASTING
 - Shi et al., **ConvLSTM** precipitation nowcasting, NeurIPS 2015 · [arXiv:1506.04214](#)
 - Ravuri et al., **Skilful nowcasting with deep generative models**, Nature 597 (2021) · [doi](#)
 - Gao et al., **Earthformer** (SEVIR benchmark), NeurIPS 2022 · [arXiv:2207.05833](#)
- SENSORS & WEATHER MODELS
 - Dowell et al., **HRRR Part I**, Weather & Forecasting 37 (2022) · [doi](#)
 - Goodman et al., **GOES-R GLM**, Atmos. Research 125 (2013) · [doi](#)
 - NOAA/NASA, **GLM Lightning Cluster-Filter ATBD** (330 ms / 16.5 km flashes) · [pdf](#)
 - Blaylock, **Herbie** NWP download tool, Zenodo · [doi](#)
- OUR WORK
 - Tadit frozen results and scorecard, [results/RESULTS.md](#) · live site [tadit.pages.dev](#)
- VERIFICATION & PHYSICS
 - Schaefer, **Critical success index** as a warning-skill score, Weather & Forecasting 1990
 - Greene & Clark, **Vertically integrated liquid (VIL)** from radar, Monthly Weather Review 1972
 - Maddox, **Mesoscale convective complexes** (−52 °C cloud-top threshold), BAMS 1980



Live prototype: click here
[tadit.pages.dev](#)



Code & results: click here
[github.com/adit1290codesrank/sih2026](#)



India Check vs Real Lightning

FY-4A satellite lightning, CSI for hours 1 / 2 / 3 (persistence in brackets)

Bhubaneswar · GK2A **0.24 / 0.22 / 0.20** (0.20 / 0.01 / 0.00)

Bhubaneswar · INSAT **0.24 / 0.16 / 0.13** (0.20 / 0.01 / 0.00)

Guwahati · INSAT **hour 6: MODERATE risk** storm hit the city then



Real Storms We Forecast

Bhubaneswar, **2 Sep 2023**: 61,000 lightning strikes in two hours ([Organiser](#)); 10 killed in six districts ([Deccan Herald](#))
Kolkata, **9 May 2024**: thundersquall and hailstorms lash the city ([ThePrint](#))



Government Sources

NCRB, **Accidental Deaths & Suicides in India 2023** (lightning deaths) · [data.gov.in](#)
IMD **station-wise nowcast** service (3 h horizon) · [mausam.imd.gov.in](#)
Cabinet approves **Mission Mausam, 2024** · [pmindia.gov.in](#)