

Risk and Decisions

Paolo Turrini

Department of Computing, Imperial College London

Introduction to Artificial Intelligence

The lectures

- The agent and the world (**Knowledge Representation**)
 - Actions and knowledge
 - Inference
- Good decisions (**Risk and Decisions**)
 - Chance
 - Gains
- Good decisions in time (**Markov Decision Processes**)
 - Chance and gains in time
 - Patience
 - Finding the best strategy
- Learning from experience (**Reinforcement Learning**)
 - Finding a reasonable strategy

Risk and Decisions

Knowing what to expect

Today

- Probabilities: a crash course
- Bayes' rule and conditional independence
- Back to the Wumpus World

The book



Stuart Russell and Peter Norvig

Artificial Intelligence: a modern approach

Chapters 13-14

Holiday

You are back from a holiday on an exotic island, and your doctor has bad news and good news. The bad news is that you've been diagnosed a serious disease and the test is 99% accurate. The good news is that the disease is very rare (1 in 10.000 get it).

Holiday

You are back from a holiday on an exotic island, and your doctor has bad news and good news. The bad news is that you've been diagnosed a serious disease and the test is 99% accurate. The good news is that the disease is very rare (1 in 10.000 get it).

How worried should you be?

Probability basics

Begin with a set Ω —the sample space

Probability basics

Begin with a set Ω —the **sample space**
e.g., 6 possible rolls of a dice.

Probability basics

Begin with a set Ω —the **sample space**
e.g., 6 possible rolls of a dice.

$w \in \Omega$ is a **sample point/possible world/atomic event**

Probability basics

A **probability space** or **probability model** is a sample space Ω with an assignment $P(w)$ for every $w \in \Omega$ s.t.

Probability basics

A **probability space** or **probability model** is a sample space Ω with an assignment $P(w)$ for every $w \in \Omega$ s.t.

$$0 \leq P(w) \leq 1$$

Probability basics

A **probability space** or **probability model** is a sample space Ω with an assignment $P(w)$ for every $w \in \Omega$ s.t.

$$0 \leq P(w) \leq 1$$

$$\sum_w P(w) = 1$$

Probability basics

A **probability space** or **probability model** is a sample space Ω with an assignment $P(w)$ for every $w \in \Omega$ s.t.

$$0 \leq P(w) \leq 1$$

$$\sum_w P(w) = 1$$

e.g., $P(1) = P(2) = P(3) = P(4) = P(5) = P(6) = 1/6$.

Events

An event A is any subset of Ω

Events

An event A is any subset of Ω

$$P(A) = \sum_{\{w \in A\}} P(w)$$

Events

An event A is any subset of Ω

$$P(A) = \sum_{\{w \in A\}} P(w)$$

E.g.,

$$P(\text{dice roll} < 4) = P(1) + P(2) + P(3) = 1/6 + 1/6 + 1/6 = 1/2$$

Random variables

A **random variable** is a function from sample points to some range, e.g., \mathbb{R} , $[0, 1]$, $\{true, false\}$...

Random variables

A **random variable** is a function from sample points to some range,
e.g., \mathbb{R} , $[0, 1]$, $\{true, false\}$...

e.g., $Odd(1) = true$.

Random variables

A **random variable** is a function from sample points to some range, e.g., \mathbb{R} , $[0, 1]$, $\{\text{true}, \text{false}\}$...

e.g., $\text{Odd}(1) = \text{true}$.

P induces a **probability distribution** for any random variable X :

$$P(X = x_i) = \sum_{\{w: X(w) = x_i\}} P(w)$$

Random variables

A **random variable** is a function from sample points to some range, e.g., \mathbb{R} , $[0, 1]$, $\{\text{true}, \text{false}\}$...

e.g., $\text{Odd}(1) = \text{true}$.

P induces a **probability distribution** for any random variable X :

$$P(X = x_i) = \sum_{\{w: X(w) = x_i\}} P(w)$$

e.g.,

$$P(\text{Odd} = \text{true}) = P(1) + P(3) + P(5) = 1/6 + 1/6 + 1/6 = 1/2$$

Conditional probability

Definition of conditional probability:

$$P(a|b) = \frac{P(a \wedge b)}{P(b)} \text{ if } P(b) \neq 0$$

Conditional probability

Definition of conditional probability:

$$P(a|b) = \frac{P(a \wedge b)}{P(b)} \text{ if } P(b) \neq 0$$

Product rule gives an alternative formulation:

$$P(a \wedge b) = P(a|b)P(b) = P(b|a)P(a)$$

Conditional probability

Definition of conditional probability:

$$P(a|b) = \frac{P(a \wedge b)}{P(b)} \text{ if } P(b) \neq 0$$

Product rule gives an alternative formulation:

$$P(a \wedge b) = P(a|b)P(b) = P(b|a)P(a) \text{ but then...}$$

Bayes' Rule

Theorem (Bayes' Rule)

$$P(a|b) = \frac{P(b|a)P(a)}{P(b)}$$

Bayes' Rule

Useful for assessing **causal** probability from **diagnostic** probability:

$$P(Cause|Effect) = \frac{P(Effect|Cause)P(Cause)}{P(Effect)}$$

Bayes' Rule

Useful for assessing **causal** probability from **diagnostic** probability:

$$P(\textit{Cause}|\textit{Effect}) = \frac{P(\textit{Effect}|\textit{Cause})P(\textit{Cause})}{P(\textit{Effect})}$$

E.g., let *c* be cold, *s* be sore throat:

$$P(c|s) = \frac{P(s|c)P(c)}{P(s)} = \frac{0.9 \times 0.001}{0.005} = 0.18$$

Bayes' rule

$$P(c|s) = \frac{P(s|c)P(c)}{P(s)} = \frac{0.9 \times 0.001}{0.005} = 0.18$$

Bayes' rule

$$P(c|s) = \frac{P(s|c)P(c)}{P(s)} = \frac{0.9 \times 0.001}{0.005} = 0.18$$

We might not know the prior probability of the evidence $P(S)$

Bayes' rule

$$P(c|s) = \frac{P(s|c)P(c)}{P(s)} = \frac{0.9 \times 0.001}{0.005} = 0.18$$

We might not know the prior probability of the evidence $P(S)$
In this case...

Bayes' rule

$$P(c|s) = \frac{P(s|c)P(c)}{P(s)} = \frac{0.9 \times 0.001}{0.005} = 0.18$$

We might not know the prior probability of the evidence $P(S)$
In this case...

- we compute the posterior probability for each value of the query variable ($c, \neg c$)

Bayes' rule

$$P(c|s) = \frac{P(s|c)P(c)}{P(s)} = \frac{0.9 \times 0.001}{0.005} = 0.18$$

We might not know the prior probability of the evidence $P(S)$
In this case...

- we compute the posterior probability for each value of the query variable ($c, \neg c$)
- and then normalise by a normalisation constant α

Bayes' rule

$$P(c|s) = \frac{P(s|c)P(c)}{P(s)} = \frac{0.9 \times 0.001}{0.005} = 0.18$$

We might not know the prior probability of the evidence $P(S)$
In this case...

- we compute the posterior probability for each value of the query variable ($c, \neg c$)
- and then normalise by a normalisation constant α

$$P(C|s) = \alpha \langle P(s|c)P(c), P(s|\neg c)P(\neg c) \rangle$$

Bayes' rule with random variables

Theorem (Bayes' rule with random variables)

$$P(X|Y) = \alpha P(Y|X)P(X)$$

Holiday solved

You are back from a holiday on an exotic island, and your doctor has bad news and good news. The bad news is that you've been diagnosed a serious disease and the test is 99% accurate. The good news is that the disease is very rare (1 in 10.000 get it).

Holiday solved

You are back from a holiday on an exotic island, and your doctor has bad news and good news. The bad news is that you've been diagnosed a serious disease and the test is 99% accurate. The good news is that the disease is very rare (1 in 10.000 get it).

E.g., let d be disease, p be that you scored positive at the test:

Holiday solved

You are back from a holiday on an exotic island, and your doctor has bad news and good news. The bad news is that you've been diagnosed a serious disease and the test is 99% accurate. The good news is that the disease is very rare (1 in 10.000 get it).

E.g., let d be disease, p be that you scored positive at the test:

$$P(d|p)$$

Holiday solved

You are back from a holiday on an exotic island, and your doctor has bad news and good news. The bad news is that you've been diagnosed a serious disease and the test is 99% accurate. The good news is that the disease is very rare (1 in 10.000 get it).

E.g., let d be disease, p be that you scored positive at the test:

$$P(d|p) = \frac{P(p|d)P(d)}{P(p|d)P(d) + P(p|\neg d)P(\neg d)}$$

Holiday solved

You are back from a holiday on an exotic island, and your doctor has bad news and good news. The bad news is that you've been diagnosed a serious disease and the test is 99% accurate. The good news is that the disease is very rare (1 in 10.000 get it).

E.g., let d be disease, p be that you scored positive at the test:

$$P(d|p) = \frac{P(p|d)P(d)}{P(p|d)P(d) + P(p|\neg d)P(\neg d)} = \frac{0.99 \times 0.0001}{0.99 \times 0.0001 + 0.01 \times 0.9999}$$

Holiday solved

You are back from a holiday on an exotic island, and your doctor has bad news and good news. The bad news is that you've been diagnosed a serious disease and the test is 99% accurate. The good news is that the disease is very rare (1 in 10.000 get it).

E.g., let d be disease, p be that you scored positive at the test:

$$P(d|p) = \frac{P(p|d)P(d)}{P(p|d)P(d) + P(p|\neg d)P(\neg d)} = \frac{0.99 \times 0.0001}{0.99 \times 0.0001 + 0.01 \times 0.9999} = 0.0098$$

Holiday solved

You are back from a holiday on an exotic island, and your doctor has bad news and good news. The bad news is that you've been diagnosed a serious disease and the test is 99% accurate. The good news is that the disease is very rare (1 in 10.000 get it).

E.g., let d be disease, p be that you scored positive at the test:

$$P(d|p) = \frac{P(p|d)P(d)}{P(p|d)P(d) + P(p|\neg d)P(\neg d)} = \frac{0.99 \times 0.0001}{0.99 \times 0.0001 + 0.01 \times 0.9999} = 0.0098$$

Notice: the posterior probability of disease is still very small!

Independence

A and B are independent iff

$$\mathbf{P}(A|B) = \mathbf{P}(A) \quad \text{or} \quad \mathbf{P}(B|A) = \mathbf{P}(B) \quad \text{or} \quad \mathbf{P}(A, B) = \mathbf{P}(A)\mathbf{P}(B)$$

$$\begin{aligned} &P(cavity) \\ = &P(cavity|Weather) \\ = &P(cavity|CristianoRonaldo\text{scores}) \end{aligned}$$

Combining evidence

Start with the joint distribution:

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	.108	.012	.072	.008
\neg <i>cavity</i>	.016	.064	.144	.576

$$P(\text{Cavity} | \text{toothache} \wedge \text{catch}) =$$

Combining evidence

Start with the joint distribution:

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	.108	.012	.072	.008
\neg <i>cavity</i>	.016	.064	.144	.576

$$P(Cavity | toothache \wedge catch) = \alpha \langle 0.108, 0.016 \rangle$$

Combining evidence

Start with the joint distribution:

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	.108	.012	.072	.008
\neg <i>cavity</i>	.016	.064	.144	.576

$$P(Cavity | toothache \wedge catch) = \alpha \langle 0.108, 0.016 \rangle =$$

Combining evidence

Start with the joint distribution:

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	.108	.012	.072	.008
\neg <i>cavity</i>	.016	.064	.144	.576

$$\mathbf{P}(\text{Cavity} | \text{toothache} \wedge \text{catch}) = \alpha \langle 0.108, 0.016 \rangle = \langle 0.871, 0.129 \rangle$$

Combining evidence

Start with the joint distribution:

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	.108	.012	.072	.008
\neg <i>cavity</i>	.016	.064	.144	.576

- It doesn't scale up to a large number of variables
- Can we simplify?

Combining evidence

We can't use absolute independence:

Combining evidence

We can't use absolute independence:

- Toothache and Catch are **not** independent: If the probe catches in the tooth then it is likely the tooth has a cavity, which means that toothache is likely too.

Combining evidence

We can't use absolute independence:

- Toothache and Catch are **not** independent: If the probe catches in the tooth then it is likely the tooth has a cavity, which means that toothache is likely too.
- But they are independent **given** the presence or the absence of cavity! Toothache depends on the state of the nerves in the tooth, catch depends on the dentist's skills, to which toothache is irrelevant

Conditional independence

- ① $P(\text{catch}|\text{toothache}, \text{cavity}) = P(\text{catch}|\text{cavity})$, the same independence holds if I haven't got a cavity:

Conditional independence

- ① $P(\text{catch}|\text{toothache}, \text{cavity}) = P(\text{catch}|\text{cavity})$, the same independence holds if I haven't got a cavity:
- ② $P(\text{catch}|\text{toothache}, \neg \text{cavity}) = P(\text{catch}|\neg \text{cavity})$

Conditional independence

- ① $P(\text{catch}|\text{toothache}, \text{cavity}) = P(\text{catch}|\text{cavity})$, the same independence holds if I haven't got a cavity:
- ② $P(\text{catch}|\text{toothache}, \neg \text{cavity}) = P(\text{catch}|\neg \text{cavity})$

Catch is conditionally independent of *Toothache* given *Cavity*:

Conditional independence

$$\mathbf{P}(\textit{Catch} | \textit{Toothache}, \textit{Cavity}) = \mathbf{P}(\textit{Catch} | \textit{Cavity})$$

Conditional independence

$$\mathbf{P}(\textit{Catch} | \textit{Toothache}, \textit{Cavity}) = \mathbf{P}(\textit{Catch} | \textit{Cavity})$$

Equivalent statements:

Conditional independence

$$\mathbf{P}(\textit{Catch} | \textit{Toothache}, \textit{Cavity}) = \mathbf{P}(\textit{Catch} | \textit{Cavity})$$

Equivalent statements:

$$\mathbf{P}(\textit{Toothache} | \textit{Catch}, \textit{Cavity}) = \mathbf{P}(\textit{Toothache} | \textit{Cavity})$$

Conditional independence

$$\mathbf{P}(\textit{Catch} | \textit{Toothache}, \textit{Cavity}) = \mathbf{P}(\textit{Catch} | \textit{Cavity})$$

Equivalent statements:

$$\mathbf{P}(\textit{Toothache} | \textit{Catch}, \textit{Cavity}) = \mathbf{P}(\textit{Toothache} | \textit{Cavity})$$

$$\mathbf{P}(\textit{Toothache}, \textit{Catch} | \textit{Cavity}) =$$

$$\mathbf{P}(\textit{Toothache} | \textit{Cavity})\mathbf{P}(\textit{Catch} | \textit{Cavity})$$

Conditional independence contd.

Write out full joint distribution using chain rule:

$P(\textit{Toothache}, \textit{Catch}, \textit{Cavity})$

Conditional independence contd.

Write out full joint distribution using chain rule:

$$\begin{aligned} & \mathbf{P}(Toothache, Catch, Cavity) \\ &= \mathbf{P}(Toothache|Catch, Cavity)\mathbf{P}(Catch, Cavity) \end{aligned}$$

Conditional independence contd.

Write out full joint distribution using chain rule:

$$\begin{aligned} & \mathbf{P}(Toothache, Catch, Cavity) \\ &= \mathbf{P}(Toothache|Catch, Cavity)\mathbf{P}(Catch, Cavity) \\ &= \mathbf{P}(Toothache|Catch, Cavity)\mathbf{P}(Catch|Cavity)\mathbf{P}(Cavity) \end{aligned}$$

Conditional independence contd.

Write out full joint distribution using chain rule:

$$\begin{aligned} & \mathbf{P}(Toothache, Catch, Cavity) \\ &= \mathbf{P}(Toothache|Catch, Cavity)\mathbf{P}(Catch, Cavity) \\ &= \mathbf{P}(Toothache|Catch, Cavity)\mathbf{P}(Catch|Cavity)\mathbf{P}(Cavity) \\ &= \mathbf{P}(Toothache|Cavity)\mathbf{P}(Catch|Cavity)\mathbf{P}(Cavity) \end{aligned}$$

Conditional independence contd.

Write out full joint distribution using chain rule:

$$\begin{aligned} & \mathbf{P}(\textit{Toothache}, \textit{Catch}, \textit{Cavity}) \\ &= \mathbf{P}(\textit{Toothache} | \textit{Catch}, \textit{Cavity}) \mathbf{P}(\textit{Catch}, \textit{Cavity}) \\ &= \mathbf{P}(\textit{Toothache} | \textit{Catch}, \textit{Cavity}) \mathbf{P}(\textit{Catch} | \textit{Cavity}) \mathbf{P}(\textit{Cavity}) \\ &= \mathbf{P}(\textit{Toothache} | \textit{Cavity}) \mathbf{P}(\textit{Catch} | \textit{Cavity}) \mathbf{P}(\textit{Cavity}) \end{aligned}$$

I.e., $2 + 2 + 1 = 5$ independent numbers (first and second steps remove two). Else $8-1=7$. The gain is bigger the more the combinations.

Conditional independence contd.

In most cases, the use of conditional independence reduces the size of the representation of the joint distribution from exponential in n to linear in n .

Conditional independence contd.

In most cases, the use of conditional independence reduces the size of the representation of the joint distribution from exponential in n to linear in n .

Conditional independence is our most basic and robust form of knowledge about uncertain environments.

Bayes' Rule and conditional independence

$$P(\textit{Cavity} | \textit{toothache} \wedge \textit{catch})$$

Bayes' Rule and conditional independence

$$\begin{aligned} & \mathbf{P}(Cavity | toothache \wedge catch) \\ &= \alpha \mathbf{P}(toothache \wedge catch | Cavity) \mathbf{P}(Cavity) \end{aligned}$$

Bayes' Rule and conditional independence

$$\begin{aligned} & \mathbf{P}(\textit{Cavity} | \textit{toothache} \wedge \textit{catch}) \\ &= \alpha \mathbf{P}(\textit{toothache} \wedge \textit{catch} | \textit{Cavity}) \mathbf{P}(\textit{Cavity}) \\ &= \alpha \mathbf{P}(\textit{toothache} | \textit{Cavity}) \mathbf{P}(\textit{catch} | \textit{Cavity}) \mathbf{P}(\textit{Cavity}) \end{aligned}$$

Bayes' Rule and conditional independence

$$\begin{aligned} & \mathbf{P}(\text{Cavity} | \text{toothache} \wedge \text{catch}) \\ &= \alpha \mathbf{P}(\text{toothache} \wedge \text{catch} | \text{Cavity}) \mathbf{P}(\text{Cavity}) \\ &= \alpha \mathbf{P}(\text{toothache} | \text{Cavity}) \mathbf{P}(\text{catch} | \text{Cavity}) \mathbf{P}(\text{Cavity}) \end{aligned}$$

This is an example of a **naive Bayes** model:



$$\mathbf{P}(\text{Cause}, \text{Effect}_1, \dots, \text{Effect}_n) = \mathbf{P}(\text{Cause}) \prod_i \mathbf{P}(\text{Effect}_i | \text{Cause})$$

Bayes' Rule and conditional independence

$$\begin{aligned}
 & \mathbf{P}(\text{Cavity} | \text{toothache} \wedge \text{catch}) \\
 &= \alpha \mathbf{P}(\text{toothache} \wedge \text{catch} | \text{Cavity}) \mathbf{P}(\text{Cavity}) \\
 &= \alpha \mathbf{P}(\text{toothache} | \text{Cavity}) \mathbf{P}(\text{catch} | \text{Cavity}) \mathbf{P}(\text{Cavity})
 \end{aligned}$$


This is an example of a **naive Bayes** model:

$$\mathbf{P}(\text{Cause}, \text{Effect}_1, \dots, \text{Effect}_n) = \mathbf{P}(\text{Cause}) \prod_i \mathbf{P}(\text{Effect}_i | \text{Cause})$$




The Wumps World


The Wumpus World




The Wumpus World

OK			
OK 	OK		


The Wumpus World




The Wumpus World




The Wumpus World




The Wumpus World




The Wumpus World



The Wumpus World



The Wumpus World



Wumpus World

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

Wumpus World

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

$P_{ij} = \text{true}$ iff $[i, j]$ contains a pit

$B_{ij} = \text{true}$ iff $[i, j]$ is breezy

Specifying the probability model

Specifying the probability model

Include only $B_{1,1}$, $B_{1,2}$, $B_{2,1}$ in the probability model!

Specifying the probability model

Include only $B_{1,1}$, $B_{1,2}$, $B_{2,1}$ in the probability model!

The full joint distribution is $\mathbf{P}(P_{1,1}, \dots, P_{4,4}, B_{1,1}, B_{1,2}, B_{2,1})$

Specifying the probability model

Include only $B_{1,1}$, $B_{1,2}$, $B_{2,1}$ in the probability model!

The full joint distribution is $\mathbf{P}(P_{1,1}, \dots, P_{4,4}, B_{1,1}, B_{1,2}, B_{2,1})$

Apply product rule:

$$\mathbf{P}(B_{1,1}, B_{1,2}, B_{2,1} \mid P_{1,1}, \dots, P_{4,4}) \mathbf{P}(P_{1,1}, \dots, P_{4,4})$$

(Do it this way to get $P(\textit{Effect} \mid \textit{Cause})$.)

Specifying the probability model

Include only $B_{1,1}, B_{1,2}, B_{2,1}$ in the probability model!

The full joint distribution is $\mathbf{P}(P_{1,1}, \dots, P_{4,4}, B_{1,1}, B_{1,2}, B_{2,1})$

Apply product rule:

$$\mathbf{P}(B_{1,1}, B_{1,2}, B_{2,1} \mid P_{1,1}, \dots, P_{4,4}) \mathbf{P}(P_{1,1}, \dots, P_{4,4})$$

(Do it this way to get $P(\textit{Effect} \mid \textit{Cause})$.)

First term: 1 if pits are adjacent to breezes, 0 otherwise

Specifying the probability model

Include only $B_{1,1}$, $B_{1,2}$, $B_{2,1}$ in the probability model!

The full joint distribution is $\mathbf{P}(P_{1,1}, \dots, P_{4,4}, B_{1,1}, B_{1,2}, B_{2,1})$

Apply product rule:

$$\mathbf{P}(B_{1,1}, B_{1,2}, B_{2,1} \mid P_{1,1}, \dots, P_{4,4}) \mathbf{P}(P_{1,1}, \dots, P_{4,4})$$

(Do it this way to get $P(\textit{Effect} \mid \textit{Cause})$.)

First term: 1 if pits are adjacent to breezes, 0 otherwise

Second term: pits are placed randomly, probability 0.2 per square:

Specifying the probability model

Include only $B_{1,1}, B_{1,2}, B_{2,1}$ in the probability model!

The full joint distribution is $\mathbf{P}(P_{1,1}, \dots, P_{4,4}, B_{1,1}, B_{1,2}, B_{2,1})$

Apply product rule:

$$\mathbf{P}(B_{1,1}, B_{1,2}, B_{2,1} \mid P_{1,1}, \dots, P_{4,4}) \mathbf{P}(P_{1,1}, \dots, P_{4,4})$$

(Do it this way to get $P(\textit{Effect} \mid \textit{Cause})$.)

First term: 1 if pits are adjacent to breezes, 0 otherwise

Second term: pits are placed randomly, probability 0.2 per square:

$$\mathbf{P}(P_{1,1}, \dots, P_{4,4}) = \prod_{i,j=1,1}^{4,4} \mathbf{P}(P_{i,j}) = 0.2^n \times 0.8^{16-n}$$

for n pits.

Observations and query

We know the following facts:

$$b = \neg b_{1,1} \wedge b_{1,2} \wedge b_{2,1}$$

Observations and query

We know the following facts:

$$b = \neg b_{1,1} \wedge b_{1,2} \wedge b_{2,1}$$

$$explored = \neg p_{1,1} \wedge \neg p_{1,2} \wedge \neg p_{2,1}$$

Observations and query

We know the following facts:

$$b = \neg b_{1,1} \wedge b_{1,2} \wedge b_{2,1}$$

$$explored = \neg p_{1,1} \wedge \neg p_{1,2} \wedge \neg p_{2,1}$$

Query is $\mathbf{P}(P_{1,3} | explored, b)$

Observations and query

We know the following facts:

$$b = \neg b_{1,1} \wedge b_{1,2} \wedge b_{2,1}$$

$$explored = \neg p_{1,1} \wedge \neg p_{1,2} \wedge \neg p_{2,1}$$

Query is $\mathbf{P}(P_{1,3} | explored, b)$

Define *Unexplored* = P_{ij} s other than $P_{1,3}$ and *Explored*

Complexity

For inference by enumeration, we have

$$\mathbf{P}(P_{1,3}|\textit{explored}, b) = \alpha \sum_{\textit{unexplored}} \mathbf{P}(P_{1,3}, \textit{unexplored}, \textit{explored}, b)$$

Complexity

For inference by enumeration, we have

$$\mathbf{P}(P_{1,3}|\textit{explored}, b) = \alpha \sum_{\textit{unexplored}} \mathbf{P}(P_{1,3}, \textit{unexplored}, \textit{explored}, b)$$

- There are 12 unknown squares

Complexity

For inference by enumeration, we have

$$\mathbf{P}(P_{1,3}|\textit{explored}, b) = \alpha \sum_{\textit{unexplored}} \mathbf{P}(P_{1,3}, \textit{unexplored}, \textit{explored}, b)$$

- There are 12 unknown squares
- The summation contains $2^{12} = 4096$ terms

Complexity

For inference by enumeration, we have

$$\mathbf{P}(P_{1,3}|\textit{explored}, b) = \alpha \sum_{\textit{unexplored}} \mathbf{P}(P_{1,3}, \textit{unexplored}, \textit{explored}, b)$$

- There are 12 unknown squares
- The summation contains $2^{12} = 4096$ terms

In general the summation grows exponentially with the number of squares!

Complexity

For inference by enumeration, we have

$$\mathbf{P}(P_{1,3}|\textit{explored}, b) = \alpha \sum_{\textit{unexplored}} \mathbf{P}(P_{1,3}, \textit{unexplored}, \textit{explored}, b)$$

- There are 12 unknown squares
- The summation contains $2^{12} = 4096$ terms

In general the summation grows exponentially with the number of squares!


And now?

Using conditional independence

Basic insight: observations are conditionally independent of other hidden squares given neighbouring hidden squares


Using conditional independence

Basic insight: observations are conditionally independent of other hidden squares given neighbouring hidden squares



Using conditional independence


Basic insight: observations are conditionally independent of other hidden squares given neighbouring hidden squares



Define $Unexplored = Fringe \cup Other$

Using conditional independence

Basic insight: observations are conditionally independent of other hidden squares given neighbouring hidden squares




Define $Unexplored = Fringe \cup Other$

$$P(b|P_{1,3}, Explored, Unexplored) = P(b|P_{1,3}, Explored, Fringe)$$

Using conditional independence

Basic insight: observations are conditionally independent of other hidden squares given neighbouring hidden squares



Define $Unexplored = Fringe \cup Other$

$P(b|P_{1,3}, Explored, Unexplored) = P(b|P_{1,3}, Explored, Fringe)$

Manipulate query into a form where we can use this!

Using conditional independence

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

$$P(P_{1,3} | \text{explored}, b)$$

Using conditional independence

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

$$\mathbf{P}(P_{1,3} | explored, b) = \alpha \sum_{unexplored} \mathbf{P}(P_{1,3}, unexplored, explored, b)$$

Using conditional independence

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

Inference by enumeration

$$\mathbf{P}(P_{1,3} | explored, b) = \alpha \sum_{unexplored} \mathbf{P}(P_{1,3}, unexplored, explored, b)$$

Using conditional independence

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

$$\propto \sum_{unexplored} \mathbf{P}(P_{1,3}, unexplored, explored, b)$$

Using conditional independence

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

$$\begin{aligned}
 & \alpha \sum_{unexplored} \mathbf{P}(P_{1,3}, unexplored, explored, b) \\
 &= \alpha \sum_{unexplored} \mathbf{P}(b|explored, P_{1,3}, unexplored) \times \\
 & \times \mathbf{P}(P_{1,3}, explored, unexplored)
 \end{aligned}$$

Using conditional independence

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

Product rule

$$\begin{aligned}
 & \alpha \sum_{unexplored} \mathbf{P}(P_{1,3}, unexplored, explored, b) \\
 &= \alpha \sum_{unexplored} \mathbf{P}(b|explored, P_{1,3}, unexplored) \times \\
 & \times \mathbf{P}(P_{1,3}, explored, unexplored)
 \end{aligned}$$

Using conditional independence

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

$$\propto \sum_{unexplored} \mathbf{P}(b|P_{1,3}, unexplored, explored) \mathbf{P}(P_{1,3}, unexplored, explored)$$

Using conditional independence

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

$$\begin{aligned}
 & \alpha \sum_{unexplored} \mathbf{P}(b|P_{1,3}, unexplored, explored) \mathbf{P}(P_{1,3}, unexplored, explored) \\
 & = \alpha \sum_{fringe} \sum_{other} \mathbf{P}(b|explored, P_{1,3}, fringe, other) \times \\
 & \times \mathbf{P}(P_{1,3}, explored, fringe, other)
 \end{aligned}$$

Using conditional independence

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

Distinguishing the unknown

$$\begin{aligned}
 & \alpha \sum_{unexplored} \mathbf{P}(b|P_{1,3}, unexplored, explored) \mathbf{P}(P_{1,3}, unexplored, explored) \\
 & = \alpha \sum_{fringe} \sum_{other} \mathbf{P}(b|explored, P_{1,3}, fringe, other) \times \\
 & \times \mathbf{P}(P_{1,3}, explored, fringe, other)
 \end{aligned}$$

Using conditional independence

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

$$\propto \sum_{fringe} \sum_{other} \mathbf{P}(b | explored, P_{1,3}, fringe, other) \times \\ \times \mathbf{P}(P_{1,3}, explored, fringe, other)$$

Using conditional independence

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

$$\begin{aligned}
 & \alpha \sum_{fringe} \sum_{other} \mathbf{P}(b|explored, P_{1,3}, fringe, other) \times \\
 & \times \mathbf{P}(P_{1,3}, explored, fringe, other) \\
 & = \alpha \sum_{fringe} \sum_{other} \mathbf{P}(b|explored, P_{1,3}, fringe) \times \\
 & \times \mathbf{P}(P_{1,3}, explored, fringe, other)
 \end{aligned}$$

Using conditional independence

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

Conditional Independence

$$\begin{aligned}
 & \alpha \sum_{fringe} \sum_{other} \mathbf{P}(b|explored, P_{1,3}, fringe, other) \times \\
 & \times \mathbf{P}(P_{1,3}, explored, fringe, other) \\
 & = \alpha \sum_{fringe} \sum_{other} \mathbf{P}(b|explored, P_{1,3}, fringe) \times \\
 & \times \mathbf{P}(P_{1,3}, explored, fringe, other)
 \end{aligned}$$

Using conditional independence

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

$$\propto \sum_{fringe} \sum_{other} \mathbf{P}(b|explored, P_{1,3}, fringe) \times \\ \times \mathbf{P}(P_{1,3}, explored, fringe, other)$$

Using conditional independence

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

$$\alpha \sum_{fringe} \sum_{other} \mathbf{P}(b|explored, P_{1,3}, fringe) \times \\ \times \mathbf{P}(P_{1,3}, explored, fringe, other)$$

$$= \alpha \sum_{fringe} \mathbf{P}(b|explored, P_{1,3}, fringe) \times \\ \times \sum_{other} \mathbf{P}(P_{1,3}, explored, fringe, other)$$

Using conditional independence

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

Pushing the sums inwards

$$\begin{aligned}
 & \alpha \sum_{fringe} \sum_{other} \mathbf{P}(b|explored, P_{1,3}, fringe) \times \\
 & \times \mathbf{P}(P_{1,3}, explored, fringe, other) \\
 & = \alpha \sum_{fringe} \mathbf{P}(b|explored, P_{1,3}, fringe) \times \\
 & \times \sum_{other} \mathbf{P}(P_{1,3}, explored, fringe, other)
 \end{aligned}$$

Using conditional independence

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

$$\alpha \sum_{fringe} P(b|explored, P_{1,3}, fringe) \times \sum_{other} P(P_{1,3}, explored, fringe, other)$$

Using conditional independence

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

$$\begin{aligned}
 & \alpha \sum_{fringe} \mathbf{P}(b|explored, P_{1,3}, fringe) \times \\
 & \sum_{other} \mathbf{P}(P_{1,3}, explored, fringe, other) \\
 &= \alpha \sum_{fringe} \mathbf{P}(b|explored, P_{1,3}, fringe) \times \\
 & \sum_{other} \mathbf{P}(P_{1,3}) P(explored) P(fringe) P(other)
 \end{aligned}$$

Using conditional independence

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

Independence

$$\begin{aligned}
 & \alpha \sum_{fringe} \mathbf{P}(b|explored, P_{1,3}, fringe) \times \\
 & \sum_{other} \mathbf{P}(P_{1,3}, explored, fringe, other) \\
 & = \alpha \sum_{fringe} \mathbf{P}(b|explored, P_{1,3}, fringe) \times \\
 & \sum_{other} \mathbf{P}(P_{1,3}) P(explored) P(fringe) P(other)
 \end{aligned}$$

Using conditional independence

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

$$\propto \sum_{fringe} P(b|explored, P_{1,3}, fringe) \times \\ \times \sum_{other} P(P_{1,3})P(explored)P(fringe)P(other)$$

Using conditional independence

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

$$\begin{aligned}
 & \propto \sum_{fringe} \mathbf{P}(b|explored, P_{1,3}, fringe) \times \\
 & \times \sum_{other} \mathbf{P}(P_{1,3}) P(explored) P(fringe) P(other) \\
 & = \alpha P(explored) \mathbf{P}(P_{1,3}) \times \\
 & \times \sum_{fringe} \mathbf{P}(b|explored, P_{1,3}, fringe) P(fringe) \sum_{other} P(other)
 \end{aligned}$$

Using conditional independence

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

Reordering
and pushing sums inwards

$$\begin{aligned}
 & \propto \sum_{fringe} \mathbf{P}(b|explored, P_{1,3}, fringe) \times \\
 & \times \sum_{other} \mathbf{P}(P_{1,3}) P(explored) P(fringe) P(other) \\
 & = \alpha P(explored) \mathbf{P}(P_{1,3}) \times \\
 & \times \sum_{fringe} \mathbf{P}(b|explored, P_{1,3}, fringe) P(fringe) \sum_{other} P(other)
 \end{aligned}$$

Using conditional independence

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

$$\propto P(\text{explored})P(P_{1,3}) \times \\ \times \sum_{\text{fringe}} P(b|\text{explored}, P_{1,3}, \text{fringe})P(\text{fringe}) \sum_{\text{other}} P(\text{other})$$

Using conditional independence

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1

$$\begin{aligned}
 &\propto P(\text{explored}) \mathbf{P}(P_{1,3}) \times \\
 &\times \sum_{\text{fringe}} \mathbf{P}(b | \text{explored}, P_{1,3}, \text{fringe}) P(\text{fringe}) \sum_{\text{other}} P(\text{other}) \\
 &= \alpha' \mathbf{P}(P_{1,3}) \sum_{\text{fringe}} \mathbf{P}(b | \text{explored}, P_{1,3}, \text{fringe}) P(\text{fringe})
 \end{aligned}$$


Using conditional independence

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 B OK	2,2	3,2	4,2
1,1 OK	2,1 B OK	3,1	4,1


Simplifying

$$\begin{aligned}
 &\propto P(\text{explored}) \mathbf{P}(P_{1,3}) \times \\
 &\times \sum_{\text{fringe}} \mathbf{P}(b|\text{explored}, P_{1,3}, \text{fringe}) P(\text{fringe}) \sum_{\text{other}} P(\text{other}) \\
 &= \alpha' \mathbf{P}(P_{1,3}) \sum_{\text{fringe}} \mathbf{P}(b|\text{explored}, P_{1,3}, \text{fringe}) P(\text{fringe})
 \end{aligned}$$


Using conditional independence contd.




$$0.2 \times 0.2 = 0.04$$




$$0.2 \times 0.8 = 0.16$$



$$0.8 \times 0.2 = 0.16$$




$$0.2 \times 0.2 = 0.04$$




$$0.2 \times 0.8 = 0.16$$

$P(b|explored, P_{1,3}, fringe)$


Using conditional independence contd.




$$0.2 \times 0.2 = 0.04$$




$$0.2 \times 0.8 = 0.16$$



$$0.8 \times 0.2 = 0.16$$



$$0.2 \times 0.2 = 0.04$$




$$0.2 \times 0.8 = 0.16$$


$P(b|explored, P_{1,3}, fringe)$

- = 1 when the frontier is consistent with the observations
- = 0 otherwise


Using conditional independence contd.




$$0.2 \times 0.2 = 0.04$$




$$0.2 \times 0.8 = 0.16$$



$$0.8 \times 0.2 = 0.16$$



$$0.2 \times 0.2 = 0.04$$



$$0.2 \times 0.8 = 0.16$$

$P(b|explored, P_{1,3}, fringe)$

- = 1 when the frontier is consistent with the observations
- = 0 otherwise

We can sum over the *possible configurations* for the frontier variables that are consistent with the known facts.

Using conditional independence contd.

1,3		
1,2	B	2,2
	OK	
1,1	2,1	3,1
	OK	OK

$$0.2 \times 0.2 = 0.04$$

1,3		
1,2	B	2,2
	OK	
1,1	2,1	3,1
	OK	OK

$$0.2 \times 0.8 = 0.16$$

1,3		
1,2	B	2,2
	OK	
1,1	2,1	3,1
	OK	OK

$$0.8 \times 0.2 = 0.16$$

1,3		
1,2	B	2,2
	OK	
1,1	2,1	3,1
	OK	OK


$$0.2 \times 0.2 = 0.04$$

1,3		
1,2	B	2,2
	OK	
1,1	2,1	3,1
	OK	OK


$$0.2 \times 0.8 = 0.16$$

$$P(P_{1,3} | \text{explored}, b) =$$


Using conditional independence contd.




$$0.2 \times 0.2 = 0.04$$




$$0.2 \times 0.8 = 0.16$$



$$0.8 \times 0.2 = 0.16$$



$$0.2 \times 0.2 = 0.04$$




$$0.2 \times 0.8 = 0.16$$


$P(P_{1,3} | explored, b) =$

$\alpha' \langle 0.2(0.04 + 0.16 + 0.16), 0.8(0.04 + 0.16) \rangle$


Using conditional independence contd.




$$0.2 \times 0.2 = 0.04$$




$$0.2 \times 0.8 = 0.16$$



$$0.8 \times 0.2 = 0.16$$



$$0.2 \times 0.2 = 0.04$$



$$0.2 \times 0.8 = 0.16$$

$P(P_{1,3} | explored, b) =$

$$\alpha' \langle 0.2(0.04 + 0.16 + 0.16), 0.8(0.04 + 0.16) \rangle$$

$$\approx \langle 0.31, 0.69 \rangle$$

Using conditional independence contd.

1,3		
1,2	B	2,2
	OK	
1,1	2,1	3,1
	OK	

$$0.2 \times 0.2 = 0.04$$

1,3		
1,2	B	2,2
	OK	
1,1	2,1	3,1
	OK	

$$0.2 \times 0.8 = 0.16$$

1,3		
1,2	B	2,2
	OK	
1,1	2,1	3,1
	OK	

$$0.8 \times 0.2 = 0.16$$

1,3		
1,2	B	2,2
	OK	
1,1	2,1	3,1
	OK	

$$0.2 \times 0.2 = 0.04$$

1,3		
1,2	B	2,2
	OK	
1,1	2,1	3,1
	OK	

$$0.2 \times 0.8 = 0.16$$


$$\mathbf{P}(P_{1,3} | explored, b) =$$

$$\alpha' \langle 0.2(0.04 + 0.16 + 0.16), 0.8(0.04 + 0.16) \rangle$$


$$\approx \langle 0.31, 0.69 \rangle$$

$$\mathbf{P}(P_{2,2} | explored, b) \approx \langle 0.86, 0.14 \rangle$$


Using conditional independence contd.




$$0.2 \times 0.2 = 0.04$$




$$0.2 \times 0.8 = 0.16$$



$$0.8 \times 0.2 = 0.16$$



$$0.2 \times 0.2 = 0.04$$



$$0.2 \times 0.8 = 0.16$$

$$\mathbf{P}(P_{1,3} | explored, b) =$$

$$\alpha' \langle 0.2(0.04 + 0.16 + 0.16), 0.8(0.04 + 0.16 + 0.16) \rangle$$

$$\approx \langle 0.31, 0.69 \rangle$$

$$\mathbf{P}(P_{2,2} | explored, b) \approx \langle 0.86, 0.14 \rangle$$



Today's class

- Probabilities and conditional probabilities
- Independence and conditional independence
- Estimating chances of possible outcomes

Coming next

- Combining chances and rewards
- Maximising the expected reward