



Partitions

To *partition* a set is to break it up into disjoint nonempty subsets.



Let me show you an important example of a partition on my set of shapes, which will show up later.



Here's the formal definition.

Definition. Let α be a set. A *partition* of α is a set $\mathcal{C} = \{X_1, X_2, X_3, \dots\}$ of subsets of α called *blocks*, satisfying the following axioms:

- 1 Every $X_i \in \mathcal{C}$ is non-empty;
- 2 The union of all the X_i 's is α ;
- 3 Distinct X_i are disjoint. That is if $X_i \cap X_j \neq \emptyset$ then $X_i = X_j$.

Here's the formal definition.

Definition. Let α be a set. A *partition* of α is a set $\mathcal{C} = \{X_1, X_2, X_3, \dots\}$ of subsets of α called *blocks*, satisfying the following axioms:

- 1 Every $X_i \in \mathcal{C}$ is non-empty;
- 2 The union of all the X_i 's is α ;
- 3 Distinct X_i are disjoint. That is if $X_i \cap X_j \neq \emptyset$ then $X_i = X_j$.

Another way of thinking about it – a partition of α is a collection of nonempty subsets of α called blocks with the property that every element of α is in *exactly one* of the blocks.

Here's the formal definition.

Definition. Let α be a set. A *partition* of α is a set $\mathcal{C} = \{X_1, X_2, X_3, \dots\}$ of subsets of α called *blocks*, satisfying the following axioms:

- 1 Every $X_i \in \mathcal{C}$ is non-empty;
- 2 The union of all the X_i 's is α ;
- 3 Distinct X_i are disjoint. That is if $X_i \cap X_j \neq \emptyset$ then $X_i = X_j$.

Another way of thinking about it – a partition of α is a collection of nonempty subsets of α called blocks with the property that every element of α is in *exactly one* of the blocks.

Now let me explain why I have introduced this notion.

Being the same thing

A mathematical idea can sometimes be looked at in two different ways.

Being the same thing

A mathematical idea can sometimes be looked at in two different ways.

Each way tells the truth, but might give different insights.

Being the same thing

A mathematical idea can sometimes be looked at in two different ways.

Each way tells the truth, but might give different insights.

For example, one way of looking at a situation might make a calculation much easier than another way.

Being the same thing

A mathematical idea can sometimes be looked at in two different ways.

Each way tells the truth, but might give different insights.

For example, one way of looking at a situation might make a calculation much easier than another way.

Predicates on a set X and subsets of X are two ways of thinking about the same idea.

Being the same thing

A mathematical idea can sometimes be looked at in two different ways.

Each way tells the truth, but might give different insights.

For example, one way of looking at a situation might make a calculation much easier than another way.

Predicates on a set X and subsets of X are two ways of thinking about the same idea.

A binary relation on X and its graph, a subset of $X \times X$, are two ways of thinking about the same idea.

Being the same thing

A mathematical idea can sometimes be looked at in two different ways.

Each way tells the truth, but might give different insights.

For example, one way of looking at a situation might make a calculation much easier than another way.

Predicates on a set X and subsets of X are two ways of thinking about the same idea.

A binary relation on X and its graph, a subset of $X \times X$, are two ways of thinking about the same idea.

It turns out that equivalence relations and partitions are two ways of thinking about the same idea.


Given a partition of α into blocks, here is how to make an equivalence relation on α :

Given a partition of α into blocks, here is how to make an equivalence relation on α :

We say two elements a, b of α are related if they're in the same block.

Given a partition of α into blocks, here is how to make an equivalence relation on α :


We say two elements a, b of α are related if they're in the same block.



For example, in the picture above, x and y would be related, and so would z and z , but x would not be related to z .

Given a partition of α into blocks, here is how to make an equivalence relation on α :


We say two elements a, b of α are related if they're in the same block.



For example, in the picture above, x and y would be related, and so would z and z , but x would not be related to z .

What are the three things to check to ensure this is an equivalence relation? Can you check them? Pause the video and have a go.

Example: if we consider the partition of my plastic shapes, we see that the corresponding equivalence relation is my favourite equivalence relation! Two shapes are related iff they're the same colour.



Example: we can partition the natural numbers into X_1 , the odd ones, and X_0 the even ones.

Example: we can partition the natural numbers into X_1 , the odd ones, and X_0 the even ones.

The corresponding equivalence relation: two natural numbers are equivalent if and only if they are both odd, or both even.

Example: we can partition the natural numbers into X_1 , the odd ones, and X_0 the even ones.

The corresponding equivalence relation: two natural numbers are equivalent if and only if they are both odd, or both even.

In other words, if they have the same remainder when you divide them by 2.

In this video we learnt about partitions, and how to make an equivalence relation on a set α from a partition on α .

In this video we learnt about partitions, and how to make an equivalence relation on a set α from a partition on α .

In the final video we will explain the reverse construction.

In this video we learnt about partitions, and how to make an equivalence relation on a set α from a partition on α .

In the final video we will explain the reverse construction.

These constructions are the way we pass between our two different ways of understanding the idea of an equivalence relation.