

# INTEGERS: VIDEO XI

## SOME RESULTS ABOUT PRIMES

# Bézout's Identity

## Theorem (Bézout's Identity)

*Let  $a, b \in \mathbb{Z} - \{0\}$ . Then there exist  $x, y \in \mathbb{Z}$ , such that  $ax + by = \gcd(a, b)$ .*

**Idea of Proof:** **Exercise!**

# Bézout's Identity

## Theorem (Bézout's Identity)

*Let  $a, b \in \mathbb{Z} - \{0\}$ . Then there exist  $x, y \in \mathbb{Z}$ , such that  $ax + by = \gcd(a, b)$ .*

### Idea of Proof: **Exercise!**

- Use the **Well-Ordering Principle (WOP)**
  - Look at the set  $S = \{n \in \mathbb{N} \mid n = ax + by > 0, x, y \in \mathbb{Z}\} \subseteq \mathbb{N}$

# Bézout's Identity

## Theorem (Bézout's Identity)

*Let  $a, b \in \mathbb{Z} - \{0\}$ . Then there exist  $x, y \in \mathbb{Z}$ , such that  $ax + by = \gcd(a, b)$ .*

### Idea of Proof: **Exercise!**

- Use the **Well-Ordering Principle (WOP)**
  - Look at the set  $S = \{n \in \mathbb{N} \mid n = ax + by > 0, x, y \in \mathbb{Z}\} \subseteq \mathbb{N}$
  - Show that  $S \neq \emptyset$ .

# Bézout's Identity

## Theorem (Bézout's Identity)

*Let  $a, b \in \mathbb{Z} - \{0\}$ . Then there exist  $x, y \in \mathbb{Z}$ , such that  $ax + by = \gcd(a, b)$ .*

### Idea of Proof: **Exercise!**

- Use the **Well-Ordering Principle (WOP)**
  - Look at the set  $S = \{n \in \mathbb{N} \mid n = ax + by > 0, x, y \in \mathbb{Z}\} \subseteq \mathbb{N}$
  - Show that  $S \neq \emptyset$ .
  - By the WOP, there exists a **least element**  $d = ax_0 + by_0 > 0$ ,  $x_0, y_0 \in \mathbb{Z}$ !

# Bézout's Identity

## Theorem (Bézout's Identity)

*Let  $a, b \in \mathbb{Z} - \{0\}$ . Then there exist  $x, y \in \mathbb{Z}$ , such that  $ax + by = \gcd(a, b)$ .*

### Idea of Proof: **Exercise!**

- Use the **Well-Ordering Principle (WOP)**
  - Look at the set  $S = \{n \in \mathbb{N} \mid n = ax + by > 0, x, y \in \mathbb{Z}\} \subseteq \mathbb{N}$
  - Show that  $S \neq \emptyset$ .
  - By the WOP, there exists a **least element**  $d = ax_0 + by_0 > 0$ ,  $x_0, y_0 \in \mathbb{Z}$ !
- Show that  $d|a$  (similarly  $d|b$ ) using **the Quotient-Remainder Theorem**:
  - By QRT  $a = dq + r$ ,  $0 \leq r < d$

# Bézout's Identity

## Theorem (Bézout's Identity)

Let  $a, b \in \mathbb{Z} - \{0\}$ . Then there exist  $x, y \in \mathbb{Z}$ , such that  $ax + by = \gcd(a, b)$ .

### Idea of Proof: **Exercise!**

- Use the **Well-Ordering Principle (WOP)**
  - Look at the set  $S = \{n \in \mathbb{N} \mid n = ax + by > 0, x, y \in \mathbb{Z}\} \subseteq \mathbb{N}$
  - Show that  $S \neq \emptyset$ .
  - By the WOP, there exists a **least element**  $d = ax_0 + by_0 > 0$ ,  $x_0, y_0 \in \mathbb{Z}$ !
- Show that  $d|a$  (similarly  $d|b$ ) using **the Quotient-Remainder Theorem**:
  - By QRT  $a = dq + r$ ,  $0 \leq r < d$
  - Deduce **plugging in**  $d = ax_0 + by_0$ , that  $r \in S$ , if  $r > 0$ : **contradiction** to  $d$  least element!  
Therefore, we need to have  $r = 0$ .

# Bézout's Identity

## Theorem (Bézout's Identity)

Let  $a, b \in \mathbb{Z} - \{0\}$ . Then there exist  $x, y \in \mathbb{Z}$ , such that  $ax + by = \gcd(a, b)$ .

### Idea of Proof: **Exercise!**

- Use the **Well-Ordering Principle (WOP)**
  - Look at the set  $S = \{n \in \mathbb{N} \mid n = ax + by > 0, x, y \in \mathbb{Z}\} \subseteq \mathbb{N}$
  - Show that  $S \neq \emptyset$ .
  - By the WOP, there exists a **least element**  $d = ax_0 + by_0 > 0$ ,  $x_0, y_0 \in \mathbb{Z}$ !
- Show that  $d|a$  (similarly  $d|b$ ) using **the Quotient-Remainder Theorem**:
  - By QRT  $a = dq + r$ ,  $0 \leq r < d$
  - Deduce **plugging in**  $d = ax_0 + by_0$ , that  $r \in S$ , if  $r > 0$ : **contradiction** to  $d$  least element!  
Therefore, we need to have  $r = 0$ .
- Show finally that  $d$  is the greatest common divisor of  $a$  and  $b$ .



# ...and a useful application

## Theorem (Bézout's Identity)

*Let  $a, b \in \mathbb{Z} - \{0\}$ . Then there exist  $x, y \in \mathbb{Z}$ , such that  $ax + by = \gcd(a, b)$ .*

## Corollary

*Let  $a, b \in \mathbb{Z}$ . If  $n|ab$  for some  $n \in \mathbb{N}$  and  $\gcd(n, a) = 1$ , then  $n|b$ .*

**proof**

# Fundamental Theorem of Arithmetic

## Corollary

*Let  $a, b \in \mathbb{Z}$ . If  $n|ab$  and  $\gcd(n, a) = 1$ , then  $n|b$ .*

## Theorem (Fundamental Theorem of Arithmetic)

*Every integer  $n > 1$  has a **unique prime factorization**.*

# Fundamental Theorem of Arithmetic

## Corollary

Let  $a, b \in \mathbb{Z}$ . If  $n|ab$  and  $\gcd(n, a) = 1$ , then  $n|b$ .

## Theorem (Fundamental Theorem of Arithmetic)

Every integer  $n > 1$  has a *unique prime factorization*.

### Proof:

- We already proved with the Well-Ordering Principle (Video 6):  
All  $n \in \mathbb{N}$ ,  $n > 1$  can be factored by a product of primes!

# Fundamental Theorem of Arithmetic

## Corollary

Let  $a, b \in \mathbb{Z}$ . If  $n|ab$  and  $\gcd(n, a) = 1$ , then  $n|b$ .

## Theorem (Fundamental Theorem of Arithmetic)

Every integer  $n > 1$  has a *unique prime factorization*.

### Proof:

- We already proved with the Well-Ordering Principle (Video 6):  
All  $n \in \mathbb{N}$ ,  $n > 1$  can be factored by a product of primes!
- We just need to prove the uniqueness: **Proof by contradiction!**
  - Assume there are *two distinct* prime factorizations

$$n = p_1 p_2 \dots p_k = q_1 q_2 \dots q_l, \quad k, l \in \mathbb{N}.$$

# Fundamental Theorem of Arithmetic

## Corollary

Let  $a, b \in \mathbb{Z}$ . If  $n|ab$  and  $\gcd(n, a) = 1$ , then  $n|b$ .

## Theorem (Fundamental Theorem of Arithmetic)

Every integer  $n > 1$  has a *unique prime factorization*.

### Proof:

- We already proved with the Well-Ordering Principle (Video 6):  
All  $n \in \mathbb{N}$ ,  $n > 1$  can be factored by a product of primes!
- We just need to prove the uniqueness: **Proof by contradiction!**
  - Assume there are *two distinct* prime factorizations

$$n = p_1 p_2 \dots p_k = q_1 q_2 \dots q_l, \quad k, l \in \mathbb{N}.$$

- Remove all common primes:

$$p_{i_1} p_{i_2} \dots p_{i_m} = q_{j_1} q_{j_2} \dots q_{j_m}.$$

# Fundamental Theorem of Arithmetic

## Corollary

Let  $a, b \in \mathbb{Z}$ . If  $n|ab$  and  $\gcd(n, a) = 1$ , then  $n|b$ .

## Theorem (Fundamental Theorem of Arithmetic)

Every integer  $n > 1$  has a *unique prime factorization*.

### Proof:

- We already proved with the Well-Ordering Principle (Video 6):  
All  $n \in \mathbb{N}$ ,  $n > 1$  can be factored by a product of primes!
- We just need to prove the uniqueness: **Proof by contradiction!**
  - Assume there are *two distinct* prime factorizations

$$n = p_1 p_2 \dots p_k = q_1 q_2 \dots q_l, \quad k, l \in \mathbb{N}.$$

- Remove all common primes:

$$p_{i_1} p_{i_2} \dots p_{i_m} = q_{j_1} q_{j_2} \dots q_{j_m}.$$

Now all  $p_{i_s}$  are different from all  $q_{j_t}$ !

# Fundamental Theorem of Arithmetic

## Corollary

Let  $a, b \in \mathbb{Z}$ . If  $n|ab$  and  $\gcd(n, a) = 1$ , then  $n|b$ .

## Theorem (Fundamental Theorem of Arithmetic)

Every integer  $n > 1$  has a *unique prime factorization*.

### Proof:

- We already proved with the Well-Ordering Principle (Video 6):  
All  $n \in \mathbb{N}$ ,  $n > 1$  can be factored by a product of primes!
- We just need to prove the uniqueness: **Proof by contradiction!**
  - Assume there are *two distinct* prime factorizations

$$n = p_1 p_2 \dots p_k = q_1 q_2 \dots q_l, \quad k, l \in \mathbb{N}.$$

- Remove all common primes:

$$p_{i_1} p_{i_2} \dots p_{i_m} = q_{j_1} q_{j_2} \dots q_{j_m}.$$

Now all  $p_{i_s}$  are different from all  $q_{j_t}$ !

- Obviously  $p_{i_1}$  divides the left handside, hence  $p_{i_1} | q_{j_1} q_{j_2} \dots q_{j_m}$ .

# Fundamental Theorem of Arithmetic

## Corollary

Let  $a, b \in \mathbb{Z}$ . If  $n|ab$  and  $\gcd(n, a) = 1$ , then  $n|b$ .

## Theorem (Fundamental Theorem of Arithmetic)

Every integer  $n > 1$  has a *unique prime factorization*.

### Proof:

- We already proved with the Well-Ordering Principle (Video 6):  
All  $n \in \mathbb{N}$ ,  $n > 1$  can be factored by a product of primes!
- We just need to prove the uniqueness: **Proof by contradiction!**
  - Assume there are *two distinct* prime factorizations

$$n = p_1 p_2 \dots p_k = q_1 q_2 \dots q_l, \quad k, l \in \mathbb{N}.$$

- Remove all common primes:

$$p_{i_1} p_{i_2} \dots p_{i_m} = q_{j_1} q_{j_2} \dots q_{j_m}.$$

Now all  $p_{i_s}$  are different from all  $q_{j_t}$ !

- Obviously  $p_{i_1}$  divides the left handside, hence  $p_{i_1} | q_{j_1} q_{j_2} \dots q_{j_m}$ .
- By above Corollary:  $p_{i_1} | q_{j_t}$ , for some  $j_t$ .

**Contradiction since  $q_{j_t}$  prime!**  $\Rightarrow$  The factorization is unique. □



# Fundamental Theorem of Arithmetic

## Corollary

Let  $a, b \in \mathbb{Z}$ . If  $n|ab$  and  $\gcd(n, a) = 1$ , then  $n|b$ .

## Theorem (Fundamental Theorem of Arithmetic)

Every integer  $n > 1$  has a *unique prime factorization*.

### Proof:

- We already proved with the Well-Ordering Principle (Video 6):  
All  $n \in \mathbb{N}$ ,  $n > 1$  can be factored by a product of primes!
- We just need to prove the uniqueness: **Proof by contradiction!**
  - Assume there are *two distinct* prime factorizations

$$n = p_1 p_2 \dots p_k = q_1 q_2 \dots q_l, \quad k, l \in \mathbb{N}.$$

- Remove all common primes:

$$p_{i_1} p_{i_2} \dots p_{i_m} = q_{j_1} q_{j_2} \dots q_{j_m}.$$

Now all  $p_{i_s}$  are different from all  $q_{j_t}$ !

- Obviously  $p_{i_1}$  divides the left handside, hence  $p_{i_1} | q_{j_1} q_{j_2} \dots q_{j_m}$ .
- By above Corollary:  $p_{i_1} | q_{j_t}$ , for some  $j_t$ .

**Contradiction since  $q_{j_t}$  prime!**  $\Rightarrow$  The factorization is unique.  $\square$

**Prime power factorization:**  $a = p_1^{n_1} p_2^{n_2} \dots p_k^{n_k}$ ,  $n_1 \dots n_k > 0$ ,  $p_1 < p_2 < \dots < p_k$ .

# Applications of the Fundamental Theorem

Theorem (Fundamental Theorem of Arithmetic)

*Every integer  $n > 1$  has a unique prime factorization.*

Theorem ((Infinitude of primes))

*There are infinitely many primes.*

**Proof:**

# Applications of the Fundamental Theorem

## Proposition

Let  $a, b \in \mathbb{N}$  with prime power factorization

$$a = p_1^{n_1} p_2^{n_2} \cdots p_k^{n_k} q_1^{s_1} \cdots q_l^{s_l}, \quad b = p_1^{m_1} p_2^{m_2} \cdots p_k^{m_k} r_1^{t_1} \cdots r_j^{t_j},$$

where

# Applications of the Fundamental Theorem

## Proposition

Let  $a, b \in \mathbb{N}$  with prime power factorization

$$a = p_1^{n_1} p_2^{n_2} \cdots p_k^{n_k} q_1^{s_1} \cdots q_l^{s_l}, \quad b = p_1^{m_1} p_2^{m_2} \cdots p_k^{m_k} r_1^{t_1} \cdots r_j^{t_j},$$

where

- **the primes appearing  $(p_i, q_i, r_i)$  are all distinct**
- **all exponents are positive**
- **we don't require the primes be in increasing order here!**

# Applications of the Fundamental Theorem

## Proposition

Let  $a, b \in \mathbb{N}$  with prime power factorization

$$a = p_1^{n_1} p_2^{n_2} \cdots p_k^{n_k} q_1^{s_1} \cdots q_l^{s_l}, \quad b = p_1^{m_1} p_2^{m_2} \cdots p_k^{m_k} r_1^{t_1} \cdots r_j^{t_j},$$

where

- **the primes appearing ( $p_i, q_i, r_i$ ) are all distinct**
- **all exponents are positive**
- **we don't require the primes be in increasing order here!**

Then

$$\gcd(a, b) = p_1^{\min(n_1, m_1)} p_2^{\min(n_2, m_2)} \cdots p_k^{\min(n_k, m_k)},$$

$$\operatorname{lcm}(a, b) = p_1^{\max(n_1, m_1)} p_2^{\max(n_2, m_2)} \cdots p_k^{\max(n_k, m_k)} q_1^{s_1} \cdots q_l^{s_l} r_1^{t_1} \cdots r_j^{t_j}.$$

**Example:**

# INTEGERS: VIDEO XII

## MODULAR ARITHMETIC

## Definition

Let  $a, b \in \mathbb{Z}$  and  $n \in \mathbb{N}, n > 0$ . We say that  $a$  is congruent to  $b$  modulo  $n$  if  $n \mid (a - b)$ .

**Notation:**  $a \equiv b \pmod{n}$ .

## Definition

Let  $a, b \in \mathbb{Z}$  and  $n \in \mathbb{N}, n > 0$ . We say that  $a$  is congruent to  $b$  modulo  $n$  if  $n \mid (a - b)$ .

**Notation:**  $a \equiv b \pmod{n}$ .

## Proposition

Let  $a, b, c \in \mathbb{Z}$ . Then for  $n \in \mathbb{N}, n > 0$ .

- 1  $a \equiv a \pmod{n}$ , for all  $a \in \mathbb{Z}$ .



## Definition

Let  $a, b \in \mathbb{Z}$  and  $n \in \mathbb{N}, n > 0$ . We say that  $a$  is congruent to  $b$  modulo  $n$  if  $n \mid (a - b)$ .

**Notation:**  $a \equiv b \pmod{n}$ .

## Proposition

Let  $a, b, c \in \mathbb{Z}$ . Then for  $n \in \mathbb{N}, n > 0$ .

- 1  $a \equiv a \pmod{n}$ , for all  $a \in \mathbb{Z}$ .
- 2 If  $a \equiv b \pmod{n}$ , then  $b \equiv a \pmod{n}$ .

## Definition

Let  $a, b \in \mathbb{Z}$  and  $n \in \mathbb{N}, n > 0$ . We say that  $a$  is congruent to  $b$  modulo  $n$  if  $n \mid (a - b)$ .

**Notation:**  $a \equiv b \pmod{n}$ .

## Proposition

Let  $a, b, c \in \mathbb{Z}$ . Then for  $n \in \mathbb{N}, n > 0$ .

- 1  $a \equiv a \pmod{n}$ , for all  $a \in \mathbb{Z}$ .
- 2 If  $a \equiv b \pmod{n}$ , then  $b \equiv a \pmod{n}$ .
- 3 If  $a \equiv b \pmod{n}$  and  $b \equiv c \pmod{n}$ , then  $a \equiv c \pmod{n}$ .

# Modular arithmetic

## Definition

Let  $a, b \in \mathbb{Z}$  and  $n \in \mathbb{N}, n > 0$ . We say that  $a$  is congruent to  $b$  modulo  $n$  if  $n \mid (a - b)$ .

**Notation:**  $a \equiv b \pmod{n}$ .

## Proposition

Let  $a, b, c \in \mathbb{Z}$ . Then for  $n \in \mathbb{N}, n > 0$ .

- 1  $a \equiv a \pmod{n}$ , for all  $a \in \mathbb{Z}$ .
- 2 If  $a \equiv b \pmod{n}$ , then  $b \equiv a \pmod{n}$ .
- 3 If  $a \equiv b \pmod{n}$  and  $b \equiv c \pmod{n}$ , then  $a \equiv c \pmod{n}$ .

★ This means that  $\equiv$  defines an **equivalence relation** on the set of integers!

## Definition

Let  $a, b \in \mathbb{Z}$  and  $n \in \mathbb{N}, n > 0$ . We say that  $a$  is congruent to  $b$  modulo  $n$  if  $n|(a - b)$ .

**Proof of proposition:** Let  $a, b, c \in \mathbb{Z}$ . Then for  $n \in \mathbb{N}, n > 0$ .

- 1 Reflexivity:  $a \equiv a \pmod{n}$ , for all  $a \in \mathbb{Z}$ .
- 2 Symmetry:  $a \equiv b \pmod{n}$ , then  $b \equiv a \pmod{n}$ .
- 3 Transitivity: If  $a \equiv b \pmod{n}$  and  $b \equiv c \pmod{n}$ , then  $a \equiv c \pmod{n}$ .

And we can define **congruence classes**:

$$[a]_n := \{b \in \mathbb{Z} | b \equiv a \pmod{n}\}.$$

# First results

## Proposition

Let  $a, b \in \mathbb{Z}$ ,  $n$  a positive integer. Then  $a \equiv b \pmod n$  if and only if *there exists  $k \in \mathbb{Z}$  such that  $a = b + kn$* .

**Proof:**

**Consequently:**

$$[a]_n := \{a + kn \mid k \in \mathbb{Z}\} = \{\dots a - 2n, a - n, a, a + n, a + 2n, \dots\}$$

## Proposition

Let  $a, b \in \mathbb{Z}$  and  $n$  a positive integer. Then  $a \equiv b \pmod n$  if and only if  *$a$  and  $b$  have the same remainder after division by  $n$* .

**Proof:**

# Congruence Classes

- Keep in mind!  $[a]_n := \{a + kn \mid k \in \mathbb{Z}\} = \{\dots a - 2n, a - n, a, a + n, a + 2n, \dots\}$ .

## Proposition

There are *exactly  $n$  congruence classes modulo  $n$* :  $[0]_n, [1]_n, \dots, [n-1]_n$ .

## Intuition:

- $n=1$ :  $[a]_1 := \{a + k \mid k \in \mathbb{Z}\} = \{\dots a - 2, a - 1, a, a + 1, a + 2, \dots\} = \mathbb{N}$ !

# Congruence Classes

- Keep in mind!  $[a]_n := \{a + kn \mid k \in \mathbb{Z}\} = \{\dots a - 2n, a - n, a, a + n, a + 2n, \dots\}$ .

## Proposition

There are *exactly  $n$  congruence classes modulo  $n$* :  $[0]_n, [1]_n, \dots, [n-1]_n$ .

## Intuition:

- $n=1$ :**  $[a]_1 := \{a + k \mid k \in \mathbb{Z}\} = \{\dots a - 2, a - 1, a, a + 1, a + 2, \dots\} = \mathbb{N}$ !  
All numbers are congruent modulo 1:  $[0]_1 = [1]_1 = [2]_1 = \dots$

$\Rightarrow$  only **one congruence classe mod 1**.

# Congruence Classes

- Keep in mind!  $[a]_n := \{a + kn \mid k \in \mathbb{Z}\} = \{\dots a - 2n, a - n, a, a + n, a + 2n, \dots\}$ .

## Proposition

There are *exactly  $n$  congruence classes modulo  $n$* :  $[0]_n, [1]_n, \dots, [n-1]_n$ .

## Intuition:

- $n=1$ :**  $[a]_1 := \{a + k \mid k \in \mathbb{Z}\} = \{\dots a - 2, a - 1, a, a + 1, a + 2, \dots\} = \mathbb{N}$ !  
All numbers are congruent modulo 1:  $[0]_1 = [1]_1 = [2]_1 = \dots$

$\Rightarrow$  only **one congruence classe mod 1**.

- $n=2$ :**  $[a]_2 := \{a + 2k \mid k \in \mathbb{Z}\} = \{\dots a - 4, a - 2, a, a + 2, a + 4, \dots\}$



# Congruence Classes

- Keep in mind!  $[a]_n := \{a + kn \mid k \in \mathbb{Z}\} = \{\dots a - 2n, a - n, a, a + n, a + 2n, \dots\}$ .

## Proposition

There are *exactly  $n$  congruence classes modulo  $n$* :  $[0]_n, [1]_n, \dots, [n-1]_n$ .

### Intuition:

- $n=1$ :**  $[a]_1 := \{a + k \mid k \in \mathbb{Z}\} = \{\dots a - 2, a - 1, a, a + 1, a + 2, \dots\} = \mathbb{N}$ !  
All numbers are congruent modulo 1:  $[0]_1 = [1]_1 = [2]_1 = \dots$

$\Rightarrow$  only **one congruence class mod 1**.

- $n=2$ :**  $[a]_2 := \{a + 2k \mid k \in \mathbb{Z}\} = \{\dots a - 4, a - 2, a, a + 2, a + 4, \dots\}$   
Numbers of the same class have same parity:  $[0]_2 = [2]_2 = [4]_2 = \dots$  and  $[1]_2 = [3]_2 = [5]_2 = \dots$

$\Rightarrow$  **two congruence classes mod 2**.

# Congruence Classes

- Keep in mind!  $[a]_n := \{a + kn \mid k \in \mathbb{Z}\} = \{\dots a - 2n, a - n, a, a + n, a + 2n, \dots\}$ .

## Proposition

There are *exactly  $n$  congruence classes modulo  $n$* :  $[0]_n, [1]_n, \dots, [n-1]_n$ .

### Intuition:

- $n=1$ :**  $[a]_1 := \{a + k \mid k \in \mathbb{Z}\} = \{\dots a - 2, a - 1, a, a + 1, a + 2, \dots\} = \mathbb{N}$ !  
All numbers are congruent modulo 1:  $[0]_1 = [1]_1 = [2]_1 = \dots$

$\Rightarrow$  only **one congruence class mod 1**.

- $n=2$ :**  $[a]_2 := \{a + 2k \mid k \in \mathbb{Z}\} = \{\dots a - 4, a - 2, a, a + 2, a + 4, \dots\}$   
Numbers of the same class have same parity:  $[0]_2 = [2]_2 = [4]_2 = \dots$  and  $[1]_2 = [3]_2 = [5]_2 = \dots$

$\Rightarrow$  **two congruence classes mod 2**.

## Proposition

There are *exactly  $n$  congruence classes modulo  $n$* :  $[0]_n, [1]_n, \dots, [n-1]_n$ .

### Proof of Proposition:

- Show that  $[0]_n, [1]_n, \dots, [n-1]_n$  are all different!
- Show that  $[0]_n, [1]_n, \dots, [n-1]_n$  are the only possible classes!

## Definition

A **complete system of residues modulo  $n$**  is a set of integers such that every integer is congruent modulo  $n$  to exactly one integer in the set.

### Example:

# The set of congruence classes modulo $n$ : $\mathbb{Z}_n$

## Definition

$$\mathbb{Z}_n := \{[0]_n, [1]_n, \dots, [n-1]_n\}$$

We want to define operations on our new set of equivalence classes!

Let's just do it the simplest possible way!

# The set of congruence classes modulo $n$ : $\mathbb{Z}_n$

## Definition

$$\mathbb{Z}_n := \{[0]_n, [1]_n, \dots, [n-1]_n\}$$

We want to define operations on our new set of equivalence classes!

Let's just do it the simplest possible way!

$$+ : \mathbb{Z}_n \times \mathbb{Z}_n \rightarrow \mathbb{Z}_n, ([a]_n, [b]_n) \mapsto [a + b]_n$$

$$\cdot : \mathbb{Z}_n \times \mathbb{Z}_n \rightarrow \mathbb{Z}_n, ([a]_n, [b]_n) \mapsto [ab]_n.$$

# The set of congruence classes modulo $n$ : $\mathbb{Z}_n$

## Definition

$$\mathbb{Z}_n := \{[0]_n, [1]_n, \dots, [n-1]_n\}$$

We want to define operations on our new set of equivalence classes!

Let's just do it the simplest possible way!

$$+ : \mathbb{Z}_n \times \mathbb{Z}_n \rightarrow \mathbb{Z}_n, ([a]_n, [b]_n) \mapsto [a + b]_n$$

$$\cdot : \mathbb{Z}_n \times \mathbb{Z}_n \rightarrow \mathbb{Z}_n, ([a]_n, [b]_n) \mapsto [ab]_n.$$

Are we done so easily?

# The set of congruence classes modulo $n$ : $\mathbb{Z}_n$

## Definition

$$\mathbb{Z}_n := \{[0]_n, [1]_n, \dots, [n-1]_n\}$$

We want to define operations on our new set of equivalence classes!

Let's just do it the simplest possible way!

$$+ : \mathbb{Z}_n \times \mathbb{Z}_n \rightarrow \mathbb{Z}_n, ([a]_n, [b]_n) \mapsto [a + b]_n$$

$$\cdot : \mathbb{Z}_n \times \mathbb{Z}_n \rightarrow \mathbb{Z}_n, ([a]_n, [b]_n) \mapsto [ab]_n.$$

Are we done so easily? NO!!!! WE NEED TO CHECK that the operations are independent of the representatives of the class!

# The set of congruence classes modulo $n$ : $\mathbb{Z}_n$

## Definition

$$\mathbb{Z}_n := \{[0]_n, [1]_n, \dots, [n-1]_n\}$$

We want to define operations on our new set of equivalence classes!

Let's just do it the simplest possible way!

$$+ : \mathbb{Z}_n \times \mathbb{Z}_n \rightarrow \mathbb{Z}_n, ([a]_n, [b]_n) \mapsto [a + b]_n$$

$$\cdot : \mathbb{Z}_n \times \mathbb{Z}_n \rightarrow \mathbb{Z}_n, ([a]_n, [b]_n) \mapsto [ab]_n.$$

Are we done so easily? NO!!!! WE NEED TO CHECK that the operations are independent of the representatives of the class!

## Lemma

Suppose that  $a, a', b, b' \in \mathbb{Z}$ , such that  $[a]_n = [a']_n$  and  $[b]_n = [b']_n$ . Then

$$1) [a + b]_n = [a' + b']_n \qquad 2) [ab]_n = [a'b']_n.$$

Proof (of 1)):