

Binary relations

✓	✗	✗
✓	✗	✓
✗	✗	✗

A binary relation on a set X is a predicate on $X \times X$.

A binary relation on a set X is a predicate on $X \times X$.

To put it another way, a binary relation R on a set X is a function, which takes as input an ordered pair of elements (a, b) of X , and returns a true-false statement $R(a, b)$.

A binary relation on a set X is a predicate on $X \times X$.

To put it another way, a binary relation R on a set X is a function, which takes as input an ordered pair of elements (a, b) of X , and returns a true-false statement $R(a, b)$.

If P is a predicate on $X \times X$, and if $a, b \in X$, then we say a and b are *related* by the relation if $P(a, b)$ is `true`.

Let's look at some examples.

We all know that $2 + 2 = 4$ and that $2 < 3$.

We all know that $2 + 2 = 4$ and that $2 < 3$.

We know that 2 and 3 and 4 are real numbers.

We all know that $2 + 2 = 4$ and that $2 < 3$.

We know that 2 and 3 and 4 are real numbers.

We know that $+$ is a function $\mathbb{R}^2 \rightarrow \mathbb{R}$.

We all know that $2 + 2 = 4$ and that $2 < 3$.

We know that 2 and 3 and 4 are real numbers.

We know that $+$ is a function $\mathbb{R}^2 \rightarrow \mathbb{R}$.

The other two things, $=$ and $<$, are binary relations on \mathbb{R} .

We all know that $2 + 2 = 4$ and that $2 < 3$.

We know that 2 and 3 and 4 are real numbers.

We know that $+$ is a function $\mathbb{R}^2 \rightarrow \mathbb{R}$.

The other two things, $=$ and $<$, are binary relations on \mathbb{R} .

Let's think a little about why this is the case.

Say we have two real numbers a and b . Then $a < b$ is a true-false statement.

Say we have two real numbers a and b . Then $a < b$ is a true-false statement.

And $a = b$ is also a true-false statement.

Say we have two real numbers a and b . Then $a < b$ is a true-false statement.

And $a = b$ is also a true-false statement.

So $=$ and $<$ are functions which take in two real numbers and output true-false statements.

Say we have two real numbers a and b . Then $a < b$ is a true-false statement.

And $a = b$ is also a true-false statement.

So $=$ and $<$ are functions which take in two real numbers and output true-false statements.

So they're predicates on $\mathbb{R}^2 = \mathbb{R} \times \mathbb{R}$.


Say we have two real numbers a and b . Then $a < b$ is a true-false statement.

And $a = b$ is also a true-false statement.


So $=$ and $<$ are functions which take in two real numbers and output true-false statements.

So they're predicates on $\mathbb{R}^2 = \mathbb{R} \times \mathbb{R}$.

So by definition they're binary relations on \mathbb{R} .




The picture represents a subset of \mathbb{R}^2 .



The picture represents a subset of \mathbb{R}^2 .

For some pairs (a, b) of real numbers, the corresponding coordinate is coloured in, i.e., part of the picture of the cat.




The picture represents a subset of \mathbb{R}^2 .

For some pairs (a, b) of real numbers, the corresponding coordinate is coloured in, i.e., part of the picture of the cat.

Let $P(a, b)$ be `true` if (a, b) is part of the cat, and `false` otherwise.

In other words, let $P(a, b)$ be the true-false statement “ (a, b) is part of the cat”.



The picture represents a subset of \mathbb{R}^2 .

For some pairs (a, b) of real numbers, the corresponding coordinate is coloured in, i.e., part of the picture of the cat.

Let $P(a, b)$ be `true` if (a, b) is part of the cat, and `false` otherwise.

In other words, let $P(a, b)$ be the true-false statement “ (a, b) is part of the cat”.

Then P is a predicate on \mathbb{R}^2 so it's a binary relation on \mathbb{R} .

The graph of a binary relation

Recall that the idea of a subset of α and a predicate on α are two different ways of thinking about the same data.

The graph of a binary relation

Recall that the idea of a subset of α and a predicate on α are two different ways of thinking about the same data.

A binary relation on X is a predicate on X^2 .

The graph of a binary relation

Recall that the idea of a subset of α and a predicate on α are two different ways of thinking about the same data.

A binary relation on X is a predicate on X^2 .

The *graph* of a binary relation is the subset of X^2 associated to this predicate.

That cat drawing didn't obey the vertical line test!

That cat drawing didn't obey the vertical line test!

But that's OK – relations are more general than functions.

Click [here](#) to see a mathoverflow post where someone puts in a lot of work to come up with a formula for a nice pictorial relation.

Counting

The final thing in this video: how many binary relations are there on a set with n elements?

Counting

The final thing in this video: how many binary relations are there on a set with n elements?

Pause the video and see if you can work it out.

Counting

The final thing in this video: how many binary relations are there on a set with n elements?

Pause the video and see if you can work it out.

A binary relation on X is a true-false statement for every element of X^2 .

When we're counting, we think of propositions as extensional – a proposition is determined by its truth value.

Counting

The final thing in this video: how many binary relations are there on a set with n elements?

Pause the video and see if you can work it out.

A binary relation on X is a true-false statement for every element of X^2 .

When we're counting, we think of propositions as extensional – a proposition is determined by its truth value.

So for every element of X^2 (i.e., for every pair of elements of X) we need to decide whether the relation is `true` or `false`.

Counting

The final thing in this video: how many binary relations are there on a set with n elements?

Pause the video and see if you can work it out.

A binary relation on X is a true-false statement for every element of X^2 .

When we're counting, we think of propositions as extensional – a proposition is determined by its truth value.

So for every element of X^2 (i.e., for every pair of elements of X) we need to decide whether the relation is `true` or `false`.

We have two choices of answer for each of n^2 questions.

Counting

The final thing in this video: how many binary relations are there on a set with n elements?

Pause the video and see if you can work it out.

A binary relation on X is a true-false statement for every element of X^2 .

When we're counting, we think of propositions as extensional – a proposition is determined by its truth value.

So for every element of X^2 (i.e., for every pair of elements of X) we need to decide whether the relation is `true` or `false`.

We have two choices of answer for each of n^2 questions.

So the answer is 2^{n^2} .

Counting

The final thing in this video: how many binary relations are there on a set with n elements?

Pause the video and see if you can work it out.

A binary relation on X is a true-false statement for every element of X^2 .

When we're counting, we think of propositions as extensional – a proposition is determined by its truth value.

So for every element of X^2 (i.e., for every pair of elements of X) we need to decide whether the relation is `true` or `false`.

We have two choices of answer for each of n^2 questions.

So the answer is 2^{n^2} .

NB does that mean $2^{(n^2)}$ or $(2^n)^2$?