

# Partial orders

A *partial order* on a set  $X$  is a binary relation which is all three of

- 1 reflexive;
- 2 antisymmetric;
- 3 transitive.

The standard example is  $\leq$  on the natural numbers or the real numbers.

Pause the video and check that you can prove that  $\leq$  is a partial order, or at least reduce it to statements which could be assumed without question at school.

Here's the surprise though. There are other examples of partial orders which look quite different to the real numbers.

Here's the surprise though. There are other examples of partial orders which look quite different to the real numbers.

Let  $\alpha$  be a set and consider the set  $\omega$  of subsets of  $\alpha$ . Then  $\subseteq$  is a binary relation on  $\omega$ . What does this mean? Pause the video and try and figure out what I just meant.

Here's the surprise though. There are other examples of partial orders which look quite different to the real numbers.

Let  $\alpha$  be a set and consider the set  $\omega$  of subsets of  $\alpha$ . Then  $\subseteq$  is a binary relation on  $\omega$ . What does this mean? Pause the video and try and figure out what I just meant.

It means that if  $X, Y \in \omega$  then  $X \subseteq Y$  should be a true-false statement.

Here's the surprise though. There are other examples of partial orders which look quite different to the real numbers.

Let  $\alpha$  be a set and consider the set  $\omega$  of subsets of  $\alpha$ . Then  $\subseteq$  is a binary relation on  $\omega$ . What does this mean? Pause the video and try and figure out what I just meant.

It means that if  $X, Y \in \omega$  then  $X \subseteq Y$  should be a true-false statement.

Now I claim that  $\subseteq$  is a partial order on  $\omega$ . What does that mean? Let's take a look.

Looking at the definition, we see that we need to prove that the binary relation  $\subseteq$  on the set  $\omega$  is reflexive, antisymmetric and transitive. So what does that mean?

Looking at the definition, we see that we need to prove that the binary relation  $\subseteq$  on the set  $\omega$  is reflexive, antisymmetric and transitive. So what does that mean?

$$\forall X, X \subseteq X;$$

$$\forall X Y, (X \subseteq Y \wedge Y \subseteq X) \implies X = Y;$$

$$\forall X Y Z, (X \subseteq Y \wedge Y \subseteq Z) \implies X \subseteq Z;$$

Three levels of a computer game.

Here  $X, Y, Z$  are running through all subsets of  $\alpha$ .

Those statements about sets correspond to the following three statements of logic:

$$P \implies P;$$

$$(P \implies Q \wedge Q \implies P) \implies (P \iff Q);$$

$$(P \implies Q \wedge Q \implies R) \implies (P \implies R).$$

All three can be checked by trying all the cases.



Those statements about sets correspond to the following three statements of logic:

$$P \implies P;$$

$$(P \implies Q \wedge Q \implies P) \implies (P \iff Q);$$

$$(P \implies Q \wedge Q \implies R) \implies (P \implies R).$$

All three can be checked by trying all the cases.


Or you can prove them constructively if you find it fun.

So  $\subseteq$  is a partial order on the set of subsets of  $\alpha$ .

So  $\subseteq$  is a partial order on the set of subsets of  $\alpha$ .

And this is funny because if  $X$  and  $Y$  are random subsets of  $\alpha$  then probably both  $X \subseteq Y$  and  $Y \subseteq X$  are false!

Antisymmetry doesn't mind that.



If you don't like the possibility of “incomparable objects”, you might prefer to study *total orders*. A total order is a partial order (let's call it  $\leq$ ) on a set  $X$  satisfying the additional axiom of “totality”:

$$\forall a, b \in X, a \leq b \vee b \leq a.$$

The set of subsets of  $\alpha$  usually does not have this property.

If you don't like the possibility of “incomparable objects”, you might prefer to study *total orders*. A total order is a partial order (let's call it  $\leq$ ) on a set  $X$  satisfying the additional axiom of “totality”:

$$\forall a, b \in X, a \leq b \vee b \leq a.$$

The set of subsets of  $\alpha$  usually does not have this property.

Can you give an example of  $\alpha$  for which  $\subseteq$  is a total order on the set of subsets of  $\alpha$ ?

The naturals, integers, rationals and reals are all totally ordered, but the complexes have no natural total order.

The naturals, integers, rationals and reals are all totally ordered, but the complexes have no natural total order.

I'll finish this video by mentioning an even stronger condition: A *well-ordered set* is a set  $X$  equipped with a total order  $\leq$  which furthermore satisfies that there are no infinite strictly decreasing sequences  $x_1 > x_2 > x_3 > x_4 > \dots$ . Here  $a > b$  is defined to mean  $\neg(a \leq b)$ .

Consequence: every non-empty subset has a least element.

The naturals, integers, rationals and reals are all totally ordered, but the complexes have no natural total order.

I'll finish this video by mentioning an even stronger condition: A *well-ordered set* is a set  $X$  equipped with a total order  $\leq$  which furthermore satisfies that there are no infinite strictly decreasing sequences  $x_1 > x_2 > x_3 > x_4 > \dots$ . Here  $a > b$  is defined to mean  $\neg(a \leq b)$ .

Consequence: every non-empty subset has a least element.

The natural numbers are an example.



The naturals, integers, rationals and reals are all totally ordered, but the complexes have no natural total order.

I'll finish this video by mentioning an even stronger condition: A *well-ordered set* is a set  $X$  equipped with a total order  $\leq$  which furthermore satisfies that there are no infinite strictly decreasing sequences  $x_1 > x_2 > x_3 > x_4 > \dots$ . Here  $a > b$  is defined to mean  $\neg(a \leq b)$ .

Consequence: every non-empty subset has a least element.

The natural numbers are an example.

It's why induction works.

The naturals, integers, rationals and reals are all totally ordered, but the complexes have no natural total order.

I'll finish this video by mentioning an even stronger condition: A *well-ordered set* is a set  $X$  equipped with a total order  $\leq$  which furthermore satisfies that there are no infinite strictly decreasing sequences  $x_1 > x_2 > x_3 > x_4 > \dots$ . Here  $a > b$  is defined to mean  $\neg(a \leq b)$ .

Consequence: every non-empty subset has a least element.

The natural numbers are an example.

It's why induction works.

Next time: equivalence relations.