

# INTEGERS: VIDEO VIII

## MOTIVATION AND FIRST DEFINITIONS

# Motivation for the integers

**Important material:**

**EQUIVALENCE RELATIONS, EQUIVALENCE CLASSES, DIVISORS**

# Motivation for the integers

**Important material:**

**EQUIVALENCE RELATIONS, EQUIVALENCE CLASSES, DIVISORS**

- Until now we only have addition and multiplication...

# Motivation for the integers

**Important material:**

**EQUIVALENCE RELATIONS, EQUIVALENCE CLASSES, DIVISORS**

- Until now we only have addition and multiplication...
- **So how do we solve for  $x$  equations like**

$$x + 4 = 2?$$

**We want to have " $x = 2 - 4$ "!**

# Motivation for the integers

**Important material:**

**EQUIVALENCE RELATIONS, EQUIVALENCE CLASSES, DIVISORS**

- Until now we only have addition and multiplication...
- **So how do we solve for  $x$  equations like**

$$x + 4 = 2?$$

**We want to have " $x = 2 - 4$ "!**

Idea: define " $2 - 4$ " as the ordered pair  $(2, 4) \in \mathbb{N} \times \mathbb{N}$  satisfying certain rules?

# Motivation for the integers

**Important material:**

**EQUIVALENCE RELATIONS, EQUIVALENCE CLASSES, DIVISORS**

- Until now we only have addition and multiplication...
- **So how do we solve for  $x$  equations like**

$$x + 4 = 2?$$

**We want to have " $x = 2 - 4$ "!**

Idea: define " $2 - 4$ " as the ordered pair  $(2, 4) \in \mathbb{N} \times \mathbb{N}$  satisfying certain rules?

Problem:  $2 - 4 = 3 - 5 = 17 - 19 = \dots \Rightarrow (2, 4) = (3, 5) = (17, 19)$  so it is not unique.

# Motivation for the integers

**Important material:**

**EQUIVALENCE RELATIONS, EQUIVALENCE CLASSES, DIVISORS**

- Until now we only have addition and multiplication...
- **So how do we solve for  $x$  equations like**

$$x + 4 = 2?$$

**We want to have " $x = 2 - 4$ "!**

Idea: define " $2 - 4$ " as the ordered pair  $(2, 4) \in \mathbb{N} \times \mathbb{N}$  satisfying certain rules?

Problem:  $2 - 4 = 3 - 5 = 17 - 19 = \dots \Rightarrow (2, 4) = (3, 5) = (17, 19)$  so it is not unique.

★ Note:  $a - b = c - d \Leftrightarrow a + d = b + c$  ★

What would be your next idea....?

# Definition of the integers

New Idea:

**Define the equivalence relation** on  $\mathbb{N} \times \mathbb{N}$

$$(a, b) \sim (c, d) \text{ if and only if } a + d = b + c.$$



# Definition of the integers

New Idea:

**Define the equivalence relation** on  $\mathbb{N} \times \mathbb{N}$

$$(a, b) \sim (c, d) \text{ if and only if } a + d = b + c.$$

$$a - b := cl((a, b)) = \{(x, y) \in \mathbb{N} \times \mathbb{N} \mid (x, y) \sim (a, b)\}$$

# Definition of the integers

New Idea:

**Define the equivalence relation** on  $\mathbb{N} \times \mathbb{N}$

$$(a, b) \sim (c, d) \text{ if and only if } a + d = b + c.$$

$$a - b := cl((a, b)) = \{(x, y) \in \mathbb{N} \times \mathbb{N} \mid (x, y) \sim (a, b)\}$$

**$\mathbb{Z}$  is the set of all equivalence classes  $a - b$ .**

# Definition of the integers

New Idea:

**Define the equivalence relation** on  $\mathbb{N} \times \mathbb{N}$

$$(a, b) \sim (c, d) \text{ if and only if } a + d = b + c.$$

$$a - b := cl((a, b)) = \{(x, y) \in \mathbb{N} \times \mathbb{N} \mid (x, y) \sim (a, b)\}$$

**$\mathbb{Z}$  is the set of all equivalence classes  $a - b$ .**

Equivalence classes  $a - 0$  will be written  $a$ ,  $0 - b$  will be written  $-b$

# Definition of the integers

New Idea:

**Define the equivalence relation** on  $\mathbb{N} \times \mathbb{N}$

$$(a, b) \sim (c, d) \text{ if and only if } a + d = b + c.$$

$$a - b := cl((a, b)) = \{(x, y) \in \mathbb{N} \times \mathbb{N} \mid (x, y) \sim (a, b)\}$$

$\mathbb{Z}$  is the set of all equivalence classes  $a - b$ .

Equivalence classes  $a - 0$  will be written  $a$ ,  $0 - b$  will be written  $-b$

★ **Wanted:**

$$\begin{aligned}(a - b) + (c - d) &= (a + c) - (b + d), \\ (a - b)(c - d) &= (ac + bd) - (ad + bc).\end{aligned}$$

We define on  $\mathbb{Z}$ :

**Addition:**  $(a, b) + (c, d) := (a + c, b + d)$

**Multiplication:**  $(a, b) \cdot (c, d) := (ac + bd, ad + bc).$

# To go further

Beyond the scope of this lecture (but in the notes) we can show

- Addition and multiplication are well-defined on  $\mathbb{Z}$  (i.e. do not depend on the representant of the class).
- Any element in  $\mathbb{Z}$  is either an  $n \in \mathbb{N}$  or  $-n$ , with  $n \in \mathbb{N}$ . Therefore

$$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$$

**FROM NOW ON WE CAN USE WHATEVER RULES WE KNOW FOR ADDITION, SUBTRACTION, MULTIPLICATION ON THE INTEGERS! (Yippie)**

# INTEGERS: VIDEO IX

## GCD (hcf) /LCM AND QUOTIENT-REMAINDER THEOREM

# GCD(HCF)/LCM

Reminder (but for integers!):

## Definition

Let  $n, m \in \mathbb{Z}$ . We say that  $m$  divides  $n$  if and only if there exists a number  $k \in \mathbb{Z}$ , such that  $n = k \cdot m$ . Notation:  $m|n$ .

# GCD(HCF)/LCM

Reminder (but for integers!):

## Definition

Let  $n, m \in \mathbb{Z}$ . We say that  $m$  divides  $n$  if and only if there exists a number  $k \in \mathbb{Z}$ , such that  $n = k \cdot m$ . Notation:  $m|n$ .

## Definition

Let  $a, b \in \mathbb{Z}$

- A **common divisor** of  $a$  and  $b$  is an integer  $n \in \mathbb{Z}$  such that  $n|a$  and  $n|b$ .



# GCD(HCF)/LCM

Reminder (but for integers!):

## Definition

Let  $n, m \in \mathbb{Z}$ . We say that  $m$  divides  $n$  if and only if there exists a number  $k \in \mathbb{Z}$ , such that  $n = k \cdot m$ . Notation:  $m|n$ .

## Definition

Let  $a, b \in \mathbb{Z}$

- A **common divisor** of  $a$  and  $b$  is an integer  $n \in \mathbb{Z}$  such that  $n|a$  and  $n|b$ .

The **greatest common divisor (gcd)** of  $a$  and  $b$  is the largest such integer. Notation:  $\gcd(a,b)$ .

# GCD(HCF)/LCM

Reminder (but for integers!):

## Definition

Let  $n, m \in \mathbb{Z}$ . We say that  $m$  divides  $n$  if and only if there exists a number  $k \in \mathbb{Z}$ , such that  $n = k \cdot m$ . Notation:  $m|n$ .

## Definition

Let  $a, b \in \mathbb{Z}$

- A **common divisor** of  $a$  and  $b$  is an integer  $n \in \mathbb{Z}$  such that  $n|a$  and  $n|b$ .

The **greatest common divisor (gcd)** of  $a$  and  $b$  is the largest such integer. Notation:  $\gcd(a,b)$ .

- A **common multiple** of  $a$  and  $b$  is an integer  $n \in \mathbb{Z}$  such that  $n = ka$  and  $n = lb$  for some  $k, l \in \mathbb{Z}$ .

# GCD(HCF)/LCM

Reminder (but for integers!):

## Definition

Let  $n, m \in \mathbb{Z}$ . We say that  $m$  divides  $n$  if and only if there exists a number  $k \in \mathbb{Z}$ , such that  $n = k \cdot m$ . Notation:  $m|n$ .

## Definition

Let  $a, b \in \mathbb{Z}$

- A **common divisor** of  $a$  and  $b$  is an integer  $n \in \mathbb{Z}$  such that  $n|a$  and  $n|b$ .

The **greatest common divisor (gcd)** of  $a$  and  $b$  is the largest such integer. Notation:  $\gcd(a,b)$ .

- A **common multiple** of  $a$  and  $b$  is an integer  $n \in \mathbb{Z}$  such that  $n = ka$  and  $n = lb$  for some  $k, l \in \mathbb{Z}$ .

The **least common multiple (lcm)** is the smallest positive such integer. Notation:  $\text{lcm}(a,b)$ .

# GCD(HCF)/LCM

Reminder (but for integers!):

## Definition

Let  $n, m \in \mathbb{Z}$ . We say that  $m$  divides  $n$  if and only if there exists a number  $k \in \mathbb{Z}$ , such that  $n = k \cdot m$ . Notation:  $m|n$ .

## Definition

Let  $a, b \in \mathbb{Z}$

- A **common divisor** of  $a$  and  $b$  is an integer  $n \in \mathbb{Z}$  such that  $n|a$  and  $n|b$ .

The **greatest common divisor (gcd)** of  $a$  and  $b$  is the largest such integer. Notation:  $\gcd(a,b)$ .

- A **common multiple** of  $a$  and  $b$  is an integer  $n \in \mathbb{Z}$  such that  $n = ka$  and  $n = lb$  for some  $k, l \in \mathbb{Z}$ .

The **least common multiple (lcm)** is the smallest positive such integer. Notation:  $\text{lcm}(a,b)$ .

- $a$  and  $b$  are called **relatively prime** if  $\gcd(a, b) = 1$ .

By convention  $\gcd(0,0) = 0$

# About the gcd

Compute  $\gcd(8, 12)$  :

## Proposition

*Suppose  $a, b \in \mathbb{Z} - \{0\}$ . Then*

- a)  $\gcd(a, b) = \gcd(b, a)$ .*
- b) if  $a > 0$  and  $a|b$  then  $\gcd(a, b) = a$*
- c) if  $a = bq + r$ , for some  $q, r \in \mathbb{Z}$  then  $\gcd(a, b) = \gcd(r, b)$ .*

**Proof of c):**

# Quotient-Remainder theorem (Q-R T)

## Theorem

*Let  $a, b \in \mathbb{Z}$ ,  $b > 0$ . Then there exist unique integers  $q$  and  $r$  such that*

$$a = qb + r, \quad 0 \leq r < b$$

Examples: if  $a = 11$ ,  $b = 4$ ,  $11 = 2 \cdot 4 + 3$ , hence  $q = 2$ ,  $r = 3$

if  $a = -8$ ,  $b = 3$ ,  $-8 = -3 \cdot 3 + 1$ , hence  $q = -3$ ,  $r = 1$ .

★ : This result comes directly from long division of  $a$  by  $b$ , where  $q$  is the quotient and  $r$  the remainder.

# Quotient-Remainder theorem (Q-R T)–Proof of existence

## Theorem

Let  $a, b \in \mathbb{Z}$ ,  $b > 0$ . Then there **exists** unique **integers  $q$  and  $r$**  such that

$$a = qb + r, \quad 0 \leq r < b$$

**Proof:** To show:  $\exists r$ , such that  $r = a - qb$  and  $0 \leq r < b$ , for fixed  $a$  and  $b > 0$  and some  $q$

# Quotient-Remainder theorem (Q-R T)–Proof of existence

## Theorem

Let  $a, b \in \mathbb{Z}$ ,  $b > 0$ . Then there **exists** unique **integers  $q$  and  $r$**  such that

$$a = qb + r, \quad 0 \leq r < b$$

**Proof:** To show:  $\exists r$ , such that  $r = a - qb$  and  $0 \leq r < b$ , for fixed  $a$  and  $b > 0$  and some  $q$  **Idea:** Use the Well-Ordering principle (WOP)!

- Define the appropriate set  $S \subseteq \mathbb{N}$ :
- Show  $S$  is not empty:
- By WOP  $S$  has a **least element**  $r$ , such that  $r = a - qb$  and  $r \geq 0$
- It remains to show that  $r < b$

.....and uniqueness of  $r$  (read your notes!)



# INTEGERS: VIDEO X EUCLIDEAN ALGORITHM

# Euclidean algorithm to compute $\gcd(a, b)$

Reminder:

## Theorem (Q-R Theorem)

Let  $a, b \in \mathbb{Z}$ ,  $b > 0$ . Then there **exists** unique **integers  $q$  and  $r$**  such that  
 $a = qb + r$ ,  $0 \leq r < b$

# Euclidean algorithm to compute $\gcd(a, b)$

Reminder:

## Theorem (Q-R Theorem)

Let  $a, b \in \mathbb{Z}$ ,  $b > 0$ . Then there **exists** unique **integers  $q$  and  $r$**  such that  $a = qb + r$ ,  $0 \leq r < b$

## Proposition

Suppose  $a, b \in \mathbb{Z}$ ,  $b \neq 0$ . If  $a = bq + r$ , for some  $q, r \in \mathbb{Z}$  then  $\gcd(a, b) = \gcd(r, b)$ .

# Euclidean algorithm to compute $\gcd(a, b)$

Reminder:

## Theorem (Q-R Theorem)

Let  $a, b \in \mathbb{Z}$ ,  $b > 0$ . Then there **exists** unique **integers  $q$  and  $r$**  such that  $a = qb + r$ ,  $0 \leq r < b$

## Proposition

Suppose  $a, b \in \mathbb{Z}$ ,  $b \neq 0$ . If  $a = bq + r$ , for some  $q, r \in \mathbb{Z}$  then  $\gcd(a, b) = \gcd(r, b)$ .

**Euclidean Algorithm:** How to compute  $\gcd(a, b)$ ,  $a, b \in \mathbb{Z}$ ,  $b > 0$ ?

- Start with  $\gcd(a, b)$ . (We can assume Wlog that  $a > b$ )

# Euclidean algorithm to compute $\gcd(a, b)$

Reminder:

## Theorem (Q-R Theorem)

Let  $a, b \in \mathbb{Z}$ ,  $b > 0$ . Then there **exists** unique **integers  $q$  and  $r$**  such that  $a = qb + r$ ,  $0 \leq r < b$

## Proposition

Suppose  $a, b \in \mathbb{Z}$ ,  $b \neq 0$ . If  $a = bq + r$ , for some  $q, r \in \mathbb{Z}$  then  $\gcd(a, b) = \gcd(r, b)$ .

**Euclidean Algorithm:** How to compute  $\gcd(a, b)$ ,  $a, b \in \mathbb{Z}$ ,  $b > 0$ ?

- Start with  $\gcd(a, b)$ . (We can assume Wlog that  $a > b$ )
- By Q-R theorem  $a = bq_1 + r_1$  for some  $q_1 \in \mathbb{Z}$ ,  $0 \leq r_1 < b$ .

# Euclidean algorithm to compute $\gcd(a, b)$

Reminder:

## Theorem (Q-R Theorem)

Let  $a, b \in \mathbb{Z}$ ,  $b > 0$ . Then there **exists** unique **integers  $q$  and  $r$**  such that  $a = qb + r$ ,  $0 \leq r < b$

## Proposition

Suppose  $a, b \in \mathbb{Z}$ ,  $b \neq 0$ . If  $a = bq + r$ , for some  $q, r \in \mathbb{Z}$  then  $\gcd(a, b) = \gcd(r, b)$ .

**Euclidean Algorithm:** How to compute  $\gcd(a, b)$ ,  $a, b \in \mathbb{Z}$ ,  $b > 0$ ?

- Start with  $\gcd(a, b)$ . (We can assume Wlog that  $a > b$ )
- By Q-R theorem  $a = bq_1 + r_1$  for some  $q_1 \in \mathbb{Z}$ ,  $0 \leq r_1 < b$ .
- By Proposition above  $\gcd(a, b) = \gcd(b, r_1)$ .

# Euclidean algorithm to compute $\gcd(a, b)$

Reminder:

## Theorem (Q-R Theorem)

Let  $a, b \in \mathbb{Z}$ ,  $b > 0$ . Then there **exists** unique **integers  $q$  and  $r$**  such that  $a = qb + r$ ,  $0 \leq r < b$

## Proposition

Suppose  $a, b \in \mathbb{Z}$ ,  $b \neq 0$ . If  $a = bq + r$ , for some  $q, r \in \mathbb{Z}$  then  $\gcd(a, b) = \gcd(r, b)$ .

**Euclidean Algorithm: How to compute  $\gcd(a, b)$ ,  $a, b \in \mathbb{Z}$ ,  $b > 0$ ?**

- Start with  $\gcd(a, b)$ . (We can assume Wlog that  $a > b$ )
- By Q-R theorem  $a = bq_1 + r_1$  for some  $q_1 \in \mathbb{Z}$ ,  $0 \leq r_1 < b$ .
- By Proposition above  $\gcd(a, b) = \gcd(b, r_1)$ .
- Again by By Q-R theorem  $b = r_1q_2 + r_2$  for some  $q_2 \in \mathbb{Z}$ ,  $0 \leq r_2 < r_1$ .

# Euclidean algorithm to compute $\gcd(a, b)$

Reminder:

## Theorem (Q-R Theorem)

Let  $a, b \in \mathbb{Z}$ ,  $b > 0$ . Then there **exists** unique **integers  $q$  and  $r$**  such that  $a = qb + r$ ,  $0 \leq r < b$

## Proposition

Suppose  $a, b \in \mathbb{Z}$ ,  $b \neq 0$ . If  $a = bq + r$ , for some  $q, r \in \mathbb{Z}$  then  $\gcd(a, b) = \gcd(r, b)$ .

**Euclidean Algorithm: How to compute  $\gcd(a, b)$ ,  $a, b \in \mathbb{Z}$ ,  $b > 0$ ?**

- Start with  $\gcd(a, b)$ . (We can assume Wlog that  $a > b$ )
- By Q-R theorem  $a = bq_1 + r_1$  for some  $q_1 \in \mathbb{Z}$ ,  $0 \leq r_1 < b$ .
- By Proposition above  $\gcd(a, b) = \gcd(b, r_1)$ .
- Again by Q-R theorem  $b = r_1q_2 + r_2$  for some  $q_2 \in \mathbb{Z}$ ,  $0 \leq r_2 < r_1$ .
- By the Proposition again  $\gcd(b, r_1) = \gcd(r_1, r_2)$ .



# Euclidean algorithm to compute $\gcd(a, b)$

Reminder:

## Theorem (Q-R Theorem)

Let  $a, b \in \mathbb{Z}$ ,  $b > 0$ . Then there **exists** unique **integers  $q$  and  $r$**  such that  $a = qb + r$ ,  $0 \leq r < b$

## Proposition

Suppose  $a, b \in \mathbb{Z}$ ,  $b \neq 0$ . If  $a = bq + r$ , for some  $q, r \in \mathbb{Z}$  then  $\gcd(a, b) = \gcd(r, b)$ .

**Euclidean Algorithm:** How to compute  $\gcd(a, b)$ ,  $a, b \in \mathbb{Z}$ ,  $b > 0$ ?

- Start with  $\gcd(a, b)$ . (We can assume Wlog that  $a > b$ )
- By Q-R theorem  $a = bq_1 + r_1$  for some  $q_1 \in \mathbb{Z}$ ,  $0 \leq r_1 < b$ .
- By Proposition above  $\gcd(a, b) = \gcd(b, r_1)$ .
- Again by Q-R theorem  $b = r_1q_2 + r_2$  for some  $q_2 \in \mathbb{Z}$ ,  $0 \leq r_2 < r_1$ .
- By the Proposition again  $\gcd(b, r_1) = \gcd(r_1, r_2)$ .
- Iterate!! Since  $b > r_1 > r_2 > \dots > 0$ , eventually  $r_k = 0$  for some  $k$  and

$$\gcd(a, b) = \gcd(r_{k-1}, r_k) = \gcd(r_{k-1}, 0) = r_{k-1}$$

# Euclidean algorithm: Example

Compute  $\gcd(42, 18)$

- Start with  $\gcd(42, 18)$ .
- By Q-R theorem  $a = bq_1 + r_1$  for some  $q_1 \in \mathbb{Z}$ ,  $0 \leq r_1 < b$ .
- By our Proposition  $\gcd(a, b) = \gcd(b, r_1)$ .
- Again by By Q-R theorem  $b = r_1q_2 + r_2$  for some  $q_2 \in \mathbb{Z}$ ,  $0 \leq r_2 < r_1$ .
- By our Proposition again  $\gcd(b, r_1) = \gcd(r_1, r_2)$ .

# Euclidean algorithm: Example

Compute  $\gcd(42, 18)$

- Start with  $\gcd(42, 18)$ .
- By Q-R theorem  $a = bq_1 + r_1$  for some  $q_1 \in \mathbb{Z}$ ,  $0 \leq r_1 < b$ .
- By our Proposition  $\gcd(a, b) = \gcd(b, r_1)$ .
- Again by By Q-R theorem  $b = r_1q_2 + r_2$  for some  $q_2 \in \mathbb{Z}$ ,  $0 \leq r_2 < r_1$ .
- By our Proposition again  $\gcd(b, r_1) = \gcd(r_1, r_2)$ .
- Iteration is finished!