

VIDEO XIV

REALS 1: FIELDS

The reals: motivation

- We have now addition, subtraction, multiplication and division.

The reals: motivation

- We have now addition, subtraction, multiplication and division.
- **But we cannot solve for $x \in \mathbb{Q}$ equations like**

$$x^2 = 2!$$


Wanted: " $x = \sqrt{2}$ "

The reals: motivation

- We have now addition, subtraction, multiplication and division.
- **But we cannot solve for $x \in \mathbb{Q}$ equations like**

$$x^2 = 2!$$

Wanted: " $x = \sqrt{2}$ "




Intuitively if we represent rationals as points on the line...there are gaps!

The reals: motivation

- We have now addition, subtraction, multiplication and division.
- **But we cannot solve for $x \in \mathbb{Q}$ equations like**

$$x^2 = 2!$$

Wanted: " $x = \sqrt{2}$ "



Intuitively if we represent rationals as points on the line...there are gaps!

Example: $\sqrt{2}$ is not rational.

idea: See reals as sequences of rational numbers, i.e. π is the limit of:

$(3, 3.1, 3.14, 3.141, 3.1415, 3.14159, 3.141592, \dots, 3.14159265358979, \dots)$.

Construction of the reals

Actually there are **several ways to construct the real numbers** from \mathbb{Q} :

- as limits of **Cauchy sequences of rational numbers**.

Construction of the reals

Actually there are **several ways to construct the real numbers** from \mathbb{Q} :

- as limits of **Cauchy sequences of rational numbers**.
- with **nested intervals**

Construction of the reals

Actually there are **several ways to construct the real numbers** from \mathbb{Q} :

- as limits of **Cauchy sequences of rational numbers**.
- with **nested intervals**
- with **Dedekind cuts**.

Construction of the reals

Actually there are **several ways to construct the real numbers** from \mathbb{Q} :

- as limits of **Cauchy sequences of rational numbers**.
- with **nested intervals**
- with **Dedekind cuts**.

This is tedious and beyond the scope of this module...instead we will formulate them as a collection of axioms!!

Fields axioms

We want the reals to be a set \mathbb{R} with an addition $+: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ and a multiplication $\cdot: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ such that $\forall x, y, z \in \mathbb{R}$

Fields axioms

We want the reals to be a set \mathbb{R} with an addition $+: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ and a multiplication $\cdot: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ such that $\forall x, y, z \in \mathbb{R}$

(A) Axioms for addition

- 1 (A1) $x + y = y + x$ (addition is commutative)
- 2 (A2) $(x + y) + z = x + (y + z)$ (addition is associative)
- 3 (A3) \mathbb{R} contains an element 0 such that $0 + x = x$ (neutral element of +)
- 4 (A4) $\exists -x \in \mathbb{R}$ such that $x + (-x) = 0$. (additive inverse)

Fields axioms

We want the reals to be a set \mathbb{R} with an **addition** $+: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ and a **multiplication** $\cdot: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ such that $\forall x, y, z \in \mathbb{R}$

(A) Axioms for addition

- ➊ (A1) $x + y = y + x$ (**addition is commutative**)
- ➋ (A2) $(x + y) + z = x + (y + z)$ (**addition is associative**)
- ➌ (A3) \mathbb{R} contains an element 0 such that $0 + x = x$ (**neutral element of +**)
- ➍ (A4) $\exists -x \in \mathbb{R}$ such that $x + (-x) = 0$. (**additive inverse**)

(M) Axioms for multiplication

- ➊ (M1) $xy = yx$ (**multiplication is commutative**)
- ➋ (M2) $(xy)z = x(yz)$ (**multiplication is associative**)
- ➌ (M3) \mathbb{R} contains an element $1 \neq 0$ such that $1x = x$. (**neutral element of \cdot**)
- ➍ (M4) For each $x \neq 0 \exists x^{-1}$, such that $xx^{-1} = 1$ (**multiplicative inverse**)

Fields axioms

We want the reals to be a set \mathbb{R} with an addition $+: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ and a multiplication $\cdot: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ such that $\forall x, y, z \in \mathbb{R}$

(A) Axioms for addition

- 1 (A1) $x + y = y + x$ (addition is commutative)
- 2 (A2) $(x + y) + z = x + (y + z)$ (addition is associative)
- 3 (A3) \mathbb{R} contains an element 0 such that $0 + x = x$ (neutral element of +)
- 4 (A4) $\exists -x \in \mathbb{R}$ such that $x + (-x) = 0$. (additive inverse)

(M) Axioms for multiplication

- 1 (M1) $xy = yx$ (multiplication is commutative)
- 2 (M2) $(xy)z = x(yz)$ (multiplication is associative)
- 3 (M3) \mathbb{R} contains an element $1 \neq 0$ such that $1x = x$. (neutral element of \cdot)
- 4 (M4) For each $x \neq 0 \exists x^{-1}$, such that $xx^{-1} = 1$ (multiplicative inverse)

(D) The distributive law: $x(y + z) = xy + xz$.

Fields axioms

We want the reals to be a set \mathbb{R} with an **addition** $+: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ and a **multiplication** $\cdot: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ such that $\forall x, y, z \in \mathbb{R}$

(A) Axioms for addition

- ① (A1) $x + y = y + x$ (**addition is commutative**)
- ② (A2) $(x + y) + z = x + (y + z)$ (**addition is associative**)
- ③ (A3) \mathbb{R} contains an element 0 such that $0 + x = x$ (**neutral element of $+$**)
- ④ (A4) $\exists -x \in \mathbb{R}$ such that $x + (-x) = 0$. (**additive inverse**)

(M) Axioms for multiplication

- ① (M1) $xy = yx$ (**multiplication is commutative**)
- ② (M2) $(xy)z = x(yz)$ (**multiplication is associative**)
- ③ (M3) \mathbb{R} contains an element $1 \neq 0$ such that $1x = x$. (**neutral element of \cdot**)
- ④ (M4) For each $x \neq 0 \exists x^{-1}$, such that $xx^{-1} = 1$ (**multiplicative inverse**)

(D) **The distributive law:** $x(y + z) = xy + xz$.

A set satisfying all these axioms is called a **field**

Question

Is it enough to define the reals uniquely?

Example

Let $\mathbb{F}_2 = \{0, 1\}$, with $+: \mathbb{F}_2 \times \mathbb{F}_2 \rightarrow \mathbb{F}_2$, $\cdot: \mathbb{F}_2 \times \mathbb{F}_2 \rightarrow \mathbb{F}_2$, such that

$$0 + 0 = 0, 0 + 1 = 1 + 0 = 1, 1 + 1 = 0, \quad 0 \cdot 0 = 0, 0 \cdot 1 = 1 \cdot 0 = 0, 1 \cdot 1 = 1.$$

Check the axioms!

- 1 (A1) $x + y = y + x$
- 2 (A2) $(x + y) + z = x + (y + z)$
- 3 (A3) \exists an element 0 in \mathbb{F}_2 such that $0 + x = x$
- 4 (A4) $\exists -x \in \mathbb{R}$ such that $x + (-x) = 0$.
- 5 (M1) $x \cdot y = y \cdot x$
- 6 (M2) $(x \cdot y) \cdot z = x \cdot (y \cdot z)$
- 7 (M3) \exists an element $1 \neq 0$ such that $1 \cdot x = x$.
- 8 (M4) For each $x \neq 0 \exists x^{-1}$, such that $x \cdot x^{-1} = 1$
- 9 (D) $x \cdot (y + z) = x \cdot y + x \cdot z$.

Computation rules from the axioms

Question

Are the previous axioms enough to define the reals uniquely?

Answer to our question: No! \mathbb{Q} , \mathbb{F}_2 satisfies the axioms of fields.

Computation rules from the axioms

Question

Are the previous axioms enough to define the reals uniquely?

Answer to our question: No! \mathbb{Q} , \mathbb{F}_2 satisfies the axioms of fields.
Nevertheless from these axioms we can prove usual rules for $+$ and \cdot .

Proposition

Let $x, y, z \in \mathbb{R}$. Then

- 1) If $x + z = y + z$, then $x = y$.*
- 2) If $x \in \mathbb{R}$, then $x \cdot 0 = 0$*

★: This rules can be derived for an arbitrary field!

Question

But so how do we get \mathbb{R} then.....?

VIDEO XV

REALS 2: ORDERED FIELDS

Ordered fields axioms

Reminder:

A **partial order relation** R on a set X is a binary relation such that

- $\forall x \in X, R(x, x)$ (Reflexivity)
- $\forall x, y \in X, R(x, y) \wedge R(y, x) \Rightarrow x = y$ (Anti-symmetry)
- $\forall x, y, z \in X, R(x, y) \wedge R(y, z) \Rightarrow R(x, z)$ (Transitivity)

R is called a **total order** if additionally $\forall x, y \in X, R(x, y) \vee R(y, x)$

Ordered fields axioms

Reminder:

A **partial order relation** R on a set X is a binary relation such that

- $\forall x \in X, R(x, x)$ (Reflexivity)
- $\forall x, y \in X, R(x, y) \wedge R(y, x) \Rightarrow x = y$ (Anti-symmetry)
- $\forall x, y, z \in X, R(x, y) \wedge R(y, z) \Rightarrow R(x, z)$ (Transitivity)

R is called a **total order** if additionally $\forall x, y \in X, R(x, y) \vee R(y, x)$

Ordered fields axioms:

A field \mathbb{F} is called **ordered**, if there is a total order \leq on \mathbb{F} such that additionally if $x, y, z \in \mathbb{F}$, $x \leq y$ then

- 1 (O1) $x + z \leq y + z$
- 2 (O2) if moreover $z \geq 0$ then $xz \leq yz$

Ordered fields axioms

Reminder:

A **partial order relation** R on a set X is a binary relation such that

- $\forall x \in X, R(x, x)$ (Reflexivity)
- $\forall x, y \in X, R(x, y) \wedge R(y, x) \Rightarrow x = y$ (Anti-symmetry)
- $\forall x, y, z \in X, R(x, y) \wedge R(y, z) \Rightarrow R(x, z)$ (Transitivity)

R is called a **total order** if additionally $\forall x, y \in X, R(x, y) \vee R(y, x)$

Ordered fields axioms:

A field \mathbb{F} is called **ordered**, if there is a total order \leq on \mathbb{F} such that additionally if $x, y, z \in \mathbb{F}$, $x \leq y$ then

- 1 (O1) $x + z \leq y + z$
- 2 (O2) if moreover $z \geq 0$ then $xz \leq yz$

Question

Is this additional property enough to define the reals uniquely now?

Ordered fields axioms

Reminder:

A **partial order relation** R on a set X is a binary relation such that

- $\forall x \in X, R(x, x)$ (Reflexivity)
- $\forall x, y \in X, R(x, y) \wedge R(y, x) \Rightarrow x = y$ (Anti-symmetry)
- $\forall x, y, z \in X, R(x, y) \wedge R(y, z) \Rightarrow R(x, z)$ (Transitivity)

R is called a **total order** if additionally $\forall x, y \in X, R(x, y) \vee R(y, x)$

Ordered fields axioms:

A field \mathbb{F} is called **ordered**, if there is a total order \leq on \mathbb{F} such that additionally if $x, y, z \in \mathbb{F}$, $x \leq y$ then

- 1 (O1) $x + z \leq y + z$
- 2 (O2) if moreover $z \geq 0$ then $xz \leq yz$

Question

Is this additional property enough to define the reals uniquely now?

Answer: Still not! \mathbb{Q} satisfies this axioms! Such fields are called ordered fields.

Nevertheless from there we can derive all usual rules for the inequality!

Proposition

For all $x, y, z \in \mathbb{F}$, \mathbb{F} an ordered field,

- ➊ *if $x < y$, then $x + z < y + z$,*
- ➋ *if $x < y$ and $z > 0$, then $xz < yz$*
- ➌ *$x < 0$ if and only if $-x > 0$ and $x > 0$ if and only if $-x < 0$.*
- ➍ *$0 < 1$*

VIDEO XVI

REALS 3: THE AXIOM OF COMPLETENESS

Lowest/upper bound–supremum/infimum

Definition

- A set S is called **bounded above** if there is a number B such that $x \leq B$ for all $x \in S$.
Any such B is called an **upper bound** for S .
- A set S is called **bounded below** if there is a number B such that $x \geq B$ for all $x \in S$.
Any such B is called an **lower bound** for S .
- We call s a **least upper bound or supremum** of S if
 - ① s is an **upper bound** for S ,
 - ② if $B < s$ then B is **not an upper bound** for S .

Notation: $\sup S$

- We call s a **largest lower bound or infimum** of S if
 - ① s is a **lower bound** for S ,
 - ② if $B > s$ then B is **not a lower bound** for S .

Notation: $\inf S$

Now we can finish our axiomatic definition of the reals!

Completeness of the reals

Completeness axiom:

- ① (C) Every non-empty set S of real numbers which is bounded above has a supremum.

Completeness of the reals

Completeness axiom:

- ① (C) Every non-empty set S of real numbers which is bounded above has a supremum.
- **The completeness axiom captures our intuition that the real line has no holes in it!**

Completeness of the reals

Completeness axiom:

- ① (C) Every non-empty set S of real numbers which is bounded above has a supremum.
- The completeness axiom captures our intuition that the real line has no holes in it!
- This is **not true any more for \mathbb{Q}** . Ex: $\{x \in \mathbb{Q} | x^2 \leq 2\}$.

Completeness of the reals

Completeness axiom:

- ① (C) Every non-empty set S of real numbers which is bounded above has a supremum.

- The completeness axiom captures our intuition that the real line has no holes in it!
- This is **not true any more for \mathbb{Q}** . Ex: $\{x \in \mathbb{Q} | x^2 \leq 2\}$.

Completeness of the reals

Completeness axiom:

- ① (C) Every non-empty set S of real numbers which is bounded above has a supremum.

- **The completeness axiom captures our intuition that the real line has no holes in it!**
- **This is not true any more for \mathbb{Q} . Ex: $\{x \in \mathbb{Q} | x^2 \leq 2\}$.**

Definition

The **reals** are the field denoted by \mathbb{R} satisfying **all the 12 axioms** we have seen so far:

- axioms (A1)–(A4), (M1)–(M4) and (D) for addition and multiplication,
- axioms (O1) and (O2) for the total order,
- axiom of completeness (C).

It is called a **complete ordered field**. \mathbb{R} is the only such field.

Corollary (Archimedean property /Axiom of Eudoxos)

$$\forall x, y \in \mathbb{R}, x > 0, \exists n \in \mathbb{N}, \text{ such that } nx > y.$$

Proof: Exercise!

Lemma

Let $r \in \mathbb{R}$, $r > 0$. Then there exists a unique $n \in \mathbb{N}$, such that $n - 1 \leq r < n$

proof: Consider the set $S = \{m \in \mathbb{N} | m > r\}$

- By the Archimedean property $S \subseteq \mathbb{N}$ is not empty.
- Therefore by the Well-Ordering Principle there exists a (unique) least element $n \in S$ (i.e. $m > n, \forall m \in S$).

Corollary (Archimedean property /Axiom of Eudoxos)

$$\forall x, y \in \mathbb{R}, x > 0, \exists n \in \mathbb{N}, \text{ such that } nx > y.$$

Proof: Exercise!

Lemma

Let $r \in \mathbb{R}, r > 0$. Then there exists a unique $n \in \mathbb{N}$, such that $n - 1 \leq r < n$

proof: Consider the set $S = \{m \in \mathbb{N} | m > r\}$

- By the Archimedean property $S \subseteq \mathbb{N}$ is not empty.
- Therefore by the Well-Ordering Principle there exists a (unique) least element $n \in S$ (i.e. $m > n, \forall m \in S$).
- But $n > r$ since $n \in S$

Corollary (Archimedean property /Axiom of Eudoxos)

$$\forall x, y \in \mathbb{R}, x > 0, \exists n \in \mathbb{N}, \text{ such that } nx > y.$$

Proof: Exercise!

Lemma

Let $r \in \mathbb{R}, r > 0$. Then there exists a unique $n \in \mathbb{N}$, such that $n - 1 \leq r < n$

proof: Consider the set $S = \{m \in \mathbb{N} | m > r\}$

- By the Archimedean property $S \subseteq \mathbb{N}$ is not empty.
- Therefore by the Well-Ordering Principle there exists a (unique) least element $n \in S$ (i.e. $m > n, \forall m \in S$).
- But $n > r$ since $n \in S$
- Case 1: $n=1$, then $0 < r < n=1$.
Case 2: $n > 1$, then $n - 1 \in \mathbb{N}$, but not in S ! Hence $n - 1 \leq r < n$.

Applications

Corollary (Archimedean property /Axiom of Eudoxos)

$$\forall x, y \in \mathbb{R}, x > 0, \exists n \in \mathbb{N}, \text{ such that } nx > y.$$

Lemma

Let $x \in \mathbb{R}, x > 0$. There exists a unique $n \in \mathbb{N}$, such that $n - 1 \leq x < n$.

Proposition (Density of the rationals)

For all $x, y \in \mathbb{R}$, there exists $q \in \mathbb{Q}$, such that $x < q < y$.

Proof: Assume WLoG that $0 < x, y$, hence $y - x > 0$.

- Use Archimedean property:
- Use Above Lemma:
- Find a contradiction