

Predicates on relations.

We fix some background set X . We've seen examples of binary relations on X .

Now let's study the general theory of binary relations (binary relations will be variables).

Predicates on relations.

We fix some background set X . We've seen examples of binary relations on X .

Now let's study the general theory of binary relations (binary relations will be variables).

Remember the two notations for binary relations: sometimes we write $R(a, b)$ and sometimes we write $a \star b$.

Predicates on relations.

We fix some background set X . We've seen examples of binary relations on X .

Now let's study the general theory of binary relations (binary relations will be variables).

Remember the two notations for binary relations: sometimes we write $R(a, b)$ and sometimes we write $a \star b$.

In this lecture, we will look at four properties which a general binary relation may or may not have:

- Reflexivity;
- Symmetry;
- Antisymmetry;
- Transitivity.

Reflexivity

Let X be a set, and let \star be a binary relation on X . We will write $a \star b$ to be the true-false statement associated to a and b .

Reflexivity

Let X be a set, and let \star be a binary relation on X . We will write $a \star b$ to be the true-false statement associated to a and b .

New definition: A binary relation \star on a set X is *reflexive* if every element of X is related to itself.

Reflexivity

Let X be a set, and let \star be a binary relation on X . We will write $a \star b$ to be the true-false statement associated to a and b .

New definition: A binary relation \star on a set X is *reflexive* if every element of X is related to itself. In an exam, write this:

$$\forall a \in X, a \star a.$$

Reflexivity

Let X be a set, and let \star be a binary relation on X . We will write $a \star b$ to be the true-false statement associated to a and b .

New definition: A binary relation \star on a set X is *reflexive* if every element of X is related to itself. In an exam, write this:

$$\forall a \in X, a \star a.$$

For example, say $X = \mathbb{R}$ and \star is just $=$. If x is any real number at all, then $x = x$.

Reflexivity

Let X be a set, and let \star be a binary relation on X . We will write $a \star b$ to be the true-false statement associated to a and b .

New definition: A binary relation \star on a set X is *reflexive* if every element of X is related to itself. In an exam, write this:

$$\forall a \in X, a \star a.$$

For example, say $X = \mathbb{R}$ and \star is just $=$. If x is any real number at all, then $x = x$. So the proposition $\forall x \in \mathbb{R}, x = x$ is true.

Reflexivity

Let X be a set, and let \star be a binary relation on X . We will write $a \star b$ to be the true-false statement associated to a and b .

New definition: A binary relation \star on a set X is *reflexive* if every element of X is related to itself. In an exam, write this:

$$\forall a \in X, a \star a.$$

For example, say $X = \mathbb{R}$ and \star is just $=$. If x is any real number at all, then $x = x$. So the proposition $\forall x \in \mathbb{R}, x = x$ is true. So, by definition, $=$ is a reflexive binary relation on the reals.

Reflexivity

Let X be a set, and let \star be a binary relation on X . We will write $a \star b$ to be the true-false statement associated to a and b .

New definition: A binary relation \star on a set X is *reflexive* if every element of X is related to itself. In an exam, write this:

$$\forall a \in X, a \star a.$$

For example, say $X = \mathbb{R}$ and \star is just $=$. If x is any real number at all, then $x = x$. So the proposition $\forall x \in \mathbb{R}, x = x$ is true. So, by definition, $=$ is a reflexive binary relation on the reals.

This was first observed by Euclid! Check out [Euclid's common notions](#). Euclid's "common notion 4": "Things which coincide with one another equal one another."

Reflexivity

Let X be a set, and let \star be a binary relation on X . We will write $a \star b$ to be the true-false statement associated to a and b .

New definition: A binary relation \star on a set X is *reflexive* if every element of X is related to itself. In an exam, write this:

$$\forall a \in X, a \star a.$$

For example, say $X = \mathbb{R}$ and \star is just $=$. If x is any real number at all, then $x = x$. So the proposition $\forall x \in \mathbb{R}, x = x$ is true. So, by definition, $=$ is a reflexive binary relation on the reals.

This was first observed by Euclid! Check out [Euclid's common notions](#). Euclid's "common notion 4": "Things which coincide with one another equal one another."

Now here's some for you: is \leq a reflexive binary relation on the reals? How about $<$? Pause the video and have a go.

$x \leq x$ is true for every real number x , so \leq is reflexive.

$x \leq x$ is true for every real number x , so \leq is reflexive.

But $37 < 37$ is false, so $<$ is not reflexive.

$x \leq x$ is true for every real number x , so \leq is reflexive.

But $37 < 37$ is false, so $<$ is not reflexive.

Let's say that I asked that question in an exam, and someone answered " $x < x$ is false, so $<$ is not reflexive.". How many marks should I give them, out of two?

$x \leq x$ is true for every real number x , so \leq is reflexive.

But $37 < 37$ is false, so $<$ is not reflexive.

Let's say that I asked that question in an exam, and someone answered " $x < x$ is false, so $<$ is not reflexive.". How many marks should I give them, out of two?

They proved $\forall x \in X, \neg(x < x)$. They were supposed to prove $\exists x \in X, \neg(x < x)$.

New definition: The binary relation \star on X is *symmetric* if for all $a, b \in X$ we have $a \star b \implies b \star a$.

The thing to write in a test: $\forall a, b \in X, a \star b \implies b \star a$.

Symmetry

New definition: The binary relation \star on X is *symmetric* if for all $a, b \in X$ we have $a \star b \implies b \star a$.

The thing to write in a test: $\forall a, b \in X, a \star b \implies b \star a$.

For example, $=$ is a symmetric binary relation on all X , because if $a, b \in X$ and $a = b$, then $b = a$.

Reminder: \star on X is symmetric if (and only if):

$$\forall a, b \in X, a \star b \implies b \star a.$$

Symmetry

Reminder: \star on X is symmetric if (and only if):

$$\forall a, b \in X, a \star b \implies b \star a.$$

Example. Is \leq on \mathbb{R} symmetric? What do you think?

Reminder: \star on X is symmetric if (and only if):

$$\forall a, b \in X, a \star b \implies b \star a.$$

Example. Is \leq on \mathbb{R} symmetric? What do you think?

The opposite of $\forall a, b \in \mathbb{R}, a \leq b \implies b \leq a$ is...

Reminder: \star on X is symmetric if (and only if):

$$\forall a, b \in X, a \star b \implies b \star a.$$

Example. Is \leq on \mathbb{R} symmetric? What do you think?

The opposite of $\forall a, b \in \mathbb{R}, a \leq b \implies b \leq a$ is...

$$\exists a, b \in \mathbb{R}, \neg(a \leq b \implies b \leq a)$$

Symmetry

Reminder: \star on X is symmetric if (and only if):

$$\forall a, b \in X, a \star b \implies b \star a.$$

Example. Is \leq on \mathbb{R} symmetric? What do you think?

The opposite of $\forall a, b \in \mathbb{R}, a \leq b \implies b \leq a$ is...

$$\exists a, b \in \mathbb{R}, \neg(a \leq b \implies b \leq a)$$

So how about $a = 1$ and $b = 37$? Then $1 \leq 37$ is true, $37 \leq 1$ is false, so $1 \leq 37 \implies 37 \leq 1$ is also false. So \neg it is true!

Symmetry

Reminder: \star on X is symmetric if (and only if):

$$\forall a, b \in X, a \star b \implies b \star a.$$

Example. Is \leq on \mathbb{R} symmetric? What do you think?

The opposite of $\forall a, b \in \mathbb{R}, a \leq b \implies b \leq a$ is...

$$\exists a, b \in \mathbb{R}, \neg(a \leq b \implies b \leq a)$$

So how about $a = 1$ and $b = 37$? Then $1 \leq 37$ is true, $37 \leq 1$ is false, so $1 \leq 37 \implies 37 \leq 1$ is also false. So \neg it is true!

So \leq is not symmetric.

What about the relation $P(a, b)$ on \mathbb{N} where $P(a, b)$ is true if and only if a and b were both even or both odd?

What about the relation $P(a, b)$ on \mathbb{N} where $P(a, b)$ is true if and only if a and b were both even or both odd?

If a has the same remainder mod 2 as b , then certainly b has the same remainder as a !

What about the relation $P(a, b)$ on \mathbb{N} where $P(a, b)$ is true if and only if a and b were both even or both odd?

If a has the same remainder mod 2 as b , then certainly b has the same remainder as a !

So P is symmetric.

Antisymmetry

Let's start the “antisymmetry” section with a warning:

Be careful – “antisymmetric” does not mean “not symmetric”!

Antisymmetry

Let's start the “antisymmetry” section with a warning:

Be careful – “antisymmetric” does not mean “not symmetric”!

A binary relationship \star on X is *antisymmetric* if for every a and b , if $a \star b$ and $b \star a$ then $a = b$.

Antisymmetry

Let's start the “antisymmetry” section with a warning:

Be careful – “antisymmetric” does not mean “not symmetric”!

A binary relationship \star on X is *antisymmetric* if for every a and b , if $a \star b$ and $b \star a$ then $a = b$.

The standard example is \leq on \mathbb{R} . This relation is antisymmetric because if a and b are arbitrary real numbers and $a \leq b$ and $b \leq a$, then $a = b$.

Antisymmetry

Let's start the “antisymmetry” section with a warning:

Be careful – “antisymmetric” does not mean “not symmetric”!

A binary relationship \star on X is *antisymmetric* if for every a and b , if $a \star b$ and $b \star a$ then $a = b$.

The standard example is \leq on \mathbb{R} . This relation is antisymmetric because if a and b are arbitrary real numbers and $a \leq b$ and $b \leq a$, then $a = b$.

Recall that $=$ on X was symmetric. Is it also antisymmetric? Pause the video and figure out what this question *means*.

Antisymmetry

Let's start the “antisymmetry” section with a warning:

Be careful – “antisymmetric” does not mean “not symmetric”!

A binary relationship \star on X is *antisymmetric* if for every a and b , if $a \star b$ and $b \star a$ then $a = b$.

The standard example is \leq on \mathbb{R} . This relation is antisymmetric because if a and b are arbitrary real numbers and $a \leq b$ and $b \leq a$, then $a = b$.

Recall that $=$ on X was symmetric. Is it also antisymmetric? Pause the video and figure out what this question *means*.

It means “Say a and b are elements of X , and $a = b$ and $b = a$. Does $a = b$?”

This is definitely true! So equality is symmetric and antisymmetric.

Can you write down a binary relation on $\{1, 2, 3, 4\}$ which is not symmetric and not antisymmetric either?

Can you write down a binary relation on $\{1, 2, 3, 4\}$ which is not symmetric and not antisymmetric either?

If $Q(a, b)$ is the relation, then what has to be true about Q to make it not symmetric and not antisymmetric?

Can you write down a binary relation on $\{1, 2, 3, 4\}$ which is not symmetric and not antisymmetric either?

If $Q(a, b)$ is the relation, then what has to be true about Q to make it not symmetric and not antisymmetric?

Symmetry says $\forall a, b, Q(a, b) \implies Q(b, a)$. How can we make that fail?

Can you write down a binary relation on $\{1, 2, 3, 4\}$ which is not symmetric and not antisymmetric either?

If $Q(a, b)$ is the relation, then what has to be true about Q to make it not symmetric and not antisymmetric?

Symmetry says $\forall a, b, Q(a, b) \implies Q(b, a)$. How can we make that fail?

We could say $Q(1, 2)$ is true and $Q(2, 1)$ is false. That will make symmetry fail.

Can you write down a binary relation on $\{1, 2, 3, 4\}$ which is not symmetric and not antisymmetric either?

If $Q(a, b)$ is the relation, then what has to be true about Q to make it not symmetric and not antisymmetric?

Symmetry says $\forall a, b, Q(a, b) \implies Q(b, a)$. How can we make that fail?

We could say $Q(1, 2)$ is true and $Q(2, 1)$ is false. That will make symmetry fail.

Now how shall we define $Q(3, 4)$ and $Q(4, 3)$ to make antisymmetry fail?

Can you write down a binary relation on $\{1, 2, 3, 4\}$ which is not symmetric and not antisymmetric either?

If $Q(a, b)$ is the relation, then what has to be true about Q to make it not symmetric and not antisymmetric?

Symmetry says $\forall a, b, Q(a, b) \implies Q(b, a)$. How can we make that fail?

We could say $Q(1, 2)$ is true and $Q(2, 1)$ is false. That will make symmetry fail.

Now how shall we define $Q(3, 4)$ and $Q(4, 3)$ to make antisymmetry fail?

Let's make $Q(3, 4)$ and $Q(4, 3)$ both true. Then Q is not antisymmetric.

Can you write down a binary relation on $\{1, 2, 3, 4\}$ which is not symmetric and not antisymmetric either?

If $Q(a, b)$ is the relation, then what has to be true about Q to make it not symmetric and not antisymmetric?

Symmetry says $\forall a, b, Q(a, b) \implies Q(b, a)$. How can we make that fail?

We could say $Q(1, 2)$ is true and $Q(2, 1)$ is false. That will make symmetry fail.

Now how shall we define $Q(3, 4)$ and $Q(4, 3)$ to make antisymmetry fail?

Let's make $Q(3, 4)$ and $Q(4, 3)$ both true. Then Q is not antisymmetric.

Now let $Q(1, 3)$ and all the others just be fixed (e.g. random, or all false) and we have a relation which is not symmetric or antisymmetric.

New definition: The binary relation \star on X is *transitive* if for all $a, b, c \in X$ we have $(a \star b \wedge b \star c) \implies a \star c$.

The thing to write in a test:

$$\forall a, b, c \in X, (a \star b \wedge b \star c) \implies a \star c$$

.

Transitivity

New definition: The binary relation \star on X is *transitive* if for all $a, b, c \in X$ we have $(a \star b \wedge b \star c) \implies a \star c$.

The thing to write in a test:

$$\forall a, b, c \in X, (a \star b \wedge b \star c) \implies a \star c$$

.

Famous examples of transitivity: \leq and $<$. For example – if $a \leq b$ and $b \leq c$ then $a \leq c$.

Transitivity

New definition: The binary relation \star on X is *transitive* if for all $a, b, c \in X$ we have $(a \star b \wedge b \star c) \implies a \star c$.

The thing to write in a test:

$$\forall a, b, c \in X, (a \star b \wedge b \star c) \implies a \star c$$

.

Famous examples of transitivity: \leq and $<$. For example – if $a \leq b$ and $b \leq c$ then $a \leq c$.

If you've played the natural number game, you'll know how to prove that \leq is transitive on the natural numbers. What the proof of transitivity of \leq looks like on the real numbers depends on what you think the real numbers are.

Let X be any set, and consider the binary relation of equality. Is it transitive?

Let X be any set, and consider the binary relation of equality. Is it transitive?

What does that *mean*?

Let X be any set, and consider the binary relation of equality. Is it transitive?

What does that *mean*?

$$\forall a, b, c \in X, (a = b \wedge b = c) \implies a = c$$

.
This is Euclid's first [common notion](#)!

How many transitive relations are there on a set of size 25?

How many transitive relations are there on a set of size 25?

Nobody knows!

Only known algorithms to compute this take far too long to run on a normal computer. See [this link](#) for what is known.

Transitivity of a random relation can be hard to check.

So why don't we give it a go! Recall that transitivity is:

$$\forall a, b, c \in X, (a \star b \wedge b \star c) \implies a \star c.$$

Remember the random relation R on $\{A, B, C\}$ defined by $R(A, B)$ and $R(A, C)$ are true, and everything else is false? Is that relation transitive?

Lean can prove that by checking all the cases!

So why don't we give it a go! Recall that transitivity is:

$$\forall a, b, c \in X, (a \star b \wedge b \star c) \implies a \star c.$$

Remember the random relation R on $\{A, B, C\}$ defined by $R(A, B)$ and $R(A, C)$ are true, and everything else is false? Is that relation transitive?

Lean can prove that by checking all the cases!

Pause the video and have a go yourself.

Transitivity: $\forall x, y, z \in X, (x \star y \wedge y \star z) \implies x \star z$.

Consider the random relation R on $\{A, B, C\}$ defined by $R(A, B)$ and $R(A, C)$ are true, and everything else is false: is R transitive?

Here's a proof.

Transitivity: $\forall x, y, z \in X, (x \star y \wedge y \star z) \implies x \star z$.

Consider the random relation R on $\{A, B, C\}$ defined by $R(A, B)$ and $R(A, C)$ are true, and everything else is false: is R transitive?

Here's a proof.

Before we start, let's observe that if $R(x, y)$ is true, then x must be A (look at all the cases). Let's call this hypothesis $(*)$.

Transitivity: $\forall x, y, z \in X, (x \star y \wedge y \star z) \implies x \star z$.

Consider the random relation R on $\{A, B, C\}$ defined by $R(A, B)$ and $R(A, C)$ are true, and everything else is false: is R transitive?

Here's a proof.

Before we start, let's observe that if $R(x, y)$ is true, then x must be A (look at all the cases). Let's call this hypothesis $(*)$.

Now let's assume $R(x, y)$ and $R(y, z)$ are true, and you want to prove $R(x, z)$,

Transitivity: $\forall x, y, z \in X, (x \star y \wedge y \star z) \implies x \star z$.

Consider the random relation R on $\{A, B, C\}$ defined by $R(A, B)$ and $R(A, C)$ are true, and everything else is false: is R transitive?

Here's a proof.

Before we start, let's observe that if $R(x, y)$ is true, then x must be A (look at all the cases). Let's call this hypothesis $(*)$.

Now let's assume $R(x, y)$ and $R(y, z)$ are true, and you want to prove $R(x, z)$, Applying $(*)$ we deduce $x = A$ and $y = A$, so you know $R(A, A)$ and $R(A, z)$, and you want to prove $R(A, z)$, so you're done!

Transitivity: $\forall x, y, z \in X, (x \star y \wedge y \star z) \implies x \star z$.

Consider the random relation R on $\{A, B, C\}$ defined by $R(A, B)$ and $R(A, C)$ are true, and everything else is false: is R transitive?

Here's a proof.

Before we start, let's observe that if $R(x, y)$ is true, then x must be A (look at all the cases). Let's call this hypothesis $(*)$.

Now let's assume $R(x, y)$ and $R(y, z)$ are true, and you want to prove $R(x, z)$, Applying $(*)$ we deduce $x = A$ and $y = A$, so you know $R(A, A)$ and $R(A, z)$, and you want to prove $R(A, z)$, so you're done!

Is there something wrong with that proof? Pause the video and have a think.

I'll show you how to prove the R transitivity question in the Lean video.

In all of the rest of the videos, we will be focussing on relations which are *reflexive* and *transitive*. But symmetry and antisymmetry pull in different directions. We should only assume one. The red pill or the blue pill.