

RATIONALS: VIDEO XIII

MOTIVATION AND FIRST DEFINITIONS

The rationals: motivation

- Until now we only have addition, subtraction, multiplication and a notion of divisibility.

The rationals: motivation

- Until now we only have addition, subtraction, multiplication and a notion of divisibility.
- **But we cannot solve for x equations like**

$$4x = 2!$$

We want to have " $x = \frac{2}{4}$ "!

The rationals: motivation

- Until now we only have addition, subtraction, multiplication and a notion of divisibility.
- **But we cannot solve for x equations like**

$$4x = 2!$$

We want to have " $x = \frac{2}{4}$ "!

Idea: define $\frac{2}{4}$ as the ordered pair $(2, 4) \in \mathbb{Z} \times \mathbb{Z}$?

The rationals: motivation

- Until now we only have addition, subtraction, multiplication and a notion of divisibility.
- **But we cannot solve for x equations like**

$$4x = 2!$$

We want to have " $x = \frac{2}{4}$ "!

Idea: define $\frac{2}{4}$ as the ordered pair $(2, 4) \in \mathbb{Z} \times \mathbb{Z}$?

Problem: $\frac{2}{4} = \frac{1}{2} = \frac{-25}{-50} \dots \Rightarrow (2, 4) = (1, 2) = (-25, -50)$ so such a pair is not unique.

The rationals: motivation

- Until now we only have addition, subtraction, multiplication and a notion of divisibility.
- **But we cannot solve for x equations like**

$$4x = 2!$$

We want to have " $x = \frac{2}{4}$ "!

Idea: define $\frac{2}{4}$ as the ordered pair $(2, 4) \in \mathbb{Z} \times \mathbb{Z}$?

Problem: $\frac{2}{4} = \frac{1}{2} = \frac{-25}{-50} \dots \Rightarrow (2, 4) = (1, 2) = (-25, -50)$ so such a pair is not unique.

★ Note: $\frac{m}{n} = \frac{m'}{n'}$ if and only if $mn' = m'n$ ★

What would be your next idea....?

Definition of the rationals

Our old Idea:

Define the equivalence relation on $\mathbb{Z} \times (\mathbb{Z} - \{0\})$

$$(m, n) \sim (m', n') \text{ if and only if } mn' = m'n.$$

Definition of the rationals

Our old Idea:

Define the equivalence relation on $\mathbb{Z} \times (\mathbb{Z} - \{0\})$

$$(m, n) \sim (m', n') \text{ if and only if } mn' = m'n.$$

$$\frac{m}{n} := cl((m, n)) = \{(x, y) \in \mathbb{Z} \times (\mathbb{Z} - \{0\}) \mid (x, y) \sim (m, n)\}$$

Definition of the rationals

Our old Idea:

Define the equivalence relation on $\mathbb{Z} \times (\mathbb{Z} - \{0\})$

$$(m, n) \sim (m', n') \text{ if and only if } mn' = m'n.$$

$$\frac{m}{n} := cl((m, n)) = \{(x, y) \in \mathbb{Z} \times (\mathbb{Z} - \{0\}) \mid (x, y) \sim (m, n)\}$$

\mathbb{Q} is the set of all equivalence classes $\frac{m}{n}$.

Definition of the rationals

Our old Idea:

Define the equivalence relation on $\mathbb{Z} \times (\mathbb{Z} - \{0\})$

$$(m, n) \sim (m', n') \text{ if and only if } mn' = m'n.$$

$$\frac{m}{n} := cl((m, n)) = \{(x, y) \in \mathbb{Z} \times (\mathbb{Z} - \{0\}) \mid (x, y) \sim (m, n)\}$$

\mathbb{Q} is the set of all equivalence classes $\frac{m}{n}$.

Definition of the rationals

Our old Idea:

Define the equivalence relation on $\mathbb{Z} \times (\mathbb{Z} - \{0\})$

$$(m, n) \sim (m', n') \text{ if and only if } mn' = m'n.$$

$$\frac{m}{n} := cl((m, n)) = \{(x, y) \in \mathbb{Z} \times (\mathbb{Z} - \{0\}) \mid (x, y) \sim (m, n)\}$$

\mathbb{Q} is the set of all equivalence classes $\frac{m}{n}$.

★ **Wanted:**

$$\frac{m_1}{n_1} + \frac{m_2}{n_2} = \frac{m_1 n_2 + n_1 m_2}{n_1 n_2}, \quad \frac{m_1}{n_1} \cdot \frac{m_2}{n_2} = \frac{m_1 m_2}{n_1 n_2}$$

We define on \mathbb{Q} :

Addition: $cl(m_1, n_1) +_{\mathbb{Q}} cl(m_2, n_2) := cl(m_1 n_2 + n_1 m_2, n_1 n_2)$

Multiplication: $cl(m_1, n_1) \cdot_{\mathbb{Q}} cl(m_2, n_2) := cl(m_1 m_2, n_1 n_2).$

Are you happy with this?

Are you happy with this?

Well if yes **you shouldn't...because we haven't checked...**

Are you happy with this?

Well if yes **you shouldn't...because we haven't checked...**

Proposition

Let $(m_i, n_i), (m'_i, n'_i) \in \mathbb{Z} \times \mathbb{Z}$, $i = 1, 2$. If $(m_1, n_1) \sim (m'_1, n'_1)$ and $(m_2, n_2) \sim (m'_2, n'_2)$, then

$$\begin{aligned} cl(m_1, n_1) +_{\mathbb{Q}} cl(m_2, n_2) &= cl(m'_1, n'_1) +_{\mathbb{Q}} cl(m'_2, n'_2), \\ cl(m_1, n_1) \cdot_{\mathbb{Q}} cl(m_2, n_2) &= cl(m'_1, n'_1) \cdot_{\mathbb{Q}} cl(m'_2, n'_2) \end{aligned}$$

Are you happy with this?

Well if yes **you shouldn't...because we haven't checked...**

Proposition

Let $(m_i, n_i), (m'_i, n'_i) \in \mathbb{Z} \times \mathbb{Z}$, $i = 1, 2$. If $(m_1, n_1) \sim (m'_1, n'_1)$ and $(m_2, n_2) \sim (m'_2, n'_2)$, then

$$\begin{aligned} cl(m_1, n_1) +_{\mathbb{Q}} cl(m_2, n_2) &= cl(m'_1, n'_1) +_{\mathbb{Q}} cl(m'_2, n'_2), \\ cl(m_1, n_1) \cdot_{\mathbb{Q}} cl(m_2, n_2) &= cl(m'_1, n'_1) \cdot_{\mathbb{Q}} cl(m'_2, n'_2) \end{aligned}$$

Now you can be happy :)...

Are you happy with this?

Well if yes **you shouldn't...because we haven't checked...**

Proposition

Let $(m_i, n_i), (m'_i, n'_i) \in \mathbb{Z} \times \mathbb{Z}$, $i = 1, 2$. If $(m_1, n_1) \sim (m'_1, n'_1)$ and $(m_2, n_2) \sim (m'_2, n'_2)$, then

$$\begin{aligned} cl(m_1, n_1) +_{\mathbb{Q}} cl(m_2, n_2) &= cl(m'_1, n'_1) +_{\mathbb{Q}} cl(m'_2, n'_2), \\ cl(m_1, n_1) \cdot_{\mathbb{Q}} cl(m_2, n_2) &= cl(m'_1, n'_1) \cdot_{\mathbb{Q}} cl(m'_2, n'_2) \end{aligned}$$

Now you can be happy :)...

...almost...

Are you happy with this?

Well if yes **you shouldn't...because we haven't checked...**

Proposition

Let $(m_i, n_i), (m'_i, n'_i) \in \mathbb{Z} \times \mathbb{Z}$, $i = 1, 2$. If $(m_1, n_1) \sim (m'_1, n'_1)$ and $(m_2, n_2) \sim (m'_2, n'_2)$, then

$$\begin{aligned}cl(m_1, n_1) +_{\mathbb{Q}} cl(m_2, n_2) &= cl(m'_1, n'_1) +_{\mathbb{Q}} cl(m'_2, n'_2), \\cl(m_1, n_1) \cdot_{\mathbb{Q}} cl(m_2, n_2) &= cl(m'_1, n'_1) \cdot_{\mathbb{Q}} cl(m'_2, n'_2)\end{aligned}$$

Now you can be happy :)...

...almost...

You still need to check that $\mathbb{Z} \subseteq \mathbb{Q}$.

Are you happy with this?

Well if yes **you shouldn't...because we haven't checked...**

Proposition

Let $(m_i, n_i), (m'_i, n'_i) \in \mathbb{Z} \times \mathbb{Z}$, $i = 1, 2$. If $(m_1, n_1) \sim (m'_1, n'_1)$ and $(m_2, n_2) \sim (m'_2, n'_2)$, then

$$\begin{aligned}cl(m_1, n_1) +_{\mathbb{Q}} cl(m_2, n_2) &= cl(m'_1, n'_1) +_{\mathbb{Q}} cl(m'_2, n'_2), \\cl(m_1, n_1) \cdot_{\mathbb{Q}} cl(m_2, n_2) &= cl(m'_1, n'_1) \cdot_{\mathbb{Q}} cl(m'_2, n'_2)\end{aligned}$$

Now you can be happy :)....

...almost...

You still need to check that $\mathbb{Z} \subseteq \mathbb{Q}$. This is done with the proposition

Proposition

The map

$$i : \mathbb{Z} \rightarrow \mathbb{Q}, n \mapsto i(n) = \frac{n}{1}.$$

is injective and preserves addition and multiplication.

Ordering on \mathbb{Z} and \mathbb{Q}

For the integers, we say $a - b <_{\mathbb{Z}} c - d$ if $a + d < b + c$. This does not depend on the choice of representatives, and extends the ordering on \mathbb{N} . We can now define an ordering on \mathbb{Q} .

Definition

Let $\frac{a}{b}, \frac{c}{d} \in \mathbb{Q}$. Assuming $b, d > 0$, we say that $\frac{a}{b} <_{\mathbb{Q}} \frac{c}{d}$ if $ad < bc$.

Ordering on \mathbb{Z} and \mathbb{Q}

For the integers, we say $a - b <_{\mathbb{Z}} c - d$ if $a + d < b + c$. This does not depend on the choice of representatives, and extends the ordering on \mathbb{N} . We can now define an ordering on \mathbb{Q} .

Definition

Let $\frac{a}{b}, \frac{c}{d} \in \mathbb{Q}$. Assuming $b, d > 0$, we say that $\frac{a}{b} <_{\mathbb{Q}} \frac{c}{d}$ if $ad < bc$.

★ We use the symbol $<_{\mathbb{Q}}$ to denote our ordering relation in \mathbb{Q} so as to distinguish it from the orderings $<_{\mathbb{Z}}$ in \mathbb{Z} or $<$ in \mathbb{N} .

Ordering on \mathbb{Z} and \mathbb{Q}

For the integers, we say $a - b <_{\mathbb{Z}} c - d$ if $a + d < b + c$. This does not depend on the choice of representatives, and extends the ordering on \mathbb{N} . We can now define an ordering on \mathbb{Q} .

Definition

Let $\frac{a}{b}, \frac{c}{d} \in \mathbb{Q}$. Assuming $b, d > 0$, we say that $\frac{a}{b} <_{\mathbb{Q}} \frac{c}{d}$ if $ad < bc$.

★ We use the symbol $<_{\mathbb{Q}}$ to denote our ordering relation in \mathbb{Q} so as to distinguish it from the orderings $<_{\mathbb{Z}}$ in \mathbb{Z} or $<$ in \mathbb{N} .

Proposition

$<_{\mathbb{Q}}$ defines a total order on \mathbb{Q} .