

RELATING THE CLASSICAL $\mu = 0$ CONJECTURE WITH COATES-SUJATHA CONJECTURE A

DEBANJANA KUNDU

ABSTRACT. The main goal of this article is to provide more evidence on the relationship between the Classical Iwasawa $\mu = 0$ Conjecture and the $\mu = 0$ Conjecture for fine Selmer groups (Conjecture A). We give sufficient conditions to prove the Classical $\mu = 0$ Conjecture that improves upon previously known results. Furthermore, we prove isogeny invariance of Conjecture A in some previously unknown cases. We also provide a class of examples for which Conjecture A holds independent of the Classical $\mu = 0$ Conjecture.

1. INTRODUCTION

Classical Iwasawa theory is concerned with the growth of class groups in towers of number fields. In [7], Iwasawa showed that in a \mathbb{Z}_p -extension of a number field F , the growth of the p -part of the class group is regular. In particular,

Theorem. *There exist constant non-negative integers λ and μ and a constant integer ν such that for large enough n ,*

$$|A_n| = p^{\mu p^n + \lambda n + \nu}$$

where A_n is the class group of F_n , the n -th layer in the tower.

Further, the following conjecture was made.

Classical Conjecture. *For the cyclotomic \mathbb{Z}_p -extension F_{cyc}/F , $\mu = 0$.*

The conjecture is known to hold for Abelian extensions F/\mathbb{Q} (see [4], [15]).

In [11], Mazur introduced the Iwasawa theory of Selmer groups of Abelian varieties and described the growth of the size of the p^∞ -Selmer group in \mathbb{Z}_p -extensions F_∞/F . For an Abelian variety A/F , the dual Selmer group over F_∞ , denoted by $X(A/F_\infty)$, is a finitely generated $\Lambda(\Gamma)$ -module; here $\Gamma = \text{Gal}(F_\infty/F) \simeq \mathbb{Z}_p$ and $\Lambda(\Gamma)$ is the associated Iwasawa algebra. However, $X(A/F_\infty)$ is *not always* $\Lambda(\Gamma)$ -torsion, depending on the reduction type at p .

When $X(A/F_\infty)$ is $\Lambda(\Gamma)$ -torsion, it affords a structure theorem like in the classical case. But an analogue of the classical conjecture is known to be false. For the cyclotomic extension of \mathbb{Q} , there are examples of elliptic curves where the associated μ -invariant is positive at a prime of good ordinary reduction.

In [3], Coates and Sujatha studied a subgroup of the Selmer group, called the *fine Selmer group*. They made the following conjecture.

Date: December 13, 2019.

2010 Mathematics Subject Classification. Primary 11R23.

Key words and phrases. Iwasawa Theory, Fine Selmer groups, Conjecture A, Classical $\mu = 0$ Conjecture.

Conjecture A. *Let p be an odd prime and E be an elliptic curve over a number field F . When $F_\infty = F_{\text{cyc}}$, the Pontryagin dual of the fine Selmer group, denoted $Y(E/F_{\text{cyc}})$, is a finitely generated \mathbb{Z}_p -module i.e. $Y(E/F_{\text{cyc}})$ is $\Lambda(\Gamma)$ -torsion and the associated μ -invariant, $\mu(Y(E/F_{\text{cyc}})) = 0$.*

A priori, a relation between Galois modules coming from class groups and those coming from elliptic curves is not obvious. However, there is growing evidence that in a cyclotomic tower, fine Selmer groups and class groups have similar growth patterns [3], [10]. We provide more evidence for this by proving the following theorems.

Theorem 1.1. *Let F be a number field such that the Classical Conjecture holds. For E an elliptic curve over F with $E(F)[p] \neq 0$, Conjecture A holds for $Y(E/F_{\text{cyc}})$.*

Remark

- (i) [3, Corollary 3.6] is a particular case of our theorem when $F \supset \mu_p$. Indeed, this is a consequence of the following two facts. First, if F/\mathbb{Q} is an Abelian extension, so is $F(\mu_p)/\mathbb{Q}$ and the Classical Conjecture holds. Second, for finite extensions of number fields L/F , Conjecture A for $Y(E/L_{\text{cyc}})$ implies Conjecture A for $Y(E/F_{\text{cyc}})$.
- (ii) Tools from Galois cohomology are used to provide new evidence for the Classical Conjecture in Section 4. An application of Theorem 1.1 provides new evidence for Conjecture A.

A converse of Theorem 1.1 is true: given a number field F , the Classical Conjecture holds for F_{cyc}/F , if there exists *one* elliptic curve E/F , with $E(F)[p] \neq 0$ for which Conjecture A holds. It is known that the Classical Conjecture holds if there exists one elliptic curve E/F , with $E(F)[p] \neq 0$ for which the dual *Selmer group* is $\Lambda(\Gamma)$ -torsion and the corresponding μ -invariant is 0 (see [2], [9]). Our result weakens the hypothesis significantly and is proved using a different machinery.

Theorem 1.2. *Let E be an elliptic curve defined over the number field F . Let p be any odd prime. Further assume that $E(F)[p] \neq 0$. If Conjecture A holds for $Y(E/F_{\text{cyc}})$, then the Classical Conjecture holds for F_{cyc}/F .*

Theorems 1.1 and 1.2 prove isogeny invariance of Conjecture A in some previously unknown cases (see [17]).

Corollary 1.3. *Let F be a number field containing μ_p or be a totally real field. Let E and E' be isogenous elliptic curves such that both E and E' have non-trivial p -torsion points over F . Then, Conjecture A holds for $Y(E/F_{\text{cyc}})$ if and only if Conjecture A holds for $Y(E'/F_{\text{cyc}})$.*

Remark. All statements hold for Abelian varieties of dimension d . The only property of cyclotomic \mathbb{Z}_p -extensions required in the proofs is that primes in a certain finite set decompose finitely. The theorems are stated for elliptic curves over the cyclotomic \mathbb{Z}_p -extension as the original conjectures are in this setting.

2. PRELIMINARIES

Throughout this paper, F will denote a number field and p an odd prime.

Let A/F be a d -dimensional Abelian variety and S be a finite set of primes of F containing the Archimedean primes, primes above p , and primes where A has bad reduction. Fix an algebraic closure \bar{F}/F and set F_S to be the maximal subfield of

\overline{F} containing F which is unramified outside S . Denote the absolute Galois group $\text{Gal}(\overline{F}/F)$ by G_F and the Galois group $\text{Gal}(F_S/F)$ by $G_S(F)$.

Definitions of p^∞ -Selmer group and p^∞ -fine Selmer group are as in [19].

Definition 2.1. *The p -fine Selmer group, with respect to the finite set S , is*

$$(1) \quad R_S(A[p]/F) := \ker \left(H^1(G_S(F), A[p]) \rightarrow \bigoplus_{v \in S} H^1(F_v, A[p]) \right).$$

The p^∞ -fine Selmer group is

$$(2) \quad R(A/F) := \ker \left(H^1(G_S(F), A[p^\infty]) \rightarrow \bigoplus_{v \in S} H^1(F_v, A[p^\infty]) \right).$$

The definition of $R(A/F)$ is independent of S . For a \mathbb{Z}_p -extension F_∞/F ,

$$R(A/F_\infty) = \varinjlim_L R(A/L)$$

where the inductive limit is over all finite extensions L/F contained in F_∞ .

Recall, the Pontryagin dual of a discrete p -primary (resp. compact pro- p) Abelian group is a compact (resp. discrete) module over the associated Iwasawa algebra. For G a profinite group and M a G -module, M^G is the subgroup of elements fixed by G and M_G is the largest quotient of M with trivial G action.

Definition 2.2. *For an Abelian group N , its p -rank is the $\mathbb{Z}/p\mathbb{Z}$ -dimension of $N[p]$, denoted by $r_p(N)$. If G is a pro- p group, write $h_i(G) = r_p(H^i(G, \mathbb{Z}/p))$.*

Lemma 2.3. [10, Lemma 3.1] *Let G be a pro- p group and M be a discrete G -module which is co-finitely generated over \mathbb{Z}_p . If $h_1(G)$ is finite, $r_p(H^1(G, M))$ is finite. Furthermore, the following inequalities hold*

$$\begin{aligned} h_1(G)r_p(M^G) - r_p((M/M^G)^G) &\leq r_p(H^1(G, M)) \\ &\leq h_1(G)(\text{corank}_{\mathbb{Z}_p}(M) + \log_p(|M/M_{\text{div}}|)) \end{aligned}$$

Lemma 2.4. [10, Lemma 3.2] *Consider an exact sequence of co-finitely generated Abelian groups,*

$$W \rightarrow X \rightarrow Y \rightarrow Z.$$

Then

$$|r_p(X) - r_p(Y)| \leq 2r_p(W) + r_p(Z).$$

Definition 2.5. *The p -Hilbert S -class field of F , denoted $H_S(F)$, is the maximal Abelian unramified p -extension of F in which all primes in S split completely. By class field theory, the Galois group $\text{Gal}(H_S(F)/F) = \text{Cl}_S(F)$, is the S -class group.*

Lemma 2.6. [10, Lemma 5.2, 5.3] *Let F_{cyc}/F be the cyclotomic \mathbb{Z}_p -extension and F_n be the subfield of F_{cyc} such that $[F_n : F] = p^n$. Let A be an Abelian variety over F and S be as defined before. Then*

$$(3) \quad \left| r_p(\text{Cl}(F_n)) - r_p(\text{Cl}_S(F_n)) \right| = O(1),$$

$$(4) \quad \left| r_p(R_S(A[p]/F_n)) - r_p(R(A/F_n)) \right| = O(1).$$

3. PROOF OF THE MAIN RESULTS

We first prove Corollary 1.3. It follows from the main theorems in Section 1.

Proof of Corollary 1.3. Let F be a number field that contains μ_p or F/\mathbb{Q} be a totally real field. Let E be an elliptic curve isogenous to E' over F with the additional property that both $E(F)[p], E'(F)[p]$ are non-trivial. WLOG if Conjecture A holds for $Y(E/F_{\text{cyc}})$ then by Theorem 1.2 the Classical Conjecture holds for F_{cyc}/F . Now by Theorem 1.1 Conjecture A holds for $Y(E'/F_{\text{cyc}})$. This proves the corollary. \square

3.1. Proof of Theorem 1.2. Theorem 1.2 follows from the following lemma when A is an elliptic curve.

Lemma 3.1. *Let F_{cyc}/F be the cyclotomic \mathbb{Z}_p -extension and F_n be the subfield of F_{cyc} such that $[F_n : F] = p^n$. Let A be a d -dimensional Abelian variety over F and S be as defined before. Assume $A(F)[p] \neq 0$. Then for some positive constant k_1 that depends on $A(F)[p]$,*

$$(5) \quad k_1 r_p(\text{Cl}_S(F_n)) \leq r_p(R(A/F_n)) + O(1)$$

Proof. For the ease of notation, set $H_n = H_S(F_n)$ and $H_{n,w} = H_S(F_n)_w$. Consider the following commutative diagram:

$$\begin{array}{ccccccc} 0 & \rightarrow & R(A/F_n) & \rightarrow & H^1(G_S(F_n), A[p^\infty]) & \rightarrow & \bigoplus_{v_n} H^1(F_{n,v_n}, A[p^\infty]) \\ & & \downarrow r_n & & \downarrow f_n & & \downarrow \gamma_n \\ 0 & \rightarrow & R(A/H_n) & \rightarrow & H^1(G_S(H_n), A[p^\infty]) & \rightarrow & \bigoplus_{v_n} \bigoplus_{w|v_n} H^1(H_{n,w}, A[p^\infty]) \end{array}$$

Here v_n runs over all primes in $S(F_n)$, the finite set of primes in F_n that lie above the primes in S . Observe

$$\ker \gamma_n = \bigoplus_{v_n} \ker \gamma_{n,v_n}.$$

Each $\ker \gamma_{n,v_n} = H^1(G_{n,v_n}, A(H_{n,v_n})[p^\infty])$ where G_{n,v_n} is the decomposition group of $G_n := \text{Gal}(H_n/F_n)$. By definition of p -Hilbert S -class field, $G_{n,v_n} = 1$. So, $\ker \gamma_n = \text{coker } \gamma_n = 0$.

By inflation-restriction, $\ker(f_n) = H^1(G_n, A(H_n)[p^\infty])$ and by diagram chasing, one obtains $\ker(f_n) \hookrightarrow R(A/F_n)$. Thus,

$$r_p\left(H^1(G_n, A(H_n)[p^\infty])\right) \leq r_p(R(A/F_n)).$$

Combining this with Lemma 2.3 gives the following inequality

$$(6) \quad h_1(G_n) r_p(A(F_n)[p^\infty]) - 2d \leq r_p(R(A/F_n)).$$

By definition of S -class group, $\text{Gal}(H_n/F_n) = \text{Cl}_S(F_n)$. So

$$\begin{aligned} h_1(G_n) &= h_1(\text{Gal}(H_n/F_n)) \\ &= r_p(\text{Cl}_S(F_n)/p) \\ &= r_p(\text{Cl}_S(F_n)) \end{aligned}$$

where the last equality follows from the finiteness of the S -class group. Also,

$$\begin{aligned} r_p(A(F_n)[p^\infty]) &\geq r_p(A(F)[p^\infty]) \\ &= r_p(A(F)[p]). \end{aligned}$$

By Inequality 6 and the above discussion it follows,

$$(7) \quad r_p(A(F)[p]) r_p(\text{Cl}_S(F_n)) \leq r_p(R(A/F_n)) + O(1).$$

This proves the lemma as the hypothesis forces $r_p(A(F)[p]) \neq 0$. \square

We now provide a proof of Theorem 1.2.

Proof. By an application of [18, Lemma 13.20], Conjecture A holds for $Y(E/F_{\text{cyc}})$ if and only if $r_p(R(E/F_n)) = O(1)$. In other words, Conjecture A holds if and only if the p -rank remains bounded in the cyclotomic tower.

By hypothesis, Conjecture A holds for $Y(E/F_{\text{cyc}})$ so $r_p(R(E/F_n)) = O(1)$. Also by hypothesis, $E(F)[p] \neq 0$. Inequality 7 implies $r_p(\text{Cl}_S(F_n))$ is bounded independent of n . By Equation 3, so is $r_p(\text{Cl}(F_n))$.

This is enough to prove the Classical Conjecture. Indeed, the Classical Conjecture holds for F_{cyc}/F if and only if $r_p(\text{Cl}(F_n)/p)$ is bounded independent of n [18, Proposition 13.23]. Since class groups are finite, $r_p(\text{Cl}(F_n)) = r_p(\text{Cl}(F_n)/p)$. Thus, the Classical Conjecture is equivalent to $r_p(\text{Cl}(F_n))$ being independent of n . This finishes the proof. \square

3.2. Proof of Theorem 1.1. Recall the following well-known facts [8]. If L/F is a p -power Galois extension, $\mu(F_{\text{cyc}}/F) = 0$ implies $\mu(L_{\text{cyc}}/L) = 0$. On the other hand, for any extension L/F , $\mu(L_{\text{cyc}}) = 0$ implies $\mu(F_{\text{cyc}}) = 0$.

Case (i) Suppose $F \supset \mu_p$.

Since $E(F)[p] \neq 0$, it follows from the Weil pairing that $F(E[p])/F$ is either trivial or cyclic of order p . This is precisely the situation of [10, Theorem 5.5]. There is nothing left to prove.

Case(ii) Suppose $F \not\supset \mu_p$.

Theorem 1.1 follows from the following inequality where k_2 is a positive constant,

$$(8) \quad r_p(R(E/F_n)) \leq k_2 r_p(\text{Cl}(F_n)) + O(1).$$

Indeed, the Classical Conjecture is equivalent to $r_p(\text{Cl}(F_n))$ being bounded independent of n [18, Proposition 13.23]. If the Classical Conjecture holds, Inequality 8 for an elliptic curve E implies $r_p(R(E/F_n))$ is bounded independent of n . Therefore, Conjecture A holds.

Observe that by Equations 3 and 4, Theorem 1.1 follows from the following variant of Inequality 8,

$$(9) \quad r_p(R_S(E[p]/F_n)) \leq k_2 r_p(\text{Cl}_S(F_n)) + O(1).$$

Define $R_S(E(F_n)[p]/F_n)$ by replacing $E[p]$ with $E(F_n)[p]$ in Equation 1. $G_S(F_n)$ acts trivially on $E(F_n)[p]$; hence it is possible to relate $R_S(E(F_n)[p]/F_n)$ with $\text{Cl}_S(F_n)$ and similarly their p -ranks. Since the Galois action is trivial,

$$H^1(G_S(F_n), E(F_n)[p]) = \text{Hom}(G_S(F_n), E(F_n)[p])$$

and there are similar identifications for the local cohomology groups. It follows

$$R_S(E(F_n)[p]/F_n) = \text{Hom}(\text{Cl}_S(F_n), E(F_n)[p]) \simeq \text{Cl}_S(F_n)[p]^{r_p(E(F_n)[p])}$$

where the isomorphism is as Abelian groups. This gives the following inequality

$$\begin{aligned} r_p(R_S(E(F_n)[p]/F_n)) &= r_p(E(F_n)[p]) r_p(\text{Cl}_S(F_n)) \\ &\leq 2r_p(\text{Cl}_S(F_n)). \end{aligned}$$

Note that Inequality 9 follows from the above inequality provided the p -ranks of $R_S(E[p]/F_n)$ and $R_S(E(F_n)[p]/F_n)$ have the same order of growth in F_{cyc} . This is the content of the following lemma. This completes the proof of Theorem 1.1.

Lemma 3.2. *Let F be a totally real number field. Let E be an elliptic curve over F with $E(F)[p] \neq 0$. Let F_{cyc}/F be the cyclotomic \mathbb{Z}_p -extension and suppose the Classical $\mu = 0$ Conjecture holds for F_{cyc} . Let S be as defined before. Then*

$$(10) \quad \left| r_p(R_S(E(F_n)[p]/F_n)) - r_p(R_S(E[p]/F_n)) \right| = O(1).$$

Proof. Notice that if $E(F)[p] = E[p]$, then the Lemma is trivial. The focus is on the case $E(F)[p] \neq 0, E[p]$. Set $B_n = E(F_n)[p]$. Consider the commutative diagram

$$\begin{array}{ccccccc} 0 & \rightarrow & R_S(B_n/F_n) & \rightarrow & H^1(G_S(F_n), B_n) & \rightarrow & \bigoplus_{v_n} H^1(F_{n,v_n}, B_n) \\ & & \downarrow s_n & & \downarrow f_n & & \downarrow g_n \\ 0 & \rightarrow & R_S(E[p]/F_n) & \rightarrow & H^1(G_S(F_n), E[p]) & \rightarrow & \bigoplus_{v_n} H^1(F_{n,v_n}, E[p]) \end{array}$$

where v_n runs over all the primes in the finite set $S(F_n)$.

By hypothesis, E has an F_n -rational p -torsion point. This gives the short exact sequence

$$(11) \quad 0 \rightarrow B_n \rightarrow E[p] \rightarrow \mu_p \rightarrow 0.$$

This is because, if E has an F_n -rational p -torsion point, this point gives an injection $\mathbb{Z}/p\mathbb{Z} \hookrightarrow E[p]$. Therefore,

$$0 \rightarrow \mathbb{Z}/p\mathbb{Z} \rightarrow E[p] \rightarrow M \rightarrow 0.$$

By Cartier duality and the Weil pairing, the above short exact sequence turns into

$$0 \rightarrow M^\vee \rightarrow E[p] \rightarrow \mu_p \rightarrow 0,$$

where μ_p is viewed as a quotient of $E[p]$. Since the Weil pairing is alternating, the orthogonal complement of $\mathbb{Z}/p\mathbb{Z}$ is $\mathbb{Z}/p\mathbb{Z}$, thus $M^\vee = \mathbb{Z}/p\mathbb{Z}$ as a subgroup of $E[p]$.

Taking the $G_S(F_n)$ -cohomology of Equation 11, $\ker(f_n) \subseteq H^0(G_S(F_n), \mu_p)$. Since μ_p is finite, therefore $r_p(\ker(f_n)) = O(1)$ and hence $r_p(\ker(s_n)) = O(1)$. A similar argument for the local cohomology gives $r_p(\ker(g_n)) = O(1)$.

By Lemma 2.3 applied to the map s_n ,

$$\begin{aligned} \left| r_p(R_S(B_n/F_n)) - r_p(R_S(E[p]/F_n)) \right| &\leq 2r_p(\ker(s_n)) + r_p(\text{coker}(s_n)) \\ &= r_p(\text{coker}(s_n)) + O(1). \end{aligned}$$

If $r_p(\text{coker}(s_n)) = O(1)$, the proof is complete. Observe, $\text{coker}(f_n) \subseteq H^1(G_S(F_n), \mu_p)$ and $\text{coker}(g_n) \subseteq \bigoplus_{v_n} H^1(F_{n,v_n}, \mu_p)$. Further, note that

$$r_p \left(\ker \left(H^1(G_S(F_n), \mu_p) \rightarrow \bigoplus_{v_n} H^1(F_{n,v_n}, \mu_p) \right) \right) = O(1) \Rightarrow r_p(\text{coker}(s_n)) = O(1).$$

For ease of notation, refer to the kernel as a *fine Selmer group*, $R_S(\mu_p/F_n)$.

By hypothesis, the Classical Conjecture holds for F_{cyc}/F . By [17, Proposition 4.10(1)] it is equivalent to $H^2(G_S(F_{\text{cyc}}), \mu_p) = 0$. This latter statement is often

referred to as *Conjecture A for μ_p* . Using equivalent reformulations (see [3], [17]),

$$\begin{aligned} H^2(G_S(F_{\text{cyc}}), \mu_p) = 0 &\Leftrightarrow \varprojlim_n \left(H^2(G_S(F_n), \mu_p) \right) \text{ is finite} \\ &\Leftrightarrow R_S(\mu_p/F_{\text{cyc}}) \text{ is finite} \\ &\Leftrightarrow \varinjlim_n R_S(\mu_p/F_n) \text{ is finite} \\ &\Leftrightarrow R_S(\mu_p/F_n) \text{ is finite and bounded} \end{aligned}$$

In particular, $r_p(\text{coker}(s_n)) = O(1)$ which is what we needed to prove. \square

4. ILLUSTRATING THE RESULTS WITH EXAMPLES

In this section, we show that the Classical Conjecture holds for p -rational number fields. This allows us to provide evidence for Conjecture A.

Remark. For *totally real* p -rational fields, the classical Iwasawa module is trivial [13]. In particular, classical $\mu = \lambda = 0$ for such fields. It appears that a proof of the Classical Conjecture for *all* p -rational fields has not been written down in literature. We suspect it is because examples of non-Abelian p -rational fields have come to light only recently (see [1]).

Let F be a number field. Let S be a finite set of primes of F *containing* the primes above p and the Archimedean primes. The weak Leopoldt conjecture in the classical setting is the assertion

$$(12) \quad H^2(\text{Gal}(F_S/F_{\text{cyc}}), \mathbb{Q}_p/\mathbb{Z}_p) = 0.$$

It holds for the cyclotomic extension of a number field [12, Theorem 10.3.25]. If Equation 12 holds for a finite set S as mentioned above, it holds for the set $S = \Sigma = S_p \cup S_\infty$ where S_p is the set of primes of F above p and S_∞ are the Archimedean primes [12, Theorem 11.3.2]. Therefore, the weak Leopoldt Conjecture is independent of the choice of S . From here on, fix $S = \Sigma$.

Let F_{S_p} denote the maximal p -ramified extension of F . Consider the Galois group $\text{Gal}(F_{S_p}/F)$ and let $\mathcal{G}_{S_p}(F) = \text{Gal}(F_{S_p}(p)/F)$ be its maximal pro- p quotient.

Definition 4.1. [13] *Let F be a number field, p be an odd prime. F is called p -rational if and only if $\mathcal{G}_{S_p}(F)$ is pro- p -free.*

Given a number field F , it is conjectured to be p -rational for all primes outside a set of Dirichlet density 0 [6].

The following theorem is well-known.

Theorem 4.2. [12, Theorem 11.3.7] *The Classical Conjecture holds for F_{cyc} if and only if $\mathcal{G}_\Sigma(F_{\text{cyc}}) = \text{Gal}(F_\Sigma(p)/F_{\text{cyc}})$ is a free pro- p group.*

Definition 4.3. [14, Page 23] *A pro- p group G is free if and only if its p -cohomological dimension $\text{cd}_p(G) \leq 1$.*

By a standard fact in Galois cohomology of pro- p groups [14, Chapter I, Section 4, Proposition 21], an equivalent formulation of Theorem 4.2 is the following: the Classical Conjecture holds for F_{cyc} if and only if

$$(13) \quad H^2(\mathcal{G}_\Sigma(F_{\text{cyc}}), \mathbb{Z}/p\mathbb{Z}) = 0.$$

Corollary 4.4. *Let p be an odd prime and F be a p -rational number field. The Classical Conjecture holds for F_{cyc}/F .*

Proof. Since $p \neq 2$, we replace S_p by Σ in the definition of p -rational fields. For p -rational number fields, $\mathcal{G}_\Sigma(F) = \text{Gal}(F_\Sigma(p)/F)$ has p -cohomological dimension at most 1. Equivalently,

$$H^2(\mathcal{G}_\Sigma(F), \mathbb{Z}/p\mathbb{Z}) = 0.$$

Since $\mathcal{G}_\Sigma(F_{\text{cyc}})$ is a closed normal subgroup of $\mathcal{G}_\Sigma(F)$; by [14, Proposition 14]

$$\text{cd}_p(\mathcal{G}_\Sigma(F_{\text{cyc}})) \leq \text{cd}_p(\mathcal{G}_\Sigma(F)) \leq 1.$$

Thus,

$$H^2(\mathcal{G}_\Sigma(F_{\text{cyc}}), \mathbb{Z}/p\mathbb{Z}) = 0.$$

By Equation 13, the result follows. \square

This provides new evidence for Conjecture A.

Corollary 4.5. *Let F be a p -rational number field such that either*

- (i) $F \supseteq \mu_p$ or
- (ii) F is a totally real number field.

Suppose E is an elliptic curve over F with $E(F)[p] \neq 0$. Then Conjecture A holds for $Y(E/F_{\text{cyc}})$.

Proof. This follows from Theorem 1.1 along with Corollary 4.4. \square

In special cases, a proof of Conjecture A can be provided *independent* of the Classical Conjecture. For this, we need the following equivalent formulation of Conjecture A, established independently by Greenberg [5] and Sujatha [16].

Proposition 4.6. *Assume the weak Leopoldt conjecture for elliptic curves holds, i.e. $H^2(G_S(F_{\text{cyc}}), E[p^\infty]) = 0$. Then Conjecture A for $Y(E/F_{\text{cyc}})$ is equivalent to the assertion*

$$H^2(G_S(F_{\text{cyc}}), E[p]) = 0.$$

Proposition 4.7. *Let p be an odd prime, F be a p -rational field, and E be an elliptic curve with good reduction everywhere over F (or bad reduction at primes above p) such that $E[p] \subset E(F)$. Then Conjecture A holds for $Y(E/F_{\text{cyc}})$.*

Proof. Choose $S = \Sigma = S_p \cup S_\infty$. By p -rationality of F and the isomorphism of the inflation map [12, Corollary 10.4.8], we have

$$(14) \quad H^2(\mathcal{G}_\Sigma(F), E[p]) = H^2(G_\Sigma(F), E[p]) = 0.$$

By an application of the Hochschild-Serre Spectral sequence, we obtain the following exact sequence [12, Page 119]

$$H^2(\mathcal{G}_\Sigma(F), E[p]) \rightarrow H^0(\Gamma, H^2(\mathcal{G}_\Sigma(F_{\text{cyc}}), E[p])) \rightarrow 0,$$

where $\Gamma = \text{Gal}(F_{\text{cyc}}/F)$. The first term is 0, thus $H^0(\Gamma, H^2(\mathcal{G}_\Sigma(F_{\text{cyc}}), E[p]))$ is trivial. Since $H^2(\mathcal{G}_\Sigma(F_{\text{cyc}}), E[p])$ is a discrete module, it must be 0. Once again by the isomorphism of the inflation map,

$$0 = H^2(\mathcal{G}_\Sigma(F_{\text{cyc}}), E[p]) = H^2(G_\Sigma(F_{\text{cyc}}), E[p]).$$

By Proposition 4.6, Conjecture A holds for $Y(E/F_{\text{cyc}})$. \square

Remark. It appears to be hard to weaken the hypothesis $E[p] \subseteq F$, in proving Conjecture A independent of the Classical Conjecture.

ACKNOWLEDGEMENTS

The author thanks Kumar Murty for his support and many helpful suggestions and discussions over the course of preparation of this article. Lots of thanks to Sujatha Ramadorai and Prof. Nguyen Quang Do for discussions on p -rational fields; to Kestutis Cesnavicius, Gaurav Patil, and Meng Fai Lim for answering questions. Thanks also to Ali Altug for his questions when initial results were presented at the BU-Keio Number Theory Conference (2019) which helped improve the exposition.

REFERENCES

1. Razvan Barbucescu and Jishnu Ray, *Some remarks and experiments on Greenberg's p -rationality conjecture*, arXiv preprint arXiv:1706.04847 (2017).
2. Kestutis Cesnavicius, *Selmer groups and class groups*, Compositio Mathematica **151** (2015), no. 3, 416–434.
3. John Coates and Ramdorai Sujatha, *Fine Selmer groups of elliptic curves over p -adic Lie extensions*, Mathematische Annalen **331** (2005), no. 4, 809–839.
4. Bruce Ferrero and Lawrence C Washington, *The Iwasawa invariant μ_p vanishes for abelian number fields*, Annals of Mathematics (1979), 377–395.
5. Ralph Greenberg, *Iwasawa theory, projective modules, and modular representations*, American Math. Soc., 2010.
6. ———, *Galois representations with open image*, Annales mathématiques du Québec **40** (2016), no. 1, 83–119.
7. Kenkichi Iwasawa, *On Γ -extensions of algebraic number fields*, Bulletin of the American Mathematical Society **65** (1959), no. 4, 183–226.
8. ———, *On the μ -invariants of \mathbb{Z}_l -extensions*, Algebraic Geometry and Commutative Algebra (1973), 1–11.
9. Meng Fai Lim and V Kumar Murty, *Growth of selmer groups of cm abelian varieties*, Canadian Journal of Mathematics **67** (2015), no. 3, 654–666.
10. ———, *The growth of fine Selmer groups*, J. Ramanujan Math. Society **31** (2016), no. 1, 79–94.
11. Barry Mazur, *Rational points of abelian varieties with values in towers of number fields*, Inventiones mathematicae **18** (1972), no. 3-4, 183–266.
12. Jürgen Neukirch, Alexander Schmidt, and Kay Wingberg, *Cohomology of number fields, volume 323 of grundlehren der mathematischen wissenschaften [fundamental principles of mathematical sciences]*, 2008.
13. T Nguyen Quang Do and A Movahhedi, *On the arithmetic of p -rational number fields*, Séminaire de Théorie des Nombres, Paris 1987–88, 1990, pp. 155 to –200.
14. Jean-Pierre Serre, *Galois cohomology*, Springer Science & Business Media, 2013.
15. Warren Sinnott, *On the μ -invariant of the Γ -transform of a rational function*, Inventiones mathematicae **75** (1984), no. 2, 273–282.
16. R Sujatha, *Elliptic curves and iwasawa's $\mu = 0$ conjecture*, Quadratic Forms, Linear Algebraic Groups, and Cohomology, Springer, 2010, pp. 125–135.
17. R Sujatha and M Witte, *Fine Selmer groups and isogeny invariance*, Conference on Geometry, Algebra, Number Theory, and their Information Technology Applications, Springer, 2016, pp. 419–444.
18. Lawrence C Washington, *Introduction to cyclotomic fields*, vol. 83, Springer Science & Business Media, 1997.
19. Christian Wuthrich, *The fine Selmer group and height pairings*, Ph.D. thesis, University of Cambridge, 2004.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, BAHEN CENTRE, 40 ST. GEORGE ST., ROOM 6290, TORONTO, ONTARIO, CANADA, M5S 2E4
E-mail address: dkundu@math.utoronto.ca