

ಏಪ್ರಿಲ್

- ಬಿಳಿ ಕಾಂಡ ಕೊರಕ ಪೀಡಿತ ಅರೇಬಿಕಾ ಗಿಡಗಳನ್ನು ಗುರುತಿಸಿ, ಮಟ್ಟ ಮಾಡುವುದು/ಬೇರು ಸಹಿತ ಕಿತ್ತು ಸುಟ್ಟು ಹಾಕುವ ಕೆಲಸ ಮುಗಿಸಬೇಕು
- ಎಲೆ ತುಕ್ಕು ರೋಗಕ್ಕೆ ಕಾಂಟಾಪ್ (400 ಮಿಲಿ / 200 ಲೀಟರ್ ನೀರು) ಅಥವಾ ಬೆಲಟಾನ್ (160 ಗ್ರಾಂ / 200 ಲೀಟರ್ ನೀರು) ಸಿಂಪರಣೆ ಮುಗಿಸಬೇಕು
- ಅರೇಬಿಕಾ ಕಾಂಡ ಹಾಗೂ ತಾಯಿ ರೆಕ್ಕೆಗಳಿಗೆ ಕ್ಲೋರೋಪೈರಿಪಾಸ್ (600 ಮಿಲಿ / 200 ಲೀಟರ್ ನೀರು) ಸಿಂಪಡಿಸುವುದು
- ಸಾಕಷ್ಟು ತೇವಾಂಶವಿದ್ದಲ್ಲಿ, ಕಾಂಪೋಸ್ಟ್ ಅಥವಾ ಸಾವಯವ ಗೊಬ್ಬರ ಹಾಕುವುದು
- ಪಾತಿಯಲ್ಲಿ ಗಿಡಗಳ ಕಾಂಡಗಳ ಸಾಯುವಿಕೆ ಕಂಡುಬಂದಲ್ಲಿ, ಟೆಲ್ಟ್ (0.8 ಮಿಲಿ / ಲೀಟರ್ ನೀರು) ತಿಂಗಳಿಗೊಮ್ಮೆ ಗಿಡಗಳನ್ನು ನೆಡಲು ಬಳಸುವವರೆಗೆ ಸಿಂಪಡಿಸಬೇಕು
- ಹೊಸ ತೋಟಗಳನ್ನು ಮಾಡುವಲ್ಲಿ ಸಾಲು ಹಿಡಿದು ಗುಂಡಿಗಳನ್ನು ತೆಗೆಯುವುದು

April

- **Complete tracing and collar pruning/Uprooting/Burning of White Stem Borer affected Arabica plants. If stumps to be stored, immerse them in water for 10 days**
- **Complete the spraying of Baylaton (160 gram/200 litres of water) or Contaf (400 ml / 200 litres of water) for susceptible varieties like S. 795 and Cauvery to control leaf rust**
- **Spray chloropyriphos (600 ml / 200 litres of water) to main stem and thick primary branches of Arabica plants**
- **Apply compost or organic manures, if sufficient moisture is available in soil**
- **Spray Dithane M-45 or Indofil M-45 @ 5 gram / liter if cercospora leaf spot is noticed**
- **Spray Tilt (0.8 ml per liter), if stem necrosis is noticed. Repeat spray once in a month till they attain field planting stage**
- **Take up line marking and opening of pits in new clearings**