

## ಸೆಪ್ಟೆಂಬರ್

- ಹೊಸ ತೋಟಗಳಲ್ಲಿ ಕಾಫಿ ಹಾಗೂ ಮರಗಳ ಸಸಿಗಳನ್ನು ನೆಡುವುದು
- ಎಲೆ ತುಕ್ಕು ರೋಗ ನಿಯಂತ್ರಣಕ್ಕಾಗಿ ಹಿಂಗಾರಿನ 0.5 % ಬೋಡೊ ದ್ರಾವಣ ಅಥವಾ ಕಾಂಟಾಫ್ (400 ಮಿಲಿ/200 ಲೀಟರ್ ನೀರು) ಸಿಂಪಡಿಸುವುದು
- ಬಿಳಿ ಕಾಂಡ ಕೊರಕ ಪೀಡಿತ ಅರೇಬಿಕಾ ಗಿಡಗಳನ್ನು ಗುರುತಿಸಿ, ಮಟ್ಟ ಮಾಡುವುದು/ಬೇರು ಸಹಿತ ಕಿತ್ತು ಸುಟ್ಟು ಹಾಕುವುದು. ಕಾಂಡ ಹಾಗೂ ತಾಯಿ ರೆಕ್ಕೆಗಳಿಗೆ ಕ್ಲೋರೋಪೈರಿಫಾಸ್ (600 ಮಿಲಿ / 200 ಲೀಟರ್ ನೀರು) ಸಿಂಪಡಿಸುವುದು
- ಹಿಂಗಾರಿನ ಗೊಬ್ಬರ ಹಾಕುವುದು
- ಕಾಯಿ ಕೊರಕ ಪೀಡಿತ ಕಾಯಿಗಳನ್ನು ತೆಗೆದು ಸುಟ್ಟು ಹಾಕುವುದು, ಬ್ರೋಕಾ ಟ್ರಾಪ್ಸ್ ಅಳವಡಿಸುವುದು
- ರೆಕ್ಕೆ ಬೋರಲು ಪೀಡಿತ ರೆಕ್ಕೆಗಳನ್ನು ಗುರುತಿಸಿ ತೆಗೆದು ಸುಟ್ಟು ಹಾಕುವುದು. ರೊಬಸ್ಟ್ ಸಣ್ಣ ಗಿಡಗಳಿಗೆ ಟಿಲ್ಟ್ (160 ಮಿಲಿ / 200 ಲೀಟರ್ ನೀರು) ರೆಕ್ಕೆ ಬೋರಲು ನಿಯಂತ್ರಣಕ್ಕಾಗಿ ಸಿಂಪಡಿಸುವುದು

## September

- Planting coffee and shade trees saplings in young clearings
- Spray 0.5 % Bordeaux Mixture or Contaf (@ 400 ml / 200 litres) against leaf rust during post monsoon
- Tracing and collar pruning/Uprooting/Burning of White Stem Borer affected Arabica plants. Spray chloropyriphos (600 ml / 200 litres of water) to main stem and thick primary branches of Arabica plants
- Post monsoon fertilizer application
- Collect and burn berry borer infested berries, install berry borer traps
- Trace, collect and burn shot hole affected branches, spray Tilt (160 ml / 200 litres) in young robusta plants to control against shot hole borer