

## ಮಾರ್ಚ್

- ಬಿಳಿ ಕಾಂಡ ಕೊರಕ ಪೀಡಿತ ಅರೇಬಿಕಾ ಗಿಡಗಳನ್ನು ಗುರುತಿಸಿ, ಮಟ್ಟು ಮಾಡುವುದು/ಬೇರು ಸಹಿತ ಕಿತ್ತು ಸುಟ್ಟು ಹಾಕುವುದು. ಬಡ್ಡೆಗಳನ್ನು ಶೇಖರಿಸುವುದಾದಲ್ಲಿ, ಕಿತ್ತ ಬಡ್ಡೆಗಳನ್ನು ಹತ್ತು ದಿನಗಳ ಕಾಲ ನೀರಿನಲ್ಲಿ ನೆನೆ ಹಾಕಬೇಕು
- ಎಲೆ ತುಕ್ಕು ರೋಗಕ್ಕೆ ಸುಲಭವಾಗಿ ತುತ್ತಾಗುವ ಕಾವೇರಿ, ಎಸ್. 795 ತಳಿಗಳಿಗೆ ಕಾಂಟಾಪ್ (400 ಮಿಲಿ / 200 ಲೀಟರ್ ನೀರು) ಅಥವಾ ಬೆಲಟಾನ್ (160 ಗ್ರಾಂ / 200 ಲೀಟರ್ ನೀರು) ಸಿಂಪಡಿಸಿ
- ರೊಬಸ್ತಾ ಕೊಯ್ಲು ಹಾಗೂ ಸಂಸ್ಕರಣೆ ಮುಗಿಸುವುದು
- ಅರೇಬಿಕಾ ಹಾಗೂ ರೊಬಸ್ತಾ ಗಿಡ ಕಸ ಪೂರ್ಣಗೊಳಿಸುವುದು
- ರೊಬಸ್ತಾದಲ್ಲಿ ಬ್ಯಾಕ್ಟಿಂಗ್ ನೀರಾವರಿ ನೀಡುವುದು
- ಹೊಸ ತೋಟಗಳಲ್ಲಿ ಸಣ್ಣ ಕಾಫಿ ಗಿಡಗಳಿಗೆ ನೀರಾವರಿ
- ಕಂಬಳಿ ಹುಳುಗಳ ಹಾವಳಿ ಇರುವಲ್ಲಿ, ಕೋಶಾವಸ್ಥೆ ಹಾಗೂ ಮೊಟ್ಟೆಗಳನ್ನು ಆಯ್ದು ಸುಟ್ಟು ಹಾಕುವುದು

## March

- Tracing and collar pruning/Uprooting/Burning of White Stem Borer affected Arabica plants. If stumps to be stored, immerse them in water for 10 days
- Spray Baylaton (160 gram/200 litres of water) or Contaf (400 ml / 200 litres of water) for susceptible varieties like S. 795 and Cauvery to control leaf rust
- Complete harvesting and processing of robusta, collection of left overs and fallen berries
- Complete pruning of Arabica and robusta
- Backing irrigation of robusta
- Collect and burn pupae of caterpillars in endemic areas, control measures against mealy bugs, if found
- Irrigation of young plants