GASES, LIQUIDS, AND SOLUTIONS

$$PV = nRT$$

$$\left(P + \frac{n^2a}{V^2}\right)(V - nb) = nRT$$

$$P_A = P_{total} \times X_A, \text{ where } X_A = \frac{\text{moles A}}{\text{total moles}}$$

$$P_{total} = P_A + P_B + P_C + \dots$$

$$n = \frac{m}{M}$$

$$K = {}^{\circ}C + 273$$

$$\frac{P_1V_1}{T_1} = \frac{P_2V_2}{T_2}$$

$$D = \frac{m}{V}$$

$$u_{rms} = \sqrt{\frac{3kT}{m}} = \sqrt{\frac{3RT}{M}}$$

$$KE \text{ per molecule} = \frac{1}{2}mv^2$$

$$KE \text{ per mole} = \frac{3}{2}RT$$

$$\frac{r_1}{r_2} = \sqrt{\frac{M_2}{M_1}}$$

$$\text{molarity, } M = \text{ moles solute per liter solution}$$

$$\text{molarity, } M = \text{ moles solute per kilogram solvent}$$

$$\Delta T_f = iK_f \times \text{ molality}$$

$$\Delta T_b = iK_b \times \text{ molality}$$

$$\pi = iMRT$$

$$A = abc$$

OXIDATION-REDUCTION; ELECTROCHEMISTRY

$$Q = \frac{[\mathbf{C}]^c [\mathbf{D}]^d}{[\mathbf{A}]^a [\mathbf{B}]^b}, \text{ where } a \mathbf{A} + b \mathbf{B} \to c \mathbf{C} + d \mathbf{D}$$

$$I = \frac{q}{t}$$

$$E_{\text{cell}} = E_{\text{cell}}^\circ - \frac{RT}{n\mathcal{F}} \ln Q = E_{\text{cell}}^\circ - \frac{0.0592}{n} \log Q @ 25^\circ \mathbf{C}$$

$$\log K = \frac{nE^\circ}{0.0592}$$

P = pressureV = volumeT = temperaturen = number of moles D = densitym = massv = velocity u_{rms} = root-mean-square speed KE = kinetic energyr =rate of effusion M = molar mass $\pi = \text{osmotic pressure}$ i = van't Hoff factor K_f = molal freezing-point depression constant K_b = molal boiling-point elevation constant A = absorbancea = molar absorptivityb = path lengthc = concentrationQ = reaction quotientI = current (amperes)q = charge (coulombs)

 E° = standard reduction potential K = equilibrium constant

t = time (seconds)

Gas constant,
$$R = 8.31 \text{ J mol}^{-1} \text{ K}^{-1}$$

$$= 0.0821 \text{ L atm mol}^{-1} \text{ K}^{-1}$$

$$= 8.31 \text{ volt coulomb mol}^{-1} \text{ K}^{-1}$$
Boltzmann's constant, $k = 1.38 \times 10^{-23} \text{ J K}^{-1}$

$$K_f \text{ for H}_2\text{O} = 1.86 \text{ K kg mol}^{-1}$$

$$K_b \text{ for H}_2\text{O} = 0.512 \text{ K kg mol}^{-1}$$

$$1 \text{ atm} = 760 \text{ mm Hg}$$

$$= 760 \text{ torr}$$
STP = 0.00°C and 1.0 atm
Faraday's constant, $\mathcal{F} = 96,500$ coulombs per mole of electrons