



2009 AP[®] CHEMISTRY FREE-RESPONSE QUESTIONS (Form B)



2. A student performed an experiment to investigate the decomposition of sodium thiosulfate, $\text{Na}_2\text{S}_2\text{O}_3$, in acidic solution, as represented by the equation above. In each trial the student mixed a different concentration of sodium thiosulfate with hydrochloric acid at constant temperature and determined the rate of disappearance of $\text{S}_2\text{O}_3^{2-}(\text{aq})$. Data from five trials are given below in the table on the left and are plotted in the graph on the right.

Trial	Initial Concentration of $\text{S}_2\text{O}_3^{2-}(\text{aq})$ (M)	Initial Rate of Disappearance of $\text{S}_2\text{O}_3^{2-}(\text{aq})$ (M s^{-1})
1	0.050	0.020
2	0.075	0.030
3	0.088	0.034
4	0.112	0.045
5	0.125	0.051



- Identify the independent variable in the experiment.
- Determine the order of the reaction with respect to $\text{S}_2\text{O}_3^{2-}$. Justify your answer by using the information above.
- Determine the value of the rate constant, k , for the reaction. Include units in your answer. Show how you arrived at your answer.
- In another trial the student mixed 0.10 M $\text{Na}_2\text{S}_2\text{O}_3$ with hydrochloric acid. Calculate the amount of time it would take for the concentration of $\text{S}_2\text{O}_3^{2-}$ to drop to 0.020 M .
- On the graph above, sketch the line that shows the results that would be expected if the student repeated the five trials at a temperature lower than that during the first set of trials.