Answer EITHER Question 2 below OR Question 3 printed on page 8. Only one of these two questions will be graded. If you start both questions, be sure to cross out the question you do not want graded. The Section II score weighting for the question you choose is 20 percent.

$$CO(g) + \frac{1}{2}O_2(g) \rightarrow CO_2(g)$$

2. The combustion of carbon monoxide is represented by the equation above.

(a) Determine the value of the standard enthalpy change, ΔH°_{rxn}, for the combustion of $CO(g)$ at 298 K using the following information.

$$C(s) + \frac{1}{2}O_2(g) \rightarrow CO(g) \qquad \Delta H^\circ_{298} = -110.5 \text{ kJ mol}^{-1}$$

$$C(s) + O_2(g) \rightarrow CO_2(g) \qquad \Delta H^\circ_{298} = -393.5 \text{ kJ mol}^{-1}$$

(b) Determine the value of the standard entropy change, ΔS°_{rxn}, for the combustion of $CO(g)$ at 298 K using the information in the following table.

Substance	S°_{298} (J mol^{-1} K^{-1})
$CO(g)$	197.7
$CO_2(g)$	213.7
$O_2(g)$	205.1

(c) Determine the standard free energy change, ΔG°_{rxn}, for the reaction at 298 K. Include units with your answer.

(d) Is the reaction spontaneous under standard conditions at 298 K ? Justify your answer.

(e) Calculate the value of the equilibrium constant, K_{eq}, for the reaction at 298 K.

GO ON TO THE NEXT PAGE.