### ADVANCED PLACEMENT CHEMISTRY EQUATIONS AND CONSTANTS

### ATOMIC STRUCTURE

$$E = hv c = \lambda v$$

$$\lambda = \frac{h}{mv} p = mv$$

$$E_n = \frac{-2.178 \times 10^{-18}}{n^2} \text{ joule}$$

## **EQUILIBRIUM**

$$K_{a} = \frac{[H^{+}] [A^{-}]}{[HA]}$$

$$K_{b} = \frac{[OH^{-}] [HB^{+}]}{[B]}$$

$$K_{w} = [OH^{-}] [H^{+}] = 1.0 \times 10^{-14} @ 25^{\circ}C$$

$$= K_{a} \times K_{b}$$

$$pH = -\log [H^{+}], pOH = -\log [OH^{-}]$$

$$14 = pH + pOH$$

$$pH = pK_{a} + \log \frac{[A^{-}]}{[HA]}$$

$$pOH = pK_{b} + \log \frac{[HB^{+}]}{[B]}$$

$$pK_{a} = -\log K_{a}, pK_{b} = -\log K_{b}$$

$$K_{p} = K_{c}(RT)^{\Delta n},$$

where  $\Delta n$  = moles product gas – moles reactant gas

### THERMOCHEMISTRY/KINETICS

$$\Delta S^{\circ} = \sum S^{\circ} \text{ products } -\sum S^{\circ} \text{ reactants}$$

$$\Delta H^{\circ} = \sum \Delta H_{f}^{\circ} \text{ products } -\sum \Delta H_{f}^{\circ} \text{ reactants}$$

$$\Delta G^{\circ} = \sum \Delta G_{f}^{\circ} \text{ products } -\sum \Delta G_{f}^{\circ} \text{ reactants}$$

$$\Delta G^{\circ} = \sum H^{\circ} - T\Delta S^{\circ}$$

$$= -RT \ln K = -2.303 RT \log K$$

$$= -n \mathcal{F} E^{\circ}$$

$$\Delta G = \Delta G^{\circ} + RT \ln Q = \Delta G^{\circ} + 2.303 RT \log Q$$

$$q = mc\Delta T$$

$$C_{p} = \frac{\Delta H}{\Delta T}$$

$$\ln[A]_{t} - \ln[A]_{0} = -kt$$

$$\frac{1}{[A]_{t}} - \frac{1}{[A]_{0}} = kt$$

$$\ln k = \frac{-E_{a}}{R} \left(\frac{1}{T}\right) + \ln A$$

$$E = \text{energy}$$
  $v = \text{velocity}$   
 $v = \text{frequency}$   $n = \text{principal quantum number}$   
 $\lambda = \text{wavelength}$   $m = \text{mass}$   
 $p = \text{momentum}$   
Speed of light,  $c = 3.0 \times 10^8 \,\text{m s}^{-1}$   
Planck's constant,  $h = 6.63 \times 10^{-34} \,\text{J s}$ 

Boltzmann's constant, 
$$k = 1.38 \times 10^{-23} \text{ J K}^{-1}$$
  
Avogadro's number  $= 6.022 \times 10^{23} \text{ mol}^{-1}$   
Electron charge,  $e = -1.602 \times 10^{-19}$  coulomb

1 electron volt per atom =  $96.5 \text{ kJ mol}^{-1}$ 

# **Equilibrium Constants**

 $K_a$  (weak acid)  $K_b$  (weak base)

 $K_w$  (water)

 $K_p$  (gas pressure)

 $K_c$  (molar concentrations)

 $S^{\circ}$  = standard entropy

 $H^{\circ}$  = standard enthalpy

 $G^{\circ}$  = standard free energy

 $E^{\circ}$  = standard reduction potential

T = temperature

n = moles

m = mass

q = heat

c =specific heat capacity

 $C_p$  = molar heat capacity at constant pressure

 $E_a$  = activation energy

k = rate constant

A =frequency factor

Faraday's constant,  $\mathcal{F} = 96,500$  coulombs per mole of electrons

Gas constant, 
$$R = 8.31 \text{ J mol}^{-1} \text{ K}^{-1}$$
  
= 0.0821 L atm mol<sup>-1</sup> K<sup>-1</sup>  
= 8.31 volt coulomb mol<sup>-1</sup> K<sup>-1</sup>