



2010 AP[®] CHEMISTRY FREE-RESPONSE QUESTIONS (Form B)



2. A galvanic cell and the balanced equation for the spontaneous cell reaction are shown above. The two reduction half-reactions for the overall reaction that occurs in the cell are shown in the table below.

Half-Reaction	E° (V) at 298 K
$\text{Fe}^{3+}(aq) + e^{-} \rightarrow \text{Fe}^{2+}(aq)$	+0.77
$\text{MnO}_4^{-}(aq) + 8 \text{H}^{+}(aq) + 5 e^{-} \rightarrow \text{Mn}^{2+}(aq) + 4 \text{H}_2\text{O}(l)$	+1.49

- On the diagram, clearly label the cathode.
- Calculate the value of the standard potential, E° , for the spontaneous cell reaction.
- How many moles of electrons are transferred when 1.0 mol of $\text{MnO}_4^{-}(aq)$ is consumed in the overall cell reaction?
- Calculate the value of the equilibrium constant, K_{eq} , for the cell reaction at 25°C. Explain what the magnitude of K_{eq} tells you about the extent of the reaction.

Three solutions, one containing $\text{Fe}^{2+}(aq)$, one containing $\text{MnO}_4^{-}(aq)$, and one containing $\text{H}^{+}(aq)$, are mixed in a beaker and allowed to react. The initial concentrations of the species in the mixture are 0.60 M $\text{Fe}^{2+}(aq)$, 0.10 M $\text{MnO}_4^{-}(aq)$, and 1.0 M $\text{H}^{+}(aq)$.

- When the reaction mixture has come to equilibrium, which species has the higher concentration, $\text{Mn}^{2+}(aq)$ or $\text{MnO}_4^{-}(aq)$? Explain.
- When the reaction mixture has come to equilibrium, what are the molar concentrations of $\text{Fe}^{2+}(aq)$ and $\text{Fe}^{3+}(aq)$?