# CLIMATEPULSE: TEMPORAL FUSION TRANSFORMERS AND SPATIO-TEMPORAL GRAPH NETWORKS IN CLIMATE MODELING

**A**

**Minor Project ([COURSE CODE]) Report**

Submitted in the partial fulfillment of the requirement for the degree of

**Bachelor of Technology**

**in**

**Computer and Communication Engineering**

By:

**ADITRI PAUL**

**2428010209**

Under the supervision of:

Dr. Hema Shekhawat (Assistant Professor)

**[SUBMISSION MONTH] 2026**

---

Department of Computer and Communication Engineering

School of Computing and Intelligent Systems

Manipal University Jaipur

VPO. Dehmi Kalan, Jaipur, Rajasthan, India – 303007

# STUDENT DECLARATION

This is to certify that the work entitled **“CLIMATEPULSE: TEMPORAL FUSION TRANSFORMERS AND SPATIO-TEMPORAL GRAPH NETWORKS IN CLIMATE MODELING”** submitted by **Aditri Paul (2428010209)** to Manipal University Jaipur for the award of the degree of Bachelor of Technology in Computer and Communication Engineering is a bonafide record of the work carried out by her under my supervision and guidance from July 2026 to [END MONTH] 2026.

**Dr. Hema Shekhawat** Internal Supervisor,

Department of Computer and Communication Engineering

Manipal University Jaipur

# PROJECT COMPLETION CERTIFICATE

**DEPARTMENT OF COMPUTER AND COMMUNICATION ENGINEERING**

**SCHOOL OF COMPUTING AND INTELLIGENT SYSTEMS**

**MANIPAL UNIVERSITY JAIPUR**

VPO. Dehmi Kalan, Jaipur, Rajasthan, India – 303007

This is to certify that **ADITRI PAUL** (Registration No. **2428010209**), a student of the Bachelor of Technology in Computer and Communication Engineering, has successfully completed the Project Based Learning [NUMBER] entitled: **“CLIMATEPULSE: TEMPORAL FUSION TRANSFORMERS AND SPATIO-TEMPORAL GRAPH NETWORKS IN CLIMATE MODELING.”**

This project was carried out during the [SEMESTER] semester in partial fulfillment of the requirements for the award of the B.Tech degree. The work has been found to be satisfactory and meets the academic standards of the institution.

**Dr. Hema Shekhawat**

Project Supervisor

Assistant Professor, Dept. of CCE

# ABSTRACT

This report details the conceptualization, design, development and evaluation of "ClimatePulse," an open-source Python pipeline that brings four climate-modelling tasks together in one reproducible workflow. Climate questions span very different time scales and data shapes: a monthly ocean index, a network of air-quality sensors, a global weather field on a sphere and a 75-year temperature record. They are usually studied with separate tools, separate data handling and separate evaluation rules, which makes results hard to compare or rebuild. ClimatePulse instead shares one data pipeline, one configuration system, one chronological split policy and one evaluation code base across four models.

The four components are: (1) a **Temporal Fusion Transformer (TFT)** that forecasts the Niño 3.4 sea-surface temperature anomaly 1 to 12 months ahead with 10–90 % uncertainty intervals, compared with a bidirectional LSTM; (2) an **A3TGCN spatio-temporal graph network** that predicts next-day PM2.5 at 75 OpenAQ monitoring stations; (3) an **encoder–processor–decoder graph network on a 2,000-node Fibonacci spherical mesh** that predicts global 2 m temperature and precipitation 6 hours ahead from ERA5 reanalysis, with an optional Niño 3.4 conditioning input; and (4) a **piecewise-linear trend with autocorrelated errors** that detects changes in the global warming rate from NASA GISTEMP data. All five data sources are public, and every model is scored against a persistence baseline on a strictly held-out test period.

On the test period, the TFT averages an RMSE of 0.782 °C over five seeds (LSTM 0.809 °C, persistence 1.036 °C); the weather model halves the persistence error for 6-hour 2 m temperature (1.21 K vs. 2.46 K); the pollution model is 10 % more accurate than persistence on concentration but shows no exceedance skill on only 85 observed test readings; and the warming-rate analysis finds a single regime change around 1976, from 0.03 to 0.21 °C per decade. An ablation shows that the Niño 3.4 conditioning of the weather model gives no measurable benefit. A Streamlit application lets users run the trained models on held-out data and see where they succeed and where they fail. Every figure and number in this report can be rebuilt from a fresh clone with one ordered sequence of `make` commands, on a CPU.

# TABLE OF CONTENTS

**ABSTRACT** **3**

**1. INTRODUCTION** **5**
1.1 Problem Statement . . . . . 5
1.2 Proposed Solution . . . . . 5

**2. SYSTEM ARCHITECTURE AND DESIGN** **7**
2.1 System Architecture . . . . . 7
2.1.1 Component Breakdown . . . . . 7
2.2 Component Deep Dive . . . . . 8
2.2.1 Model Choice and Alternatives . . . . . 9
2.2.2 Baselines and Skill Score . . . . . 10
2.3 Functionality to Module Mapping . . . . . 10
2.4 Data Flow . . . . . 11

**3. IMPLEMENTATION DETAILS** **12**
3.1 Interactive Simulation App (Streamlit) . . . . . 12
3.2 Training and Evaluation Orchestration . . . . . 13
3.3 Robustness & Error Handling . . . . . 14
3.4 How to Run: Rebuilding Every Result . . . . . 16

**4. DATA FLOW** **18**
4.1 Data Flow for Training and Evaluation . . . . . 18
4.2 Data Sources and Pipeline Stages . . . . . 18
4.3 Data Flow for the Simulation App . . . . . 19

**5. OUTCOMES AND FUTURE WORK** **20**
5.1 Project Outcomes and Conclusion . . . . . 20
5.2 Detailed Evaluation Results . . . . . 21
5.2.1 Evaluation Protocol . . . . . 21
5.2.2 El Niño: Five Seeds . . . . . 21
5.2.3 El Niño: Lead Time . . . . . 21
5.2.4 El Niño: Bias and Calibration . . . . . 22
5.2.5 Air Pollution: Next-Day PM2.5 . . . . . 23
5.2.6 Weather Model . . . . . 25
5.2.7 Warming Rate . . . . . 26
5.2.8 Training Logs . . . . . 27
5.2.9 Patent Application Status . . . . . 28
5.2.10 Conclusions by Model . . . . . 28
5.3 Future Work . . . . . 29

**6. BIBLIOGRAPHY** **30**

# 1. INTRODUCTION

## 1.1 Problem Statement

Climate phenomena operate across very different scales. The El Niño–Southern Oscillation (ENSO) evolves over months and shifts rainfall and temperature worldwide; urban air pollution changes from day to day and is shaped by local sources and weather; the global atmosphere changes every few hours; and the warming trend unfolds over decades. Deep learning has produced strong single-task models for several of these problems, but students and researchers who want to study them together face a fragmented landscape.

The key challenges in this setting include:

- **Siloed tools and data handling:** Each task normally comes with its own download scripts, file formats, preprocessing and evaluation code. Comparing models across tasks, or reusing a forecast from one model as input to another, requires rebuilding much of this machinery by hand.
- **Expensive physical models:** Numerical weather prediction and Earth system models are physically grounded but need supercomputers, which puts them out of reach for student projects and rapid experimentation.
- **Point forecasts without uncertainty:** Many data-driven forecasts return a single number. Decision makers also need to know how confident the forecast is, which calls for calibrated prediction intervals.
- **Data that does not fit a regular grid:** Air-quality sensors are irregular points on a map, and a latitude–longitude grid crowds its points near the poles. Standard convolutional networks assume a regular grid and handle both cases poorly.
- **Optimistic evaluation:** Time-series results are easily inflated by leakage, where information from the test period reaches training through overlapping windows or normalisation statistics. An earlier version of this project suffered exactly this problem (TFT skill 0.91), and a fair evaluation lowered it substantially.
- **Poor reproducibility:** Published numbers are often typed in by hand and cannot be traced back to the run that produced them.

This project addresses these issues by building a single, open-source, CPU-friendly pipeline that downloads public data, trains four complementary models, evaluates them honestly against simple baselines and regenerates every reported figure and table automatically.

## 1.2 Proposed Solution

We propose “ClimatePulse,” a modular climate-modelling framework designed around one shared data pipeline and one evaluation discipline. Each of its four modules answers a different climate question with an architecture that matches the shape of its data:

- **El Niño forecasting:** How warm will the tropical Pacific be in 1 to 12 months? A Temporal Fusion Transformer reads 60 months of history and outputs the 10th, 50th and 90th percentile forecast for each of the next 12 months.
- **Air-pollution forecasting:** How much PM2.5 will each monitoring station record tomorrow, and will it exceed the AQI-100 threshold? An A3TGCN graph network treats the stations as nodes of a graph.
- **Neural weather modelling:** What will global temperature and precipitation be in 6 hours? A graph network maps the ERA5 grid onto an evenly spaced spherical mesh, processes it there and maps it back.
- **Warming-rate analysis:** When did the speed of global warming change since 1950? A segmented linear trend with autocorrelated errors is fitted, and the number of breaks is chosen by the Bayesian Information Criterion (BIC).

The user interacts with the system in two ways:

- **Pipeline commands:** A short, ordered sequence of `make` targets downloads the data, trains every model, runs the robustness studies, generates the evaluation artifacts and compiles the documents. Configuration lives in `configs/runtime_conf.yaml` and can be overridden with environment variables.
- **Simulation app:** `make app` opens a browser application that runs the already-trained models on held-out data, so results can be explored without retraining.

Instead of a loose collection of scripts, the outputs form one traceable package:

- **Forecasts with uncertainty:** Quantile intervals for ENSO, per-station PM2.5 predictions and global weather fields.
- **Baseline-relative metrics:** Every model is scored against persistence (carrying the latest value forward), with skill $= 1 - \mathrm{MSE}_{\mathrm{model}}/\mathrm{MSE}_{\mathrm{persistence}}$.
- **Robustness studies:** A five-seed study for the El Niño models and a one-change-at-a-time ablation of the weather model.
- **Versioned snapshots:** Each run's checkpoints, metrics and figures are saved under `runtime/saved/run_<timest` and the research paper reads its figures directly from the snapshot it reports.

# 2. SYSTEM ARCHITECTURE AND DESIGN

This chapter details the technical design of ClimatePulse. It outlines the system architecture, the responsibilities of each component, the mapping from functionality to code modules, and the path data takes from download to published result.

## 2.1 System Architecture

The system follows a layered pipeline architecture with five layers: open data sources, a data pipeline that fetches, cleans and caches them, a model layer with one package per task, an evaluation layer that scores every model and writes figures and tables, and an output layer made up of the research paper, the patent application and the interactive app. Each layer communicates with the next only through files on disk (Parquet tables, checkpoints, JSON and CSV results), so any stage can be rerun or inspected on its own.





Figure 1: Layered architecture of ClimatePulse. Each arrow is a set of files on disk, produced by the `make` target shown on the right.

### 2.1.1 Component Breakdown

**Technology Stack:**

- **Programming Language:** Python 3.12 (pinned in `.python-version`, `pyproject.toml` and the `Makefile`)
- **Deep Learning:** PyTorch 2.12, PyTorch Geometric 2.8, PyTorch Geometric Temporal (A3TGCN), pytorch-forecasting 1.8 (TFT)

- **Statistical Modelling:** NumPy and SciPy (change-point search), PyMC and ArviZ (optional emissions regression)
- **Data Engineering:** pandas, xarray, zarr and gcsfs (ERA5 from Google Cloud), pyarrow (Parquet storage)
- **Configuration:** Pydantic-Settings v2 with YAML files and environment-variable overrides
- **Interactive App:** Streamlit 1.66 and Plotly
- **Hardware:** Runs on CPU; CUDA, AMD ROCm and Intel NPU (OpenVINO) installs are also supported
- **Source Code Management and Testing:** Git and GitHub; pytest unit tests (`make test`)

**Data Pipeline (`src/data/`):**

- **Responsibility:** Fetches each source through its own connector in `src/data/sources/`, reusing cached downloads where present, cleans it with a source-specific function in `processing/cleaning.py` and derives features in `processing/feature_engineering.py`. It also builds the pollution station graph (`processing/graph_builder.py`). `DataPipeline` is a stateless orchestrator; storage is handled by `storage/db.py`, which writes Parquet tables and registers every source in a JSON metadata file.

**Model Layer (`src/models/`):**

- **Responsibility:** One package per task, each with a dataset module (windowing, splits and training-only normalisation), a model module, a trainer and an evaluation module. Training is started through a single command-line entry point, `src/scripts/train_models.py`, which the `make train-*` targets call.

**Evaluation and Artifacts (`src/scripts/generate_eval_artifacts.py`):**

- **Responsibility:** Loads the trained checkpoints, scores every model on its test split against the baselines, and writes figures, CSV and JSON results and a `metrics_summary.md` to `runtime/eval_artifacts/`. `make save-run` then copies checkpoints and artifacts into a timestamped snapshot.

**Interactive App (`src/app/`):**

- **Responsibility:** Runs the trained models on held-out data in the browser. It never trains or modifies a model, and it labels each page with how far its results are validated.

## 2.2 Component Deep Dive

**Model Roster:**

The core of the project is a set of four models. Each is chosen to fit the structure of its data, and each is tested against a simpler baseline rather than assumed to be better.

1. **Temporal Fusion Transformer (El Niño):**
   - *Task:* Forecasts the monthly Niño 3.4 anomaly for each of the next 12 months from a 60-month history. Inputs are the Niño 1+2, 3, 3.4 and 4 indices, their 3-, 6- and 12-month rolling mean and standard deviation, and the calendar month (as sine and cosine) as a known future input. Gated variable-selection networks weight the inputs, an LSTM encoder–decoder captures local patterns and four-head self-attention looks back over the whole window. Three quantile heads ($\tau$ = 0.1, 0.5, 0.9) are trained with the pinball loss $\rho_\tau(u) = u\,(\tau - \mathbf{1}[u < 0])$. The model has 96,047 parameters.
   - *Baselines:* A two-layer bidirectional LSTM (142,668 parameters) and persistence.
2. **A3TGCN Graph Network (Air Pollution):**
   - *Task:* Treats each day's PM2.5 readings as a signal on a 75-node station graph built from the 10 nearest stations by great-circle distance (705 edges, weight $1/(d+1)$ with $d$ in km, edges

beyond 500 km dropped). Graph convolution shares information between nearby stations, a GRU tracks the state over 7 snapshots and soft attention weighs those snapshots before a small MLP predicts next-day PM2.5. The model has only 6,984 parameters. Missing station-days are masked out of the loss.
    - *Baselines:* Persistence for concentration, and an "always alert" rule for exceedance days (PM2.5 $\geq$ 35.5 µg/m³, i.e. AQI above 100).
3. **Fibonacci-Mesh Graph Network (Weather):**
    - *Task:* Predicts the next 6-hourly global 2 m temperature and precipitation fields from the previous two on the 5.625° ERA5 grid (2,048 points). An encoder maps the grid onto a 2,000-node Fibonacci spherical mesh, which covers the sphere evenly at any chosen size, unlike latitude–longitude grids that crowd at the poles or icosahedral meshes limited to fixed sizes. Four residual GraphSAGE layers pass messages between mesh neighbours, and a decoder maps the result back to the grid. An optional Niño 3.4 input can be added to every processor layer. Training uses a cosine-latitude-weighted MSE loss. The model has 217,346 parameters.
    - *Baselines:* Persistence, plus an ablation of the Niño 3.4 input and of the loss weighting.
4. **Segmented Trend with AR(1) Errors (Warming Rate):**
    - *Task:* Fits a continuous piecewise-linear trend to 76 annual GISTEMP global means (1950–2025), so that a change-point alters the warming *rate* rather than the temperature level. Errors follow an AR(1) process, fitted by Prais–Winsten generalised least squares, so that year-to-year correlation does not invent breaks. Every placement of 0 to 3 change-points with segments of at least 10 years is fitted, and BIC chooses how many the data justify. No neural network is involved.
    - *Comparison:* A mean-shift CUSUM test, which finds 12 spurious breaks because it tests the level rather than the slope.

### 2.2.1 Model Choice and Alternatives

Each architecture was chosen because it fits the shape of its data, and each choice was set against established approaches from the literature.

| Task | Chosen | Alternatives considered | Why the choice |
|---|---|---|---|
| El Niño | Temporal Fusion Transformer | CNN on climate-model output (Ham et al.); plain LSTM | Forecasts all 12 months at once, uses known future inputs such as the calendar month, gives calibrated quantile intervals rather than only point forecasts, and its attention can be inspected |
| Air pollution | A3TGCN | Grid CNNs; diffusion GCN (DCRNN) | Stations are irregular points, not a grid: graph convolution shares signal between nearby stations, a GRU tracks time and attention weighs recent days, in only 6,984 parameters |

| Task | Chosen | Alternatives considered | Why the choice |
|---|---|---|---|
| Weather | Mesh GNN on a Fibonacci mesh | FourCastNet (regular grid); GraphCast (icosahedral mesh) | Regular grids crowd at the poles; a Fibonacci mesh is uniform at any size, while icosahedral meshes exist only at fixed subdivision levels |
| Warming rate | Segmented trend, AR(1) errors, BIC | Mean-shift CUSUM; breaks with independent errors | Tests the rate, not the level; AR(1) errors stop autocorrelation inventing breaks (CUSUM finds 12 spurious ones) |

#### 2.2.2 Baselines and Skill Score

A model adds value only where it beats a simple reference. Climate variables change slowly, so at short leads this is a demanding test. Every model is therefore compared with **persistence**, which assumes nothing changes and needs no model at all:

- **El Niño:** each of the next 12 months equals this month's Niño 3.4 value.
- **Air pollution:** tomorrow's PM2.5 equals today's PM2.5 at the same station. For exceedance days, an **always-alert** rule (predict unhealthy air every day) is also used.
- **Weather:** the field in 6 hours equals the current field.

Skill is measured as $\text{Skill} = 1 - \text{MSE}_{\text{model}}/\text{MSE}_{\text{persistence}}$ (the convention of Barnston et al.). A value of 0 means no better than persistence, 1 would be perfect and a negative value is worse than persistence.

### 2.3 Functionality to Module Mapping

To keep the code base clean, each concern lives in its own module under `src/` (imported with `PYTHONPATH=src`):

- **Configuration:** `config.py` defines Pydantic-Settings models for data, model, NWM and logging options. `configs/runtime_conf.yaml` and `configs/dataset_conf.yaml` hold the values, and every path is derived from `PROJECT_ROOT`.
- **Data Ingestion:** `data/sources/` holds one connector per source (`noaa_ersst.py`, `openaq.py`, `nasa_gistemp.py`, `climate_trace.py`); ERA5 is fetched by `scripts/download_nwm_data.py`.
- **Processing and Storage:** `data/processing/` (cleaning, feature engineering, station graph) and `data/storage/db.py` (Parquet and JSON metadata), orchestrated by `data/pipeline.py`.
- **Models:** `models/el_nino/` (TFT and LSTM), `models/pollution/` (A3TGCN), `models/nwm/` (mesh builder, model, dataset, trainer) and `models/climate_feedback/` (change-point model and optional Bayesian regression).
- **Robustness Studies:** `scripts/run_elnino_seeds.py` (five seeds, TFT with and without the calendar year), `scripts/export_tft_forecasts.py` and `scripts/run_nwm_ablation.py`.
- **Evaluation and Figures:** `scripts/generate_eval_artifacts.py` with chart helpers in `visualization/`.
- **Simulation App:** `app/main.py` (entry point and tabs), `app/loaders.py` (all model and data access, with no Streamlit import) and `app/views/` (one module per page).
- **Tests:** `tests/` covers the El Niño split boundaries, the pollution graph and masking, the NWM dataset, the change-point model, configuration and the app loaders.

## 2.4 Data Flow

The life cycle of a complete ClimatePulse run follows a strict, ordered pipeline:

1. **Data Download:** `make download-data` runs `DataPipeline` for NOAA ERSST, OpenAQ, NASA GISTEMP and Climate TRACE; `make download-nwm-data` fetches the ERA5 subset. Cached files are reused.
2. **Model Selection:** `make train-elnino-seeds` trains the TFT (with and without the calendar-year input) and the LSTM over five seeds, and selects the TFT variant by mean *validation* loss before any test result is looked at.
3. **Training:** `make train-elnino`, `make train-pollution` and `make train-nwm` train the main models; the best checkpoint by validation loss is kept in every case.
4. **Robustness Studies:** `make export-tft-forecasts` writes the TFT's out-of-sample lead-1 forecasts, and `make nwm-ablation` trains and scores the three weather-model variants.
5. **Evaluation and Publication:** `make gen-eval-artifacts` scores every model and writes the figures and tables; `make save-run` snapshots the run; `make sync-paper-artifacts` points the paper at that snapshot; and `make gen-research-paper-pdf` compiles the paper.

# 3. IMPLEMENTATION DETAILS

## 3.1 Interactive Simulation App (Streamlit)

The front end is a Streamlit application, launched with `make app`, that runs the trained models on held-out data. It has five tabs: an overview and one tab per simulation. Only the open tab runs, so one page's models are not loaded or re-run for the others.

- **Overview:** Shows the headline results and which checkpoints were found, and whether each came from the live training output or a committed snapshot.
- **Weather:** The user picks any moment in the test period and compares ERA5 truth, the model's prediction and the error, as maps or on a rotatable 3D globe, for temperature or precipitation. The user can switch between the main model and the three ablation variants, override the Niño 3.4 input, and roll the model forward and animate it.
- **El Niño:** A slider selects any 12-month forecast from 2015 to 2026 and shows the TFT median and 10–90 % band against the LSTM, persistence and the observed values, with an animation through the test period and error by lead time.
- **Air pollution:** Predicted and observed PM2.5 are shown on a map, coloured by US EPA air-quality band, and a station can be highlighted to see its recent inputs, prediction and reading.
- **Warming rate:** The user changes the maximum number of breaks and the shortest segment length and watches the fitted trend, the BIC choice and the segment slopes update, with a straight-line fit for comparison.

**Honest badges:** Every page states how far its results are validated: "Validated" for El Niño, "Validated for one step" for the weather model, "Demonstration" for pollution and "Statistical model" for the warming rate. Rolling the weather model forward feeds it its own output, which it was never trained or tested on, so the page warns that errors grow and may exceed persistence.





Figure 2: Overview tab of the simulation app, showing the headline results and the honest reading of each model.





Figure 3: Weather tab: ERA5 truth, model prediction and error for a chosen test time.

### 3.2 Training and Evaluation Orchestration

The pipeline is driven by a `Makefile` and a single training entry point, so every step can be run on its own or as part of the full sequence.

- **Configuration:** Two layers of configuration keep secrets apart from parameters. Credentials (the OpenAQ API key) live in an untracked `.env` file; operational parameters (epochs, seeds, encoder lengths, NWM conditioning source and loss weighting) live in `configs/runtime_conf.yaml`. Any value can be overridden with an environment variable such as `MODEL_MAX_EPOCHS=5`, which makes quick tests and the Colab notebook (`run.ipynb`) possible without editing files.
- **Training setup:** All models use PyTorch, and the learning rate halves when validation loss stops improving (`ReduceLROnPlateau`, factor 0.5). Batch size is 32 for the TFT, LSTM and weather model; the A3TGCN trains on one graph window per step. The best checkpoint by validation loss is kept in every case.

| Model | Parameters | Loss | Optimiser | Stopping |
|---|---|---|---|---|
| TFT | 96,047 | Quantile (pinball) loss at $\tau$ = 0.1, 0.5, 0.9 | Adam, lr $10^{-3}$, clip 0.1 | Early stop, patience 10 |
| BiLSTM | 142,668 | MSE over 12 leads | Adam, lr $10^{-3}$, clip 1.0 | Early stop, patience 15 |
| A3TGCN | 6,984 | MSE on observed station-days only | Adam, lr $10^{-3}$, clip 1.0 | 100 epochs, best kept |
| Weather model | 217,346 | MSE weighted by cos(latitude) | AdamW, lr $10^{-3}$, decay $10^{-4}$ | Early stop, patience 10 |

- **Efficient graph batching:** In the weather model, the edge indices for a whole batch are built by offset arithmetic on tensors rather than Python loops, so the per-batch overhead does not grow with batch size.
- **Compute budget:** The networks are small (7k to 217k parameters) and train on a CPU. The main models take about 2.2 hours, almost all of it the weather model (about 2 hours, roughly 90 s per epoch); the robustness studies add about 6 hours, mostly the three ablation variants trained in parallel.
- **Snapshots:** `make save-run` copies checkpoints and artifacts to `runtime/saved/run_<timestamp>/`. The research paper's `artifacts/` folder is a symbolic link to the selected snapshot, so the paper always reads one fixed run. The results in this report come from `run_20261027_230426`, which is committed together with its checkpoints.





Figure 4: Training (blue) and validation (red, dashed) loss for the four deep-learning models.

### 3.3 Robustness & Error Handling

Forecasting projects rarely fail loudly; more often, results look better than they are. ClimatePulse therefore treats evaluation integrity as its main form of error handling. Six rules apply to every model:

- **Chronological splits:** Data are split 70/15/15 into training, validation and test periods in time order. For El Niño, training runs from January 1950 to August 2003, validation from September 2003 to February 2015, and testing from March 2015 to September 2026. No training target falls in the validation or test period.
- **Training-only statistics:** Normalisation statistics are computed on the training split only and then applied to validation and test data.
- **Persistence baselines:** Every skill score is measured against carrying the latest observed value forward, the minimum any forecast must beat.
- **Multiple seeds and validation-based selection:** El Niño results are reported as mean $\pm$ standard deviation over five seeds, and the TFT variant is chosen on validation loss, never on test results.
- **Physical units and masking:** Pollution is scored in µg/m³ at the AQI-100 threshold, and missing readings are excluded rather than imputed into the score.

- **Ablation:** The weather model's Niño 3.4 input and loss weighting are each tested one change at a time.

Further safeguards in the code:

- **Fault-isolated downloads:** Each source in `DataPipeline.run_all()` runs in its own error handler, so one failing API does not stop the others, and the Climate TRACE connector retries failed requests. Cached downloads are reused so that repeated runs do not hit the APIs again.
- **Helpful failures in the app:** If a checkpoint or data file is missing, the app raises a `DataMissing` error that names the `make` command that creates it, and the page shows that message instead of crashing. Checkpoints are found by searching the live training directory first and then each committed snapshot, newest first, so a fresh clone can run the app without retraining.
- **Unit tests:** The test suite (`make test`) checks, among other things, that El Niño validation and test targets never overlap training targets, that pollution windows and masks are built correctly, and that the change-point model recovers known breaks.




Figure 5: El Niño tab: a 12-month TFT forecast with its 10–90 % band against the LSTM, persistence and observations.





Figure 6: Warming-rate tab: adjustable change-point search with the BIC choice and segment slopes.

### 3.4 How to Run: Rebuilding Every Result

One ordered run rebuilds every figure and table. The same steps are listed in `run.sh`, and `run.ipynb` runs them on Google Colab. The committed snapshot `run_20261027_230426` lets results be checked without retraining.

```
# 0. Get the code
git clone <repository-url> && cd climate_pulse

# 1. Environment and credentials
make install-dev
cp -n .env.example .env              # set DATA_OPENAQ_API_KEY

# 2. Data
make download-data
make download-nwm-data
make test

# 3. El Niño: five seeds, then main run
make train-elnino-seeds
# set tft_use_year in configs/runtime_conf.yaml from elnino_seed_selection.json
make train-elnino

# 4. Other models
make train-pollution
make train-nwm                       # about 2 h on CPU
make export-tft-forecasts            # needs the main TFT
make nwm-ablation                    # about 5 h on CPU
```

```
# 5. Artifacts, snapshot, documents
make gen-eval-artifacts
make save-run
make sync-paper-artifacts
make gen-research-paper-pdf
```

The order matters: the ablation scores the main weather model, so `train-nwm` comes first, and exporting TFT forecasts needs the main TFT.

# 4. DATA FLOW

## 4.1 Data Flow for Training and Evaluation

The primary execution cycle turns five public datasets into trained models and published metrics:

1. **Ingestion:** NOAA ERSST v5 provides monthly sea-surface temperature from 1950, from which the Niño 1+2, 3, 3.4 and 4 indices are computed. OpenAQ v3 provides PM2.5 readings for stations in India and the United States (an API key is required). NASA GISTEMP v4 provides global temperature anomalies. Climate TRACE v7 provides country-level emissions for an optional regression that no reported result uses. ERA5 from WeatherBench2 provides 6-hourly 2 m temperature and precipitation at 5.625° for 2020–2021 (2,924 time steps).
2. **Processing:** Each source is cleaned by its own function and turned into model-ready features: rolling and seasonal features for the Niño indices (`nino_features.parquet`), daily station means and a 10-nearest-neighbour station graph for OpenAQ, annual means for GISTEMP, and a regridded, subset NetCDF file for ERA5.
3. **Windowing and Splitting:** Inside each model's dataset module, the series is cut into input–target windows and split chronologically. Validation and test windows take their input history from the preceding period, but their targets never overlap earlier splits. Normalisation statistics are fitted here, on training data only.
4. **Training:** Each model is trained on its training windows, monitored on its validation windows, and its best checkpoint by validation loss is saved.
5. **Evaluation:** Checkpoints are scored on the test windows against persistence (and the LSTM or always-alert baselines), predictions are converted back to physical units, and the metrics, figures and tables are written to `runtime/eval_artifacts/` and then frozen in a snapshot.

## 4.2 Data Sources and Pipeline Stages

All five datasets are open access and are downloaded automatically by `make download-data` (four sources) and `make download-nwm-data` (ERA5). Only OpenAQ needs a free API key.

| Dataset | What it provides | Used by | URL |
|---|---|---|---|
| NOAA ERSST v5 | Sea temperature 1950–2026, monthly on a 2° grid | El Niño | psl.noaa.gov/data/gridded/ |
| OpenAQ v3 | PM2.5, 75 stations, 2016–2025 | Pollution | openaq.org |
| NASA GISTEMP v4 | Global temperature 1950–2025 | Warming rate | data.giss.nasa.gov/gistemp/ |
| Climate TRACE v7 | Country emissions (optional regression) | Not used in any reported result | climatetrace.org |
| ERA5 / WeatherBench2 | 2 m temperature and rain, 2020–2021, 6-hourly at 5.625° (2,924 time steps) | Weather | weatherbench2.readthedocs.io |

Each model reads its own cached file, so any stage can be rerun or inspected on its own. The splits and training-only normalisation happen later, inside each model's dataset module.

| Source | Processing | Stored in `datasets/` | Used by |
|---|---|---|---|
| NOAA ERSST v5 | Clean → Niño indices → rolling and seasonal features | `nino_features.parquet` | TFT, LSTM; weather-model Niño input |
| OpenAQ v3 | Clean, add AQI; stations → 10-nearest-neighbour haversine graph | PM2.5 cache + `openaq/locations.parquet` | A3TGCN |
| NASA GISTEMP v4 | Clean → rolling and seasonal features | `gistemp_features.parquet` | Change-point model |
| Climate TRACE v7 | Clean → align with GISTEMP by year | `regression_dataset.parquet` | Bayesian regression (optional) |
| ERA5 / WeatherBench2 | Subset 2020–2021, 5.625°, temperature and rain | `era5_5deg_6hr_2020_2022.nc` | Weather model |

### 4.3 Data Flow for the Simulation App

The app reads the same checkpoints and evaluation code as the paper, so its numbers match the reported ones:

1. **Tab selection:** When the user opens a tab, Streamlit reruns the script, and only the open tab's page renders.
2. **Resolution:** `app/loaders.py` finds the required checkpoint (live directory first, then committed snapshots) and the required data file, raising `DataMissing` with the fix if either is absent.
3. **Cached loading:** Models and datasets are loaded once and cached, so moving a slider re-runs inference without reloading weights.
4. **Inference:** The page runs the model on the chosen held-out sample. Weather predictions are computed one sample at a time, which gives 1.217 K against the reported 1.214 K computed in batches of 16, because the model's layer normalisation pools statistics across a batch.
5. **Rendering:** Plotly draws the maps, globe, time series and metric cards, using the repository's white, grey and blue theme.

# 5. OUTCOMES AND FUTURE WORK

## 5.1 Project Outcomes and Conclusion

The ClimatePulse project designed, built and evaluated a reproducible pipeline that integrates four climate-modelling tasks. The headline results on the held-out test periods are:

| Model | Result | Baseline | Reading |
|---|---|---|---|
| El Niño TFT (5 seeds) | RMSE 0.782 ± 0.075 °C, skill 0.43 | LSTM 0.809 °C; persistence 1.036 °C | Beats persistence from 3 months ahead; comparable to the LSTM |
| Air-pollution A3TGCN | RMSE 46.6 µg/m³ | Persistence 52.0 µg/m³ | 10 % better on concentration; exceedance F1 0.828 equals always-alert |
| Weather mesh GNN | T2m RMSE 1.21 K (6 h) | Persistence 2.46 K | Halves persistence error; Niño 3.4 input adds nothing |
| Warming-rate change-points | One break in 1976 | Straight line; CUSUM | 0.035 → 0.211 °C per decade |

- **Functionality:** One pipeline downloads, trains, evaluates and documents all four components. Each module shares the data infrastructure, configuration and evaluation tooling, and none is a standalone script.
- **Forecast quality:** Both learned El Niño models clearly beat persistence. The TFT is the most accurate model at every lead from 3 to 12 months (12-month RMSE 0.856 °C against 0.934 °C for the LSTM and 1.376 °C for persistence in the main run), but across seeds its edge over the LSTM is smaller than its own run-to-run spread. The weather model learns real 6-hour dynamics: its error is half that of persistence, lowest in the tropics (0.84 K) and highest near the poles (1.55 K), and its precipitation error is 24 % below persistence.
- **Scientific honesty:** The evaluation exposes weaknesses as well as strengths. Both El Niño models are biased cold by about 0.3 °C because the test period (mean Niño 3.4 anomaly +0.14 °C) is warmer than the training period (−0.23 °C). The pollution test set has only 85 observed station-days, too few to show exceedance skill. The ablation shows that conditioning the 6-hour weather model on Niño 3.4 gives no benefit, and that feeding it the TFT's forecasts makes it worse (1.335 K), so the proposed cross-model coupling is not supported on two years of data.
- **Transparency:** The simulation app lets anyone explore the results interactively, and its badges and warnings make the limits of each model visible, including the sparsity of the pollution data, which headline numbers hide.
- **Reproducibility and cost:** All data are public, the models train on a CPU, and the full study can be rebuilt from a fresh clone in about 7–8 hours on CPU. The committed snapshot lets results be checked without retraining.

The work also contributed to a draft research paper and a draft patent application covering the Fibonacci-mesh weather model and its loop-free graph batching method.

Ultimately, ClimatePulse shows that a small, open, CPU-trainable pipeline can study ENSO, air quality, global weather and the warming trend under one consistent evaluation, and that holding every model to the same leakage-free standard produces results that are more modest than the architectures' reputations, but trustworthy.

## 5.2 Detailed Evaluation Results

### 5.2.1 Evaluation Protocol

Leakage, meaning test-period information reaching training, inflated an earlier version of this project (TFT skill 0.91). The six rules in Section 3.3 remove it, so the numbers below are lower than the leaky run's, but each one is a fair test on unseen data. All results come from the saved run `20261027_230426`, the same snapshot that the research paper and the patent application report.

### 5.2.2 El Niño: Five Seeds

Test period March 2015 – September 2026, leads 1–12, mean ± standard deviation over five seeds. RMSE is the average error (lower is better), ACC the correlation with observed anomalies, and bias is forecast minus observed.

| Model | RMSE (°C) | ACC | Skill | Bias (°C) |
|---|---|---|---|---|
| TFT, no year (selected) | 0.782 ± 0.075 | 0.645 ± 0.023 | 0.427 ± 0.113 | −0.321 ± 0.146 |
| TFT, with year | 0.838 ± 0.070 | 0.483 ± 0.193 | 0.341 ± 0.106 | −0.242 ± 0.141 |
| Bidirectional LSTM | 0.809 ± 0.020 | 0.562 ± 0.035 | 0.390 ± 0.029 | −0.268 ± 0.078 |
| Persistence | 1.036 | 0.383 | 0.000 | +0.062 |

Removing the calendar-year input lowered the mean validation loss from 0.331 to 0.283, so that TFT variant was selected before any test result was examined. Both learned models clearly beat persistence. The TFT's edge over the LSTM is smaller than its seed-to-seed spread (its RMSE spread is nearly four times the LSTM's), so the two should be treated as comparable. The seed-42 main run scored better than this average (RMSE 0.684 °C).





Figure 7: A 12-month forecast for one held-out sequence: TFT median and 10–90 % band, LSTM and observed Niño 3.4.

### 5.2.3 El Niño: Lead Time

RMSE by forecast lead in the main run (seed 42):

| Lead (months) | TFT | LSTM | Persistence |
|---|---|---|---|
| 1 | 0.394 | 0.324 | **0.247** |
| 3 | **0.525** | 0.542 | 0.609 |
| 6 | **0.704** | 0.862 | 1.032 |
| 9 | **0.765** | 0.927 | 1.269 |
| 12 | **0.856** | 0.934 | 1.376 |

Persistence is best at 1 month because Niño 3.4 changes slowly from month to month. The LSTM overtakes it from month 2, and the TFT is the most accurate of the three at every lead from 3 to 12 months. The learned models earn their keep from about three months out.





Figure 8: RMSE as a function of forecast lead time for the TFT, LSTM and persistence.

### 5.2.4 El Niño: Bias and Calibration

Both models run cold in a warmer test period. The mean TFT bias over five seeds is −0.32 °C (LSTM −0.27 °C), and the mean Niño 3.4 anomaly rises from −0.23 °C in the training period to +0.14 °C in the test period. The LSTM never sees the calendar year and shares the bias, so the year input is not the cause; non-stationarity is the likelier one.

An 80 % interval should contain 80 % of observations. In the main run the TFT's interval contains 75.7 %, with 16.1 % above its top and 8.2 % below its bottom; the five-seed mean coverage is about 63 %. Recent years are warmer than the training years, so fixing the bias matters more than widening the interval.





Figure 9: TFT quantile calibration: observed frequency of outcomes below the predicted 10th, 50th and 90th percentiles. Points below the diagonal indicate forecasts that are too low.





Figure 10: Six-month-ahead TFT median and 10–90 % interval against observed values across the test set. The forecast reproduces the timing of the 2015–16 and 2023–24 El Niño events but underestimates their peaks.

### 5.2.5 Air Pollution: Next-Day PM2.5

Forecasts are scored only where a reading exists. An exceedance is a day with PM2.5 ≥ 35.5 µg/m³ (AQI above 100).

| Metric | A3TGCN | Persistence | Always alert |
|---|---|---|---|
| RMSE (µg/m³) | 46.6 | 52.0 | — |
| Exceedance F1 | 0.828 | 0.881 | 0.828 |
| Precision / recall | 0.706 / 1.000 | — | — |

| Metric | A3TGCN | Persistence | Always alert |
|---|---|---|---|

Only 85 of the 1,500 test station-days (from 36 stations) have a reading, and 62 of those fall on two dates. Of all 85 readings, 70.6 % exceed the threshold, which makes always-alert a strong baseline. The model flags every observed day (60 true positives, 25 false positives), so its F1 equals an always-alert rule. Concentration forecasts beat persistence by about 10 %, but 85 test readings are too few to show that the model can spot unhealthy-air days.





Figure 11: Exceedance confusion matrix on observed test station-days: 60 true positives, 25 false positives, no false negatives and no true negatives.





Figure 12: Distribution of per-station RMSE (µg/m³) across the 36 stations with test readings.

### 5.2.6 Weather Model

On the 438 held-out test samples, the main model predicts 2 m temperature 6 hours ahead with an area-weighted (cosine-latitude) RMSE of **1.214 K**, half that of persistence (2.457 K). Its error is lowest in the tropics (0.84 K) and highest near the poles (1.55 K). For total precipitation, the RMSE is $6.82 \times 10^{-4}$ m per 6 h against $9.03 \times 10^{-4}$ for persistence, a 24 % reduction. Training ran for 81 epochs before early stopping, with the best validation loss of 0.1173 at epoch 70; the gap of about 9 % between training and validation loss at that point (0.1079 vs. 0.1173) is mild overfitting at most.





Figure 13: Global 2 m temperature for a held-out test sample: ERA5 truth (left), model prediction (middle) and error (right). Sample RMSE 1.15 K.





Figure 14: Time-averaged 2 m temperature RMSE at each grid point over the test set (global 1.21 K, tropics 0.84 K, poles 1.55 K).





Figure 15: Weather-model loss convergence and the learning-rate schedule.

An ablation changes one thing in the main model, retrains it, and scores every variant on the same test samples (one training run per variant; gaps below about 0.05 K are within seed noise):

| Variant | Niño 3.4 input | Loss | T2m RMSE (K) |
|---|---|---|---|
| Unweighted loss | observed | unweighted | **1.164** |
| Unconditioned | none (zero) | area-weighted | 1.195 |
| Main model | observed | area-weighted | 1.214 |
| Conditioned on TFT forecasts | TFT lead-1 forecast | area-weighted | 1.335 |
| Persistence | — | — | 2.457 |

Niño 3.4 adds nothing at 6 hours, and feeding in TFT forecasts makes the model worse (+0.12 K); the area-weighted loss does not help either. Over a 6-hour step, the recent atmospheric state carries far more information than a monthly ENSO index, and 2020–2021 spans a single La Niña episode, so the index barely varies.

### 5.2.7 Warming Rate

A trend of joined straight segments is fitted to 76 annual global means (GISTEMP, 1950–2025), and BIC decides how many breaks the data justify.

| Change-points $K$ | 0 | 1 | 2 | 3 |
|---|---|---|---|---|
| BIC | −105.7 | **−114.1** | −113.7 | −105.7 |

BIC selects one change-point, in **1976**, where the warming rate rises from 0.035 to 0.211 °C per decade (residual AR(1) coefficient $\phi = 0.27$). The location is loosely constrained: break years from 1970 to 1986 fit almost as well ($\Delta$BIC < 2). A second break in 2012 is nearly tied ($\Delta$BIC = 0.4; the two-break fit places breaks in 1971 and 2012, with a final rate of 0.363 °C per decade). Warming therefore sped up about six-fold in the mid-1970s, while a further acceleration after 2012 is possible but not yet established.





Figure 16: GISTEMP anomalies: monthly values (grey), annual means (black points) and the BIC-selected piecewise-linear trend with the 1976 change-point.

### 5.2.8 Training Logs

The logs below are verbatim lines from the main run on 27 October 2026; only the logger prefix (level and module) is trimmed.

```
# El Niño main run
14:31:54 Prepared TFT DataFrame: 921 timesteps, 15 features
14:31:57 Split: train=644 months, val=138 months, test=139 months;
         windows train=573, val=127, test=128
14:31:57 Created TFT model: 96,047 parameters
14:31:57 Training TFT: 100 epochs, batch_size=32
14:32:50 TFT training complete. Best val_loss=0.287493
14:32:52 Created LSTM model: 142,668 params, hidden=64, layers=2,
         bidirectional=True
14:33:03 Early stopping at epoch 21
14:33:03 LSTM training complete. Best val_loss=0.337486

# Pollution A3TGCN
14:33:17 Created PollutionSTGNN: 6,984 params, hidden=32, periods=7
14:33:17 Training PollutionSTGNN: 100 epochs, lr=0.001, device=cpu,
         params=6,984
14:33:20 Epoch 1/100: train=1.741392, val=0.380588
14:37:44 Epoch 100/100: train=0.968618, val=0.301509

# Main weather model
14:37:52 NWM conditioning=observed (nino_features.parquet),
         area_weighted_loss=True
14:37:53 Training NWM ST-GNN: 100 epochs, device=cpu, lr=0.001, wd=0.0001
14:39:11 Epoch 1/100: train_loss=0.286078, val_loss=0.193916,
         val_rmse=0.440358
16:32:23 Epoch 80/100: train_loss=0.107055, val_loss=0.117450,
```

```
         val_rmse=0.342710
16:33:47 Early stopping triggered at epoch 81.
16:33:47 NWM training complete. Best val_loss=0.117334

# Ablation (nwm_ablation/forecast/train_nwm_forecast.log)
16:53:06 NWM conditioning=forecast (nino34_tft_lead1.parquet),
         area_weighted_loss=True
22:06:03 Early stopping triggered at epoch 85.

# Evaluation (make gen-eval-artifacts)
23:04:07 NWM test (observed): T2m RMSE 1.214 K (persistence 2.457 K),
         precip RMSE 6.821e-04 (persistence 9.032e-04)
```

The TFT trained in under a minute, and its best validation loss of 0.2875 matches its checkpoint name. The pollution model ran all 100 epochs and keeps the best validation checkpoint. The main weather model trained from 14:37 to 16:33 (about 86 seconds per epoch), the three ablation variants trained in parallel from 16:53 to 22:06, and the logged test RMSE is exactly the 1.214 K reported.

### 5.2.9 Patent Application Status

The draft application has 15 claims covering the spherical-mesh weather model, a fast graph-batching method and the four-module system.

| Claims | Subject |
|---|---|
| 1–7 | Fibonacci-mesh weather model with ENSO conditioning |
| 8–9 | Loop-free batching of graph edge indices |
| 10–13 | Four-module system, TFT forecasts driving the weather model |
| 14–15 | Matching method and storage-medium claims |

The application is still a draft: inventor names remain to be completed, Section 7 now discloses the ablation results, and revised claim wording has been drafted for the patent agent. Claims built on ENSO coupling (1(d), 3, 10 and 14) conflict with the disclosed results and need revising; the mesh and batching claims are the strongest.

### 5.2.10 Conclusions by Model

| Model | What the evidence supports | Next step |
|---|---|---|
| El Niño | TFT and LSTM are comparable and beat persistence from 2–3 months ahead | Remove the shared cold bias |
| Air pollution | About 10 % below persistence on concentration | A longer download to test exceedance skill |
| Weather | Halves persistence error at 6 hours | ENSO coupling is not supported on two years of data |
| Warming rate | A regime change around 1976, from 0.03 to 0.21 °C per decade | A second break after 2012 is possible but not yet established |

The integration works and is reproducible, but the scientific claims must stay modest, and ENSO coupling of the weather model is not supported.

### 5.3 Future Work

To strengthen the scientific results and extend the system, the following enhancements are recommended:

- **Remove the El Niño cold bias:** Diagnose the shared bias and recalibrate the TFT intervals, for example with split conformal quantile regression, which guarantees the nominal coverage of the interval on exchangeable data.
- **Larger pollution record:** Download a longer and denser OpenAQ record (the current download is capped at 10,000 readings) so that exceedance skill can be measured, and add satellite $NO_2$ and methane data to extend coverage beyond the 75 stations.
- **Multi-year weather training:** Train the weather model on several years spanning different ENSO phases, and at longer lead times, before ENSO conditioning can be properly tested.
- **Other climate oscillations:** Extend the TFT to the Indian Ocean Dipole and the North Atlantic Oscillation to test whether it generalises beyond ENSO.
- **Higher resolution and GPUs:** Move from the 5.625° grid to finer ERA5 resolutions on GPU hardware, which the code already supports, so that smaller-scale weather features can be resolved.

# 6. BIBLIOGRAPHY


- Paul, A., & Shekhawat, H. (2026). ClimatePulse: Temporal Fusion Transformers and Spatio-Temporal Graph Networks in Climate Modeling. Pre-print draft, `docs/research_paper/research_paper.md`.
- Lim, B., Arık, S. Ö., Loeff, N., & Pfister, T. (2021). Temporal Fusion Transformers for Interpretable Multi-horizon Time Series Forecasting. *International Journal of Forecasting*, 37(4), 1748–1764.
- Bai, J., Zhu, J., Song, Y., Zhao, L., Hou, Z., Du, R., & Li, H. (2021). A3T-GCN: Attention Temporal Graph Convolutional Network for Traffic Forecasting. *ISPRS International Journal of Geo-Information*, 10(7), 485.
- Hochreiter, S., & Schmidhuber, J. (1997). Long Short-Term Memory. *Neural Computation*, 9(8), 1735–1780.
- Kipf, T. N., & Welling, M. (2017). Semi-Supervised Classification with Graph Convolutional Networks. *ICLR 2017*.
- Hamilton, W. L., Ying, R., & Leskovec, J. (2017). Inductive Representation Learning on Large Graphs. *NeurIPS*, 30.
- Lam, R., Sanchez-Gonzalez, A., Willson, M., et al. (2022). GraphCast: Learning Skillful Medium-Range Global Weather Forecasting. *arXiv:2212.12794*.
- Keisler, R. (2022). Forecasting Global Weather with Graph Neural Networks. *arXiv:2202.07575*.
- Ham, Y. G., Kim, J. H., & Luo, J. J. (2019). Deep Learning for Multi-year ENSO Forecasts. *Nature*, 573, 568–572.
- Barnston, A. G., Tippett, M. K., L'Heureux, M. L., Li, S., & DeWitt, D. G. (2012). Skill of Real-Time Seasonal ENSO Model Predictions During 2002–11. *Bulletin of the American Meteorological Society*, 93(5), 631–651.
- Prais, S. J., & Winsten, C. B. (1954). Trend Estimators and Serial Correlation. *Cowles Commission Discussion Paper*, Statistics No. 383.
- Schwarz, G. (1978). Estimating the Dimension of a Model. *Annals of Statistics*, 6(2), 461–464.
- Li, Y., Yu, R., Shahabi, C., & Liu, Y. (2018). Diffusion Convolutional Recurrent Neural Network: Data-Driven Traffic Forecasting. *ICLR 2018*.
- Pathak, J., Subramanian, S., Harrington, P., et al. (2022). FourCastNet: A Global Data-Driven High-Resolution Weather Model Using Adaptive Fourier Neural Operators. *arXiv:2202.11214*.
- Hersbach, H., et al. (2020). The ERA5 Global Reanalysis. *Quarterly Journal of the Royal Meteorological Society*, 146(730), 1999–2049.
- NOAA Extended Reconstructed SST v5. https://psl.noaa.gov/data/gridded/data.noaa.ersst.v5.html
- OpenAQ: Open Air Quality Data. https://openaq.org/
- NASA GISS Surface Temperature Analysis (GISTEMP v4). https://data.giss.nasa.gov/gistemp/
- Climate TRACE: Global Emissions Inventory. https://climatetrace.org/
- WeatherBench 2 Documentation. https://weatherbench2.readthedocs.io/
- PyTorch Documentation. https://pytorch.org/docs/
- PyTorch Geometric Documentation. https://pytorch-geometric.readthedocs.io/
- pytorch-forecasting Documentation. https://pytorch-forecasting.readthedocs.io/
- Streamlit Documentation. https://docs.streamlit.io/