PROJECT PROPOSAL

ClimatePulse: AI-Powered Global Climate Forecasting and Environmental Telemetry Framework

---

EXECUTIVE SUMMARY

Global climate change presents an urgent need for fast, high-precision, and multi-scale environmental forecasting tools. Traditional Numerical Weather Prediction (NWP) systems are computationally prohibitive, requiring supercomputers running for hours per forecast cycle. Conversely, existing data-driven AI weather models operate in disciplinary isolation—predicting short-term atmospheric states without integrating long-range macro-climatic forcings (such as El Niño-Southern Oscillation, or ENSO) or local urban pollution transport.

ClimatePulse addresses this challenge by proposing a unified, modular, open-source Python framework that bridges four core climate modeling dimensions: 1. **El Niño Forecasting:** Probabilistic multi-horizon forecasting of the Niño 3.4 sea surface temperature index up to 60 months ahead. 2. **Pollution Dispersion:** Spatial-temporal forecasting of $PM_{2.5}$ concentrations across a global sensor network. 3. **Neural Weather Modeling (NWM):** Global weather forecasting operating on a uniform 3D spherical latent mesh conditioned on macro-climate ENSO signals. 4. **Climate Feedback Attribution:** Statistical Bayesian regression detecting structural regime shifts in historical global temperature anomalies.

This proposal outlines the technical vision, system architecture, dataset sources, technical stack, expected evaluation metrics, and system requirements for the ClimatePulse platform.

---

1. PROBLEM STATEMENT AND MOTIVATION

1.1 Limitations of Classical Numerical Weather Prediction (NWP)

Systems such as ECMWF's Integrated Forecasting System (IFS) rely on numerical integration of primitive fluid dynamic equations over high-resolution grids. While accurate, their extreme computational latency makes real-time scenario simulation, agricultural planning, and civil protection response difficult.

1.2 Limitations of Existing Data-Driven AI Weather Models

Recent neural weather models (GraphCast, FourCastNet, Pangu-Weather, FuXi) have reduced forecast latency by orders of magnitude. However, they suffer from key structural limitations:

1. **Equirectangular Polar Distortion:** Models operating on standard latitude-longitude grids apply spatially invariant convolutions across unequal physical surface areas. Equatorial cells subtend up to 8.6× more physical area than high-latitude cells, causing unweighted models to over-allocate computational capacity to polar regions.
2. **Single-Task Unconditioned Architecture:** Prior models map atmospheric states from time $t \to t+1$ without incorporating large-scale ocean-atmosphere forcing signals (such as ENSO phases) into latent message passing. Consequently, they treat El Niño and La Niña initializations identically.
3. **Rigidity of Icosahedral Meshes:** Prior graph-based models rely on icosahedral subdivision, locking computational mesh resolution to fixed discrete node counts $(20, 80, 320, 1280, 5120)$.

research.aditripaul.org

4. **Disciplinary Silos:** No unified platform connects global climate oscillations, local air quality dispersion, and global atmospheric field forecasting in a single reproducible framework.


---

2. SYSTEM ARCHITECTURE AND MODEL SUBSYSTEMS

ClimatePulse connects four specialized AI modeling components through a single data pipeline. The key innovation is **cross-domain conditioning**: long-range El Niño forecasts feed directly into the daily Neural Weather Model as a background climate signal.

2.1 Subsystem Overview

Model / Subsystem	Core Task	Architecture	Target Data Source
El Niño Forecaster	Predict ENSO phase 3–60 months ahead	Temporal Fusion Transformer (TFT) + BiLSTM baseline	NOAA ERSST v5
Pollution Dispersion	Spatiotemporal $PM_{2.5}$ / NO_2 transport	Attention-based Temporal Graph Conv (A3TGCN)	OpenAQ v3 Telemetry
Climate Feedback	Detect warming regime changepoints	Bayesian Hierarchical Regression (PyMC)	NASA GISTEMP v4 & Climate TRACE
Neural Weather Model	Global T_{2m} & precipitation conditioned on ENSO	Spherical Fibonacci Mesh Graph Neural Network (ST-GNN)	WeatherBench 2 / ERA5



research.aditripaul.org

 © 2026 research.aditripaul.org

2.2 Subsystem Details

2.2.1 El Niño Forecaster (Ocean Oscillation Module)

- **Role:** Predicts the phase and intensity of the El Niño-Southern Oscillation (ENSO) up to 60 months in advance using historical ocean temperature records.
- **How it Works:** Uses a Temporal Fusion Transformer (TFT) with self-attention and variable selection networks to process multi-year ocean temperature series.
- **Key Output:** Delivers probabilistic prediction intervals (10th, 50th, and 90th percentiles) for the Niño 3.4 index to capture forecast uncertainty.

2.2.2 Pollution Dispersion (Air Quality Module)

- **Role:** Forecasts local $PM_{2.5}$ pollutant concentrations across a network of global monitoring stations.
- **How it Works:** Combines spatial graph neural networks (which connect nearby sensor stations based on geographic distance) with temporal recurrent units to model how pollutants diffuse over space and accumulate over time.
- **Key Output:** Generates per-station concentration forecasts and automated air quality exceedance alert warnings.

2.2.3 Neural Weather Model (Global Weather Module)

- **Role:** Predicts global surface temperatures ($T_{2\mathrm{m}}$) and total precipitation fields.
- **How it Works:**
  1. **Uniform 3D Spherical Mesh:** Converts flat latitude-longitude maps into a 2,000-node 3D spherical mesh (using a golden-angle spiral) to eliminate polar surface distortion.
  2. **Macro-Climate Conditioning:** Injects the predicted El Niño state directly into the neural network's internal processor layers so that the global forecast is dynamically shaped by the active climate phase.
  3. **Physical Area Weighting:** Weights the learning objective by the physical surface area of each grid cell to ensure fair global prediction accuracy.
- **Key Output:** Gridded global atmospheric predictions for upcoming forecast windows.

2.2.4 Climate Feedback Attribution (Regime Shift Module)

- **Role:** Analyzes long-term historical global temperature trends (1950–present) to identify structural shifts in global warming rates.
- **How it Works:** Fits a Bayesian piecewise regression model over historical surface temperature anomaly records and greenhouse gas emissions inventories.
- **Key Output:** Identifies statistical changepoints and estimates historical climate sensitivity trends.

---

3. DATASET SOURCES AND INGESTION PIPELINE

Data ingestion is constrained to the **1950–present** window for high-fidelity historical relevance. All datasets are fetched programmatically via the unified data ingestion pipeline:

research.aditripaul.org

 © 2026 research.aditripaul.org

Dataset Source	Content & Variables	Ingestion Access Mechanism	Primary Subsystem Usage
NOAA ERSST v5	Monthly sea surface temperature (SST) anomalies & derived Niño 3.4 index	OPeNDAP Protocol	El Niño Forecaster (TFT / LSTM)
OpenAQ v3	Spatiotemporal $PM_{2.5}$, NO_2, O_3 measurements from global monitoring stations	REST API (API key authenticated)	Pollution Dispersion (A3TGCN)
NASA GISTEMP v4	Long-term global surface temperature anomaly records (1880–present)	Direct HTTPS Download	Climate Feedback (Bayesian)
Climate TRACE v7	Multi-sector greenhouse gas emissions inventories (CO_2, CH_4, N_2O)	REST API	Climate Feedback (Bayesian)
WeatherBench 2 / ERA5	6-hourly 2m temperature (T_{2m}) & total precipitation (5.625° equirectangular grid)	Google Cloud Storage Zarr	Neural Weather Model (ST-GNN)

All ingested data are processed and stored using high-performance Parquet format for bulk time-series data and JSON for metadata tracking.

---

4. TECHNICAL STACK AND SYSTEM REQUIREMENTS

4.1 Technical Stack

System Layer	Libraries & Technologies
Deep Learning	PyTorch 2.12, torch-geometric 2.8, pytorch-forecasting 1.8
Bayesian Modeling	PyMC 6, ArviZ 1.2
Data Engineering	xarray 2026.7, pandas 3, pyarrow 25, zarr 3, gcsfs
Configuration	Pydantic-Settings v2, PyYAML
Logging / Artifacts	Loguru, CSV metrics logger

4.2 Hardware Compatibility

- **Hardware Acceleration:** Fully compatible with NVIDIA CUDA, AMD ROCm 7.2 (`HSA_OVERRIDE_GFX_VERSION=10.3.0`), and Intel NPU via OpenVINO.
- **Resource Footprint:** Designed for efficient training and inference on workstation hardware with $\geq$ 15 GB VRAM.

research.aditripaul.org

4.3 Build and Workflow Automation

The platform includes an automated command-line workflow for environment setup, data ingestion, training, artifact generation, and document compilation:

```
make install                    # Install environment and dependencies
make download-data              # Ingest ERSST, OpenAQ, GISTEMP, and Climate TRACE data
make download-nwm-data          # Ingest ERA5 WeatherBench 2 Zarr data
make train                      # Train all four model families
make gen-eval-artifacts         # Generate evaluation plots and summary metrics
make gen-research-paper-pdf     # Compile research paper PDF
make save-run                   # Snapshot checkpoints and evaluation artifacts
```

---

5. EXPECTED OUTCOMES AND EVALUATION STRATEGY

The framework will be evaluated against standardized meteorological benchmarks and open environmental observation datasets:

5.1 Evaluation Framework and Target Metrics

Model / Subsystem	Primary Evaluation Metrics	Target Performance Criteria	Evaluation Dataset
El Niño Forecaster (TFT)	Skill Score, RMSE, Anomaly Correlation Coefficient (ACC)	Skill Score > 0.85; RMSE < 0.40°C over 12-month lead horizon	NOAA ERSST v5 (1950–present)
Pollution ST-GNN (A3TGCN)	F_1 Score, Precision, Recall, Station RMSE	$F_1 > 0.85$ on AQI exceedance alerts ($> 100\,\mu g/m^3$)	OpenAQ Sensor Network Telemetry
Neural Weather Model (ST-GNN)	Latitude-Weighted Validation MSE, Convergence Rate	$> 35\%$ forecast error reduction relative to unweighted initialization	ERA5 WeatherBench 2 Reanalysis (5.625°)
Climate Feedback (Bayesian)	Posterior Trace Convergence, Changepoint Count	Posterior convergence ($\hat{R} \leq 1.05$) across monthly temperature series	NASA GISTEMP v4 Surface Anomaly Records

---

6. CONCLUSION

ClimatePulse establishes a novel, highly efficient paradigm for multi-modal climate forecasting. By unifying long-horizon ocean-atmosphere forecasting, local sensor-network pollution transport, and global spherical neural weather modeling into a single reproducible open-source platform, the project delivers state-of-the-art predictive performance while maintaining a low computational footprint.

research.aditripaul.org

 © 2026 research.aditripaul.org