Observed strong lenses and their matched C_bright targets

Complete 43-pair atlas • exact retained pixel payloads

Each following page places the exact observed-reference cutout used in the final authoritative analysis beside the exact bare catalogue-negative target selected for that reference. The 43 targets are the final targets inherited by C_bright.

The assignment is score-blind, without replacement, and globally optimized for maximum cardinality then minimum total robust-IQR distance. It is not 43 independent nearest-neighbour choices. Matching uses observed z magnitude, catalogue Sérsic index, axis ratio, r-band PSF FWHM, and r-band depth.

All panels show the complete 101×101-pixel (≈26.5 arcsec) g/r/z cutout with the same fixed Legacy Survey/Stein dr2_rgb mapping (m=0.03, Q=20). There is no panel-specific crop or intensity stretch. The target is shown before source insertion; C_bright subsequently rescales the retained θE=1.5-arcsec source component on this target.

Visual similarity is not a matching criterion. The atlas is a diagnostic of what the five-covariate matching does—and does not—control.


Pairs: 43
matches.parquet SHA-256: b16127143b0c9c1e62ddd7829209f2dd47a5d6e04435cf71e025c146913373ee
reference_adapter.parquet SHA-256: 43ead962c7371d42c6a9ad3f4f75861b60821c815017f79c507e7de2c253a1fb
pair_plan.parquet SHA-256: 39f3b19e232f09aef40e8d6a107bf17ec80e655f410983cb40d7707e95218e2e
brightness_design.parquet SHA-256: c09bf29230064e4a6a306bb6a6aac015c70e7e0fcf889314e58c3fb72c53b7ef

Pair 01 of 43: exact observed and matched-target pixels

global match distance = 0.108

admissible targets for this reference = 2,880

Observed confirmed lens
DESI-036.1436-00.0411 • 0361p000_6437



Matched bare target inherited by C_bright
0455m130_7409 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	19.732	19.827	-0.090	
Sérsic index	2.162	2.005	+0.052	
Axis ratio	0.527	0.532	-0.012	
r-band PSF FWHM	1.245 arcsec	1.243 arcsec	+0.007	
r-band 5σ depth	25.030 mag	25.041 mag	-0.028	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 02 of 43: exact observed and matched-target pixels

global match distance = 0.675

admissible targets for this reference = 862

Observed confirmed lens
DESI-044.2865-23.4348 • 0442m235_7484



Matched bare target inherited by C_bright
0308p025_8046 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	16.977	17.066	-0.085	
Sérsic index	4.000	2.180	+0.607	
Axis ratio	0.602	0.504	+0.278	
r-band PSF FWHM	1.139 arcsec	1.155 arcsec	-0.052	
r-band 5σ depth	25.066 mag	25.077 mag	-0.024	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 03 of 43: exact observed and matched-target pixels


global match distance = 0.603

admissible targets for this reference = 669

Observed confirmed lens
DESI-053.6254-13.1866 • 0537m132_1571



Matched bare target inherited by C_bright
1741m032_9307 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	17.046	16.606	+0.417	
Sérsic index	2.558	2.562	-0.001	
Axis ratio	0.723	0.785	-0.179	
r-band PSF FWHM	1.225 arcsec	1.170 arcsec	+0.177	
r-band 5σ depth	25.198 mag	25.347 mag	-0.356	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene

Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 04 of 43: exact observed and matched-target pixels


global match distance = 0.322

admissible targets for this reference = 4,201

Observed confirmed lens
DESI-057.2074-10.2962 • 0572m102_2469



Matched bare target inherited by C_bright
3543m095_616 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	18.401	18.337	+0.061	
Sérsic index	5.828	6.000	-0.057	
Axis ratio	0.807	0.882	-0.213	
r-band PSF FWHM	1.149 arcsec	1.214 arcsec	-0.208	
r-band 5σ depth	24.431 mag	24.469 mag	-0.090	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 05 of 43: exact observed and matched-target pixels

global match distance = 0.433

admissible targets for this reference = 1,478

Observed confirmed lens
DESI-065.6447-28.0652 • 0656m280_3774



Matched bare target inherited by C_bright
0116p032_9709 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	18.168	18.520	-0.334	
Sérsic index	4.899	4.873	+0.008	
Axis ratio	0.689	0.732	-0.124	
r-band PSF FWHM	1.230 arcsec	1.227 arcsec	+0.012	
r-band 5σ depth	25.265 mag	25.163 mag	+0.246	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 06 of 43: exact observed and matched-target pixels


global match distance = 0.578

admissible targets for this reference = 1,205

Observed confirmed lens
DESI-072.9777+00.1050 • 0728p000_6851



Matched bare target inherited by C_bright
0739m060_4888 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	16.724	16.562	+0.154	
Sérsic index	5.659	5.301	+0.119	
Axis ratio	0.677	0.528	+0.426	
r-band PSF FWHM	1.189 arcsec	1.268 arcsec	-0.252	
r-band 5σ depth	24.450 mag	24.356 mag	+0.226	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 07 of 43: exact observed and matched-target pixels


global match distance = 0.377

admissible targets for this reference = 3,893

Observed confirmed lens
DESI-073.5286-10.2227 • 0734m102_4599



Matched bare target inherited by C_bright
0736m020_870 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	18.242	18.321	-0.075	
Sérsic index	4.000	3.029	+0.324	
Axis ratio	0.269	0.287	-0.052	
r-band PSF FWHM	1.140 arcsec	1.087 arcsec	+0.171	
r-band 5σ depth	24.540 mag	24.547 mag	-0.016	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 08 of 43: exact observed and matched-target pixels


global match distance = 0.473

admissible targets for this reference = 945

Observed confirmed lens
DESI-074.9643-30.7236 • 0750m307_2398



Matched bare target inherited by C_bright
0542m162_541 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	17.479	17.689	-0.199	
Sérsic index	6.000	6.000	+0.000	
Axis ratio	0.704	0.731	-0.079	
r-band PSF FWHM	1.369 arcsec	1.263 arcsec	+0.338	
r-band 5σ depth	25.215 mag	25.110 mag	+0.252	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 09 of 43: exact observed and matched-target pixels


global match distance = 0.601

admissible targets for this reference = 1,103

Observed confirmed lens
DESI-075.2790-24.4179 • 0753m245_3999



Matched bare target inherited by C_bright
0241p017_8218 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	18.363	18.695	-0.315	
Sérsic index	6.000	5.355	+0.215	
Axis ratio	0.694	0.670	+0.069	
r-band PSF FWHM	1.203 arcsec	1.185 arcsec	+0.058	
r-band 5σ depth	25.395 mag	25.205 mag	+0.456	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 10 of 43: exact observed and matched-target pixels


global match distance = 0.223

admissible targets for this reference = 475

Observed confirmed lens
DESI-086.3066-26.5884 • 0862m265_8849



Matched bare target inherited by C_bright
3476p012_4582 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	16.659	16.629	+0.028	
Sérsic index	6.000	5.540	+0.153	
Axis ratio	0.614	0.667	-0.150	
r-band PSF FWHM	1.115 arcsec	1.118 arcsec	-0.008	
r-band 5σ depth	25.104 mag	25.127 mag	-0.054	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 11 of 43: exact observed and matched-target pixels


global match distance = 0.616

admissible targets for this reference = 168

Observed confirmed lens
DESI-087.1525-36.2427 • 0870m362_8874



Matched bare target inherited by C_bright
0487m112_573 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	15.914	16.453	-0.510	
Sérsic index	3.693	3.228	+0.155	
Axis ratio	0.812	0.758	+0.152	
r-band PSF FWHM	1.209 arcsec	1.272 arcsec	-0.199	
r-band 5σ depth	25.188 mag	25.113 mag	+0.180	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 12 of 43: exact observed and matched-target pixels


global match distance = 0.530

admissible targets for this reference = 1,139

Observed confirmed lens
DESI-090.9854-35.9683 • 0910m360_2444



Matched bare target inherited by C_bright
0250m157_4602 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	17.290	17.536	-0.233	
Sérsic index	3.662	3.346	+0.105	
Axis ratio	0.683	0.843	-0.457	
r-band PSF FWHM	1.295 arcsec	1.287 arcsec	+0.024	
r-band 5σ depth	25.099 mag	25.133 mag	-0.080	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 13 of 43: exact observed and matched-target pixels

global match distance = 0.577

admissible targets for this reference = 1,766

Observed confirmed lens
DESI-122.0852+10.5284 • 1219p105_7293



Matched bare target inherited by C_bright
1678p005_7456 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	17.106	17.641	-0.507	
Sérsic index	4.963	5.183	-0.073	
Axis ratio	0.829	0.858	-0.083	
r-band PSF FWHM	1.367 arcsec	1.312 arcsec	+0.176	
r-band 5σ depth	24.750 mag	24.825 mag	-0.181	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 14 of 43: exact observed and matched-target pixels


global match distance = 0.363

admissible targets for this reference = 3,967

Observed confirmed lens
DESI-140.8110+18.4954 • 1409p185_769



Matched bare target inherited by C_bright
1819p127_283 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	18.512	18.574	-0.058	
Sérsic index	4.986	6.000	-0.338	
Axis ratio	0.799	0.793	+0.015	
r-band PSF FWHM	1.181 arcsec	1.215 arcsec	-0.109	
r-band 5σ depth	24.221 mag	24.238 mag	-0.041	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 15 of 43: exact observed and matched-target pixels


global match distance = 0.452

admissible targets for this reference = 4,229

Observed confirmed lens
DESI-143.3887+09.3219 • 1434p092_2623



Matched bare target inherited by C_bright
2286p032_2497 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	18.271	17.932	+0.321	
Sérsic index	6.000	6.000	+0.000	
Axis ratio	0.846	0.810	+0.103	
r-band PSF FWHM	1.424 arcsec	1.512 arcsec	-0.280	
r-band 5σ depth	24.575 mag	24.621 mag	-0.109	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 16 of 43: exact observed and matched-target pixels


global match distance = 0.378

admissible targets for this reference = 471

Observed confirmed lens
DESI-160.1716+18.8477 • 1601p187_3957



Matched bare target inherited by C_bright
0659m037_4323 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	16.034	15.972	+0.059	
Sérsic index	4.644	4.739	-0.032	
Axis ratio	0.529	0.492	+0.105	
r-band PSF FWHM	1.161 arcsec	1.191 arcsec	-0.097	
r-band 5σ depth	24.176 mag	24.319 mag	-0.344	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 17 of 43: exact observed and matched-target pixels


global match distance = 0.382

admissible targets for this reference = 3,766

Observed confirmed lens
DESI-161.4114-08.8359 • 1614m087_3206



Matched bare target inherited by C_bright
1611m087_2944 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	19.433	19.326	+0.102	
Sérsic index	4.000	3.034	+0.322	
Axis ratio	0.517	0.456	+0.175	
r-band PSF FWHM	1.009 arcsec	0.998 arcsec	+0.036	
r-band 5σ depth	24.757 mag	24.760 mag	-0.008	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 18 of 43: exact observed and matched-target pixels


global match distance = 0.471

admissible targets for this reference = 3,881

Observed confirmed lens
DESI-165.4754-06.0423 • 1655m060_944



Matched bare target inherited by C_bright
0603m130_1127 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	17.792	17.498	+0.278	
Sérsic index	2.399	2.217	+0.061	
Axis ratio	0.664	0.690	-0.073	
r-band PSF FWHM	1.262 arcsec	1.376 arcsec	-0.364	
r-band 5σ depth	24.528 mag	24.553 mag	-0.060	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 19 of 43: exact observed and matched-target pixels

global match distance = 0.221

admissible targets for this reference = 3,766

Observed confirmed lens
DESI-165.6876+12.1864 • 1658p122_150



Matched bare target inherited by C_bright
0056p242_3309 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	18.708	18.748	-0.038	
Sérsic index	4.000	4.000	+0.000	
Axis ratio	0.549	0.507	+0.122	
r-band PSF FWHM	1.443 arcsec	1.387 arcsec	+0.181	
r-band 5σ depth	24.103 mag	24.105 mag	-0.003	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 20 of 43: exact observed and matched-target pixels


global match distance = 0.559

admissible targets for this reference = 1,080

Observed confirmed lens
DESI-166.9974+04.1560 • 1671p042_499



Matched bare target inherited by C_bright
0160m127_3097 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	16.611	16.735	-0.117	
Sérsic index	2.052	1.748	+0.101	
Axis ratio	0.483	0.396	+0.249	
r-band PSF FWHM	1.228 arcsec	1.078 arcsec	+0.476	
r-band 5σ depth	24.365 mag	24.366 mag	-0.002	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 21 of 43: exact observed and matched-target pixels


global match distance = 0.613

admissible targets for this reference = 292

Observed confirmed lens
DESI-168.2943+23.9443 • 1683p240_1137



Matched bare target inherited by C_bright
1677p137_3334 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	15.723	16.311	-0.557	
Sérsic index	3.519	3.013	+0.169	
Axis ratio	0.817	0.823	-0.017	
r-band PSF FWHM	1.263 arcsec	1.256 arcsec	+0.021	
r-band 5σ depth	24.158 mag	24.237 mag	-0.190	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene

Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 22 of 43: exact observed and matched-target pixels


global match distance = 1.140

admissible targets for this reference = 2,223

Observed confirmed lens
DESI-168.7683+16.7607 • 1686p167_5461



Matched bare target inherited by C_bright
1613p235_2102 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	21.079	19.966	+1.053	
Sérsic index	1.000	0.662	+0.113	
Axis ratio	0.146	0.204	-0.166	
r-band PSF FWHM	1.263 arcsec	1.197 arcsec	+0.210	
r-band 5σ depth	24.347 mag	24.211 mag	+0.326	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 23 of 43: exact observed and matched-target pixels


global match distance = 0.254

admissible targets for this reference = 4,033

Observed confirmed lens
DESI-174.5481+14.7863 • 1744p147_5445



Matched bare target inherited by C_bright
1676m040_3617 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	18.548	18.382	+0.157	
Sérsic index	4.000	4.000	+0.000	
Axis ratio	0.695	0.688	+0.021	
r-band PSF FWHM	1.188 arcsec	1.250 arcsec	-0.198	
r-band 5σ depth	24.247 mag	24.245 mag	+0.005	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 24 of 43: exact observed and matched-target pixels


global match distance = 0.432

admissible targets for this reference = 1,583

Observed confirmed lens
DESI-189.5370+15.0309 • 1894p150_8726



Matched bare target inherited by C_bright
1177p297_5606 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	16.994	16.886	+0.102	
Sérsic index	5.589	6.000	-0.137	
Axis ratio	0.790	0.691	+0.284	
r-band PSF FWHM	1.388 arcsec	1.415 arcsec	-0.085	
r-band 5σ depth	24.724 mag	24.614 mag	+0.265	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 25 of 43: exact observed and matched-target pixels


global match distance = 0.462

admissible targets for this reference = 1,718

Observed confirmed lens
DESI-190.7929+21.3329 • 1908p212_1676



Matched bare target inherited by C_bright
1406p187_1596 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	17.184	16.954	+0.218	
Sérsic index	6.000	5.595	+0.135	
Axis ratio	0.865	0.855	+0.030	
r-band PSF FWHM	1.265 arcsec	1.178 arcsec	+0.279	
r-band 5σ depth	24.297 mag	24.187 mag	+0.263	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 26 of 43: exact observed and matched-target pixels


global match distance = 0.785

admissible targets for this reference = 686

Observed confirmed lens
DESI-192.9428+01.7155 • 1928p017_9908



Matched bare target inherited by C_bright
2406p160_14667 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	18.263	18.205	+0.055	
Sérsic index	4.000	3.572	+0.143	
Axis ratio	0.762	0.872	-0.313	
r-band PSF FWHM	1.457 arcsec	1.249 arcsec	+0.665	
r-band 5σ depth	25.749 mag	25.655 mag	+0.228	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 27 of 43: exact observed and matched-target pixels


global match distance = 0.578

admissible targets for this reference = 765

Observed confirmed lens
DESI-197.5704+14.7474 • 1976p147_245



Matched bare target inherited by C_bright
2338p082_5006 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	16.372	16.464	-0.088	
Sérsic index	6.000	6.000	+0.000	
Axis ratio	0.780	0.714	+0.188	
r-band PSF FWHM	1.194 arcsec	1.349 arcsec	-0.492	
r-band 5σ depth	24.351 mag	24.443 mag	-0.221	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 28 of 43: exact observed and matched-target pixels


global match distance = 0.747

admissible targets for this reference = 443

Observed confirmed lens
DESI-200.7678+03.7216 • 2006p037_9179



Matched bare target inherited by C_bright
0366p042_5754 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	16.577	16.458	+0.113	
Sérsic index	4.750	4.269	+0.160	
Axis ratio	0.800	0.659	+0.405	
r-band PSF FWHM	1.469 arcsec	1.361 arcsec	+0.342	
r-band 5σ depth	25.150 mag	24.946 mag	+0.490	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 29 of 43: exact observed and matched-target pixels


global match distance = 0.405

admissible targets for this reference = 522

Observed confirmed lens
DESI-202.6690+04.6707 • 2026p047_4274



Matched bare target inherited by C_bright
1721m032_8015 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	16.059	16.199	-0.133	
Sérsic index	5.424	6.000	-0.192	
Axis ratio	0.864	0.819	+0.128	
r-band PSF FWHM	1.499 arcsec	1.541 arcsec	-0.134	
r-band 5σ depth	24.660 mag	24.773 mag	-0.273	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 30 of 43: exact observed and matched-target pixels


global match distance = 0.510

admissible targets for this reference = 447

Observed confirmed lens
DESI-216.8280-06.7541 • 2168m067_3815



Matched bare target inherited by C_bright
2558p135_1477 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	15.999	16.217	-0.206	
Sérsic index	6.000	6.000	+0.000	
Axis ratio	0.840	0.704	+0.387	
r-band PSF FWHM	1.388 arcsec	1.448 arcsec	-0.192	
r-band 5σ depth	24.237 mag	24.311 mag	-0.176	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 31 of 43: exact observed and matched-target pixels


global match distance = 0.315

admissible targets for this reference = 3,835

Observed confirmed lens
DESI-220.4549+14.6891 • 2204p147_3771



Matched bare target inherited by C_bright
1284p212_1624 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	18.429	18.402	+0.025	
Sérsic index	4.000	4.000	+0.000	
Axis ratio	0.788	0.875	-0.247	
r-band PSF FWHM	1.322 arcsec	1.263 arcsec	+0.187	
r-band 5σ depth	24.140 mag	24.118 mag	+0.051	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 32 of 43: exact observed and matched-target pixels


global match distance = 0.461

admissible targets for this reference = 4,161

Observed confirmed lens
DESI-227.5364+20.6236 • 2276p205_1624



Matched bare target inherited by C_bright
3356p142_4813 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	18.148	18.213	-0.061	
Sérsic index	6.000	5.613	+0.129	
Axis ratio	0.856	0.933	-0.218	
r-band PSF FWHM	1.402 arcsec	1.512 arcsec	-0.350	
r-band 5σ depth	24.468 mag	24.406 mag	+0.148	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 33 of 43: exact observed and matched-target pixels


global match distance = 0.294

admissible targets for this reference = 1,298

Observed confirmed lens
DESI-234.8707+16.8379 • 2349p167_2044



Matched bare target inherited by C_bright
3368m047_607 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	16.886	16.958	-0.067	
Sérsic index	5.457	6.000	-0.181	
Axis ratio	0.798	0.834	-0.102	
r-band PSF FWHM	1.116 arcsec	1.153 arcsec	-0.117	
r-band 5σ depth	24.260 mag	24.194 mag	+0.158	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 34 of 43: exact observed and matched-target pixels


global match distance = 0.514

admissible targets for this reference = 3,476

Observed confirmed lens
DESI-252.9173+28.0881 • 2530p280_351



Matched bare target inherited by C_bright
3393p302_2962 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	19.161	19.352	-0.181	
Sérsic index	4.000	4.000	+0.000	
Axis ratio	0.640	0.660	-0.059	
r-band PSF FWHM	1.813 arcsec	1.689 arcsec	+0.397	
r-band 5σ depth	24.374 mag	24.264 mag	+0.265	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 35 of 43: exact observed and matched-target pixels


global match distance = 0.741

admissible targets for this reference = 309

Observed confirmed lens
DESI-260.8405+23.8442 • 2608p237_3728



Matched bare target inherited by C_bright
2317p295_3494 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	15.737	15.797	-0.057	
Sérsic index	2.169	1.713	+0.152	
Axis ratio	0.629	0.705	-0.217	
r-band PSF FWHM	1.579 arcsec	1.371 arcsec	+0.662	
r-band 5σ depth	24.205 mag	24.124 mag	+0.194	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 36 of 43: exact observed and matched-target pixels


global match distance = 0.284

admissible targets for this reference = 3,946

Observed confirmed lens
DESI-310.8020-06.1649 • 3108m062_2639



Matched bare target inherited by C_bright
0674m037_5062 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	18.431	18.651	-0.209	
Sérsic index	4.000	4.000	+0.000	
Axis ratio	0.521	0.584	-0.181	
r-band PSF FWHM	1.310 arcsec	1.312 arcsec	-0.006	
r-band 5σ depth	24.196 mag	24.223 mag	-0.065	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 37 of 43: exact observed and matched-target pixels


global match distance = 0.761

admissible targets for this reference = 372

Observed confirmed lens
DESI-327.8408+13.7884 • 3277p137_6699



Matched bare target inherited by C_bright
2303p277_5802 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	15.808	16.025	-0.205	
Sérsic index	4.336	4.432	-0.032	
Axis ratio	0.561	0.597	-0.104	
r-band PSF FWHM	1.186 arcsec	1.271 arcsec	-0.270	
r-band 5σ depth	24.567 mag	24.287 mag	+0.672	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 38 of 43: exact observed and matched-target pixels


global match distance = 0.456

admissible targets for this reference = 982

Observed confirmed lens
DESI-341.0206+27.9877 • 3409p280_5325



Matched bare target inherited by C_bright
2604p252_6396 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	16.604	16.720	-0.110	
Sérsic index	3.555	2.532	+0.341	
Axis ratio	0.643	0.686	-0.123	
r-band PSF FWHM	1.448 arcsec	1.503 arcsec	-0.174	
r-band 5σ depth	24.153 mag	24.230 mag	-0.185	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 39 of 43: exact observed and matched-target pixels


global match distance = 0.507

admissible targets for this reference = 2,207

Observed confirmed lens
DESI-341.8012-02.0939 • 3418m020_1357



Matched bare target inherited by C_bright
0387p212_2812 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	17.358	17.751	-0.372	
Sérsic index	2.761	3.026	-0.089	
Axis ratio	0.937	0.866	+0.203	
r-band PSF FWHM	1.340 arcsec	1.414 arcsec	-0.236	
r-band 5σ depth	24.505 mag	24.554 mag	-0.119	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 40 of 43: exact observed and matched-target pixels

global match distance = 0.666

admissible targets for this reference = 266

Observed confirmed lens
DESI-342.9290-03.4136 • 3428m035_4814



Matched bare target inherited by C_bright
1991m045_2087 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	15.713	15.671	+0.039	
Sérsic index	6.000	6.000	+0.000	
Axis ratio	0.948	0.761	+0.534	
r-band PSF FWHM	1.373 arcsec	1.496 arcsec	-0.391	
r-band 5σ depth	24.156 mag	24.129 mag	+0.064	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 41 of 43: exact observed and matched-target pixels


global match distance = 0.429

admissible targets for this reference = 1,550

Observed confirmed lens
DESI-343.0402-04.2187 • 3431m042_1867



Matched bare target inherited by C_bright
2515p257_5020 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	16.936	16.651	+0.270	
Sérsic index	3.312	3.099	+0.071	
Axis ratio	0.677	0.730	-0.152	
r-band PSF FWHM	1.532 arcsec	1.493 arcsec	+0.123	
r-band 5σ depth	24.476 mag	24.368 mag	+0.261	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 42 of 43: exact observed and matched-target pixels


global match distance = 0.432

admissible targets for this reference = 1,232

Observed confirmed lens
DESI-345.0719+22.2249 • 3450p222_3227



Matched bare target inherited by C_bright
1788p032_4301 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	16.942	17.005	-0.059	
Sérsic index	4.731	5.027	-0.099	
Axis ratio	0.765	0.893	-0.366	
r-band PSF FWHM	1.752 arcsec	1.698 arcsec	+0.173	
r-band 5σ depth	24.144 mag	24.184 mag	-0.097	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene


Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch

Pair 43 of 43: exact observed and matched-target pixels


global match distance = 0.670

admissible targets for this reference = 425

Observed confirmed lens
DESI-345.8609+23.4760 • 3459p235_481



Matched bare target inherited by C_bright
0259p147_4049 • before source insertion



Matched covariates		observed	target	Δ(ref−target)/IQR
Observed z magnitude	15.990	16.177	-0.177	
Sérsic index	5.707	4.675	+0.344	
Axis ratio	0.699	0.587	+0.322	
r-band PSF FWHM	1.711 arcsec	1.606 arcsec	+0.335	
r-band 5σ depth	24.247 mag	24.367 mag	-0.288	

Score-blind global assignment • no replacement

Bare target shown; C_bright uses its retained θE=1.5″ parent scene

Full 101×101 field • fixed g/r/z mapping • no panel-specific stretch