

First and last name

Question 1/25

. When the correlation coefficient, r , is close to one:

- A. there is no relationship between the two variables
- B. it is impossible to tell if there is a relationship between the two variables
- C. there is a strong linear relationship between the two variables
- D. the slope of the regression line will be close to one

Question 2/25

Which of the following is an assumption underlying the use of the t-distributions?

- A. All Above
- B. The variance of the population is known
- C. s (sample standard deviation) is an unbiased estimate of the population variance.
- D. The samples are drawn from a normally distributed population

Question 3/25

The following are percentages of fat found in 5 samples of each of two brands of baby food:

A: 5.7, 4.5, 6.2, 6.3, 7.3

B: 6.3, 5.7, 5.9, 6.4, 5.1

Which of the following procedures is appropriate to test the hypothesis of equal average fat content in the two types of ice cream?

- A. Paired t-test with 5 d.f
- B. Paired t-test with 4 d.f
- C. Two samples t-test with 9 d.f
- D. Two samples t-test with 8 d.f

Question 4/25

Which of the following is not true about z score?

- A. Data points should be independent from each other. In other words, one data point isn't related or doesn't affect another data point.
- B. Your data should be normally distributed. However, for large sample sizes (over 30) this doesn't always matter.
- C. Your data should be randomly selected from a population, where each item has an equal chance of being selected.
- D. Your sample size is smaller than 30. Otherwise, use a t test.

Question 5/25

What is the relationship between sample size and the standard error of the mean?

- A. The standard error increases as the sample size increases.
- B. The standard error decreases as the sample size decreases.
- C. The standard error is unaffected by the sample size.
- D. The standard error decreases as the sample size increases.

Question 6/25

What does `scipy.stats.norm.sf(abs(z_scores))` represent?

- A. two-sided cdf
- B. one-sided cdf
- C. area under curve
- D. one-sided pdf

Question 7/25

Which of the following is a true statement, for comparing the t distributions with standard normal,

- A. The proportion of area beyond a specific value of “t” is less than the proportion of normal curve
- B. None of the Above
- C. The Normal Curve is symmetrical whereas the t-distributions are slightly skewed
- D. Greater the degree of freedom, the more the t-distribution resembles the standard normal distribution

Question 8/25

Which of the following statements sounds like a null hypothesis?

- A. The defendant is guilty
- B. There is a correlation in the population
- C. There is no difference between male and female incomes in the population
- D. The coin is not fair

Question 9/25

. Given IQ scores are approximately normally distributed with a mean of 100 and standard deviation of 15, the proportion of people with IQs above 130 is:

- A. 5%
- B. 95%
- C. 2.5%
- D. 68%

Question 10/25

In a small data sample ($N = 20$), what can we say about a z-score of 2.37?

- A. It is significant at $p < .01$
- B. It is significant at $p < .05$
- C. It is significant at $p < .001$
- D. It is non-significant

Question 11/25

. A Type I error is also known as a _____.

- A. False negative
- B. Double negative
- C. Positive negative
- D. False positive

Question 12/25

The critical value of a test statistic is determined from

- A. Calculations based on many actual repetitions of the same experiment
- B. The sampling distribution of the statistic assuming Alternative Hypothesis
- C. The sampling distribution of the statistics assuming Null Hypothesis
- D. Calculations from the data

Question 13/25

Of what is p the probability if the null hypothesis were true?

- A. p is the probability that the results would be replicated if the experiment was conducted a second time.
- B. p is the probability that the results are not due to chance, the probability that the null hypothesis (H_0) is false.
- C. p is the probability of observing a test statistic at least as big as the one we have if there were no effect in the population (i.e., the null hypothesis were true).
- D. p is the probability that the results are due to chance, the probability that the null hypothesis (H_0) is true.

Question 14/25

If you drew all possible samples from some population, calculated the mean for each of the samples, and constructed a line graph (showing the shape of the distribution) based on all of those means, what would you have?

- A. A population distribution
- B. parameter distribution
- C. A sampling distribution
- D. normal distribution

Question 15/25

For t distribution, increasing the sample size, the effect will be on

- A. The t -ratio
- B. Standard Error of the Means
- C. All of the Above
- D. Degrees of Freedom

Question 16/25

The use of the laws of probability to make inferences and draw statistical conclusions about populations based on sample data is referred to as _____.

- A. Sample statistics
- B. Inferential statistics
- C. Descriptive statistics
- D. Population statistics

Question 17/25

In statistical testing of the hypothesis, what happens to the region of rejection when the level of significance α is reduced?

- A. The rejection region is increased in size
- B. The rejection region is reduced in size
- C. The rejection region is unaltered
- D. The answer depends on the value of β

Question 18/25

Pr reasonably normally distributed with a mean of 18 and standard deviation of 6, determine the proportion of students with a 33 or higher.

- A. 0.0109
- B. 0.0217
- C. 0.0062
- D. 0.0124

Question 19/25

The .ppf() function represents

- A. percentage change in column value
- B. the probability for a given normal distribution value,
- C. the normal distribution value for which a given probability is the required value.
- D. standard normal distribution z - score

Question 20/25

Which of the following is NOT correct?

- A. The p-value measures the probability that the null hypothesis is true
- B. The probability of a type I error is controlled by the selection of the level of significance α
- C. The power of a test depends upon the sample size and the distance between the null hypothesis and the alternative hypothesis
- D. The probability of a type II error is controlled by the sample size (n)

Question 21/25

A range (set) of values within which the population parameter is expected to occur is called:

- A. confidence interval
- B. Confidence limits
- C. Level of confidence
- D. Point estimate

Question 22/25

For example, we have two different plant genotypes (genotype A and genotype B) and would like to compare if the yield of genotype A is significantly different from genotype B

- A. one sample paired t test
- B. two sample independent t test
- C. two sample unpaired t test
- D. two sample paired t test

Question 23/25

A _____ is a numerical characteristic of a sample and a _____ is a numerical characteristic of a population.

- A. Sample, population
- B. Population, sample
- C. Parameter, statistic
- D. Statistic, parameter

Question 24/25

1-alpha is the probability of

- A. Rejection Region
- B. Acceptance Region
- C. Type-II Error
- D. Type-I Error

Question 25/25

What is the parameter known as that expresses number of standard deviations a given value x falls from the mean, μ .

- A. significance level
- B. population interval
- C. z- score
- D. confidence interval