

Mixing Properties of Stable Random Fields Indexed by Amenable Groups

Parthanil Roy, Indian Statistical Institute

Joint work with **Mahan Mj (TIFR Mumbai)** and
Sourav Sarkar (University of Cambridge)

arXiv:2205.15849

Motivation and Connections

What is this work about?

Ergodic Theory


Operator Algebra

Probability Theory

Hyperbolic Geometry

Geometric Group Theory

What is this work about?



What is this work about?

Ergodic Theory

Operator Algebra

Probability Theory

Hyperbolic Geometry

Geometric Group Theory

What is this work about?

Ergodic Theory

(Nonsingular Actions)

Operator Algebra

(von Neumann Algebras)

Probability Theory

(Stable Random Fields)


Hyperbolic Geometry

(Patterson-Sullivan Theory)


Geometric Group Theory

(Amenable Groups)


What is this work about?




What is this work about?




What is this work about?




What is this work about?



What is this work about?



What is this work about?



A Crash Course on Stable Random Fields

Symmetric α -stable distribution

All random variables will be real valued and they will be defined on a common probability space (Ω, \mathcal{F}, P) . In particular, they all will be measurable maps $\Omega \rightarrow \mathbb{R}$.

Symmetric α -stable distribution

All random variables will be real valued and they will be defined on a common probability space (Ω, \mathcal{F}, P) . In particular, they all will be measurable maps $\Omega \rightarrow \mathbb{R}$.

A random variable X follows *symmetric α -stable* ($S\alpha S$) distribution ($0 < \alpha \leq 2$) with scale parameter $\sigma > 0$ (denoted by $X \sim S\alpha S(\sigma)$) if

$$E(e^{i\theta X}) = \int_{\Omega} e^{i\theta X(\omega)} dP(\omega) = e^{-\sigma^{\alpha} |\theta|^{\alpha}}, \quad \theta \in \mathbb{R}.$$

Symmetric α -stable distribution

All random variables will be real valued and they will be defined on a common probability space (Ω, \mathcal{F}, P) . In particular, they all will be measurable maps $\Omega \rightarrow \mathbb{R}$.

A random variable X follows *symmetric α -stable* ($S\alpha S$) distribution ($0 < \alpha \leq 2$) with scale parameter $\sigma > 0$ (denoted by $X \sim S\alpha S(\sigma)$) if

$$E(e^{i\theta X}) = \int_{\Omega} e^{i\theta X(\omega)} dP(\omega) = e^{-\sigma^{\alpha} |\theta|^{\alpha}}, \quad \theta \in \mathbb{R}.$$

- $X \stackrel{\mathcal{L}}{=} -X$.

Symmetric α -stable distribution

All random variables will be real valued and they will be defined on a common probability space (Ω, \mathcal{F}, P) . In particular, they all will be measurable maps $\Omega \rightarrow \mathbb{R}$.

A random variable X follows *symmetric α -stable* ($S\alpha S$) distribution ($0 < \alpha \leq 2$) with scale parameter $\sigma > 0$ (denoted by $X \sim S\alpha S(\sigma)$) if

$$E(e^{i\theta X}) = \int_{\Omega} e^{i\theta X(\omega)} dP(\omega) = e^{-\sigma^{\alpha} |\theta|^{\alpha}}, \quad \theta \in \mathbb{R}.$$

- $X \stackrel{\mathcal{L}}{=} -X$.
- $\alpha = 2 \Rightarrow X \sim \text{Gaussian}, \quad \alpha = 1 \Rightarrow X \sim \text{Cauchy}.$

Symmetric α -stable distribution

All random variables will be real valued and they will be defined on a common probability space (Ω, \mathcal{F}, P) . In particular, they all will be measurable maps $\Omega \rightarrow \mathbb{R}$.

A random variable X follows *symmetric α -stable* ($S\alpha S$) distribution ($0 < \alpha \leq 2$) with scale parameter $\sigma > 0$ (denoted by $X \sim S\alpha S(\sigma)$) if

$$E(e^{i\theta X}) = \int_{\Omega} e^{i\theta X(\omega)} dP(\omega) = e^{-\sigma^\alpha |\theta|^\alpha}, \quad \theta \in \mathbb{R}.$$

- $X \stackrel{\mathcal{L}}{=} -X$.
- $\alpha = 2 \Rightarrow X \sim \text{Gaussian}, \quad \alpha = 1 \Rightarrow X \sim \text{Cauchy}.$
- They arise as scaling limits of sums of IID symmetric random variables.

Symmetric α -stable distribution

All random variables will be real valued and they will be defined on a common probability space (Ω, \mathcal{F}, P) . In particular, they all will be measurable maps $\Omega \rightarrow \mathbb{R}$.

A random variable X follows *symmetric α -stable* ($S\alpha S$) distribution ($0 < \alpha \leq 2$) with scale parameter $\sigma > 0$ (denoted by $X \sim S\alpha S(\sigma)$) if

$$E(e^{i\theta X}) = \int_{\Omega} e^{i\theta X(\omega)} dP(\omega) = e^{-\sigma^\alpha |\theta|^\alpha}, \quad \theta \in \mathbb{R}.$$

- $X \stackrel{\mathcal{L}}{=} -X$.
- $\alpha = 2 \Rightarrow X \sim \text{Gaussian}, \quad \alpha = 1 \Rightarrow X \sim \text{Cauchy}.$
- They arise as scaling limits of sums of IID symmetric random variables.
- **Assume:** $0 < \alpha < 2$

Symmetric α -stable distribution

All random variables will be real valued and they will be defined on a common probability space (Ω, \mathcal{F}, P) . In particular, they all will be measurable maps $\Omega \rightarrow \mathbb{R}$.

A random variable X follows *symmetric α -stable* ($S\alpha S$) distribution ($0 < \alpha \leq 2$) with scale parameter $\sigma > 0$ (denoted by $X \sim S\alpha S(\sigma)$) if

$$E(e^{i\theta X}) = \int_{\Omega} e^{i\theta X(\omega)} dP(\omega) = e^{-\sigma^\alpha |\theta|^\alpha}, \quad \theta \in \mathbb{R}.$$

- $X \stackrel{\mathcal{L}}{=} -X$.
- $\alpha = 2 \Rightarrow X \sim \text{Gaussian}, \quad \alpha = 1 \Rightarrow X \sim \text{Cauchy}.$
- They arise as scaling limits of sums of IID symmetric random variables.
- **Assume:** $0 < \alpha < 2 \Rightarrow X$ is non-Gaussian.

$S_\alpha S$ random fields, $0 < \alpha < 2$

$S_\alpha S$ random fields, $0 < \alpha < 2$

Let (G, \cdot) be a countably infinite (possibly noncommutative) group with identity element e .

$S\alpha S$ random fields, $0 < \alpha < 2$

Let $(G, .)$ be a countably infinite (possibly noncommutative) group with identity element e .

A collection of random variables $\{X_g\}_{g \in G}$ is called an $S\alpha S$ random field if for all $k \geq 1$, for all $g_1, g_2, \dots, g_k \in G$ and for all $c_1, c_2, \dots, c_k \in \mathbb{R}$,

$$\sum_{i=1}^k c_i X_{g_i} \sim S\alpha S.$$

$S\alpha S$ random fields, $0 < \alpha < 2$

Let (G, \cdot) be a countably infinite (possibly noncommutative) group with identity element e .

A collection of random variables $\{X_g\}_{g \in G}$ is called an $S\alpha S$ random field if for all $k \geq 1$, for all $g_1, g_2, \dots, g_k \in G$ and for all $c_1, c_2, \dots, c_k \in \mathbb{R}$,

$$\sum_{i=1}^k c_i X_{g_i} \sim S\alpha S.$$

- The scale parameter depends on $c_1, c_2, \dots, c_k \in \mathbb{R}$.

$S\alpha S$ random fields, $0 < \alpha < 2$

Let (G, \cdot) be a countably infinite (possibly noncommutative) group with identity element e .

A collection of random variables $\{X_g\}_{g \in G}$ is called an $S\alpha S$ random field if for all $k \geq 1$, for all $g_1, g_2, \dots, g_k \in G$ and for all $c_1, c_2, \dots, c_k \in \mathbb{R}$,

$$\sum_{i=1}^k c_i X_{g_i} \sim S\alpha S.$$

- The scale parameter depends on $c_1, c_2, \dots, c_k \in \mathbb{R}$.
- We will soon describe the scale parameter for each linear combination

$S\alpha S$ random fields, $0 < \alpha < 2$

Let $(G, .)$ be a countably infinite (possibly noncommutative) group with identity element e .

A collection of random variables $\{X_g\}_{g \in G}$ is called an $S\alpha S$ random field if for all $k \geq 1$, for all $g_1, g_2, \dots, g_k \in G$ and for all $c_1, c_2, \dots, c_k \in \mathbb{R}$,

$$\sum_{i=1}^k c_i X_{g_i} \sim S\alpha S.$$

- The scale parameter depends on $c_1, c_2, \dots, c_k \in \mathbb{R}$.
- We will soon describe the scale parameter for each linear combination and this will uniquely specify the joint law of $\{X_g\}_{g \in G}$.

Kuelbs (1973) representation

Suppose $\{X_g\}_{g \in G}$ is an $S\alpha S$ random field.

Kuelbs (1973) representation

Suppose $\{X_g\}_{g \in G}$ is an $S\alpha S$ random field.

Then there exists a σ -finite standard measure space $(\Lambda, \mathcal{A}, \mu)$

Kuelbs (1973) representation

Suppose $\{X_g\}_{g \in G}$ is an $S\alpha S$ random field.

Then there exists a σ -finite standard measure space $(\Lambda, \mathcal{A}, \mu)$ and

$$\{f_g : g \in G\} \subset L^\alpha(\Lambda, \mu)$$

Kuelbs (1973) representation

Suppose $\{X_g\}_{g \in G}$ is an $S\alpha S$ random field.

Then there exists a σ -finite standard measure space $(\Lambda, \mathcal{A}, \mu)$ and

$$\{f_g : g \in G\} \subset L^\alpha(\Lambda, \mu) := \left\{ h : \Lambda \rightarrow \mathbb{R} \mid \|h\|_\alpha := \left(\int_\Lambda |h|^\alpha d\mu \right)^{1/\alpha} < \infty \right\}$$

Kuelbs (1973) representation

Suppose $\{X_g\}_{g \in G}$ is an $S\alpha S$ random field.

Then there exists a σ -finite standard measure space $(\Lambda, \mathcal{A}, \mu)$ and

$$\{f_g : g \in G\} \subset L^\alpha(\Lambda, \mu) := \left\{ h : \Lambda \rightarrow \mathbb{R} \mid \|h\|_\alpha := \left(\int_\Lambda |h|^\alpha d\mu \right)^{1/\alpha} < \infty \right\}$$

such that

$$\boxed{\sum_{i=1}^k c_i X_{g_i} \sim S\alpha S \left(\left\| \sum_{i=1}^k c_i f_{g_i} \right\|_\alpha \right)} \quad (1)$$

for all $c_1, c_2, \dots, c_k \in \mathbb{R}$ and for all $g_1, g_2, \dots, g_k \in G$.

Kuelbs (1973) representation

Suppose $\{X_g\}_{g \in G}$ is an $S\alpha S$ random field.

Then there exists a σ -finite standard measure space $(\Lambda, \mathcal{A}, \mu)$ and

$$\{f_g : g \in G\} \subset L^\alpha(\Lambda, \mu) := \left\{ h : \Lambda \rightarrow \mathbb{R} \mid \|h\|_\alpha := \left(\int_\Lambda |h|^\alpha d\mu \right)^{1/\alpha} < \infty \right\}$$

such that

$$\boxed{\sum_{i=1}^k c_i X_{g_i} \sim S\alpha S \left(\left\| \sum_{i=1}^k c_i f_{g_i} \right\|_\alpha \right)} \quad (1)$$

for all $c_1, c_2, \dots, c_k \in \mathbb{R}$ and for all $g_1, g_2, \dots, g_k \in G$.

The **converse** also holds.

Kuelbs (1973) representation

Suppose $\{X_g\}_{g \in G}$ is an $S\alpha S$ random field.

Then there exists a σ -finite standard measure space $(\Lambda, \mathcal{A}, \mu)$ and

$$\{f_g : g \in G\} \subset L^\alpha(\Lambda, \mu) := \left\{ h : \Lambda \rightarrow \mathbb{R} \mid \|h\|_\alpha := \left(\int_\Lambda |h|^\alpha d\mu \right)^{1/\alpha} < \infty \right\}$$

such that

$$\boxed{\sum_{i=1}^k c_i X_{g_i} \sim S\alpha S \left(\left\| \sum_{i=1}^k c_i f_{g_i} \right\|_\alpha \right)} \quad (1)$$

for all $c_1, c_2, \dots, c_k \in \mathbb{R}$ and for all $g_1, g_2, \dots, g_k \in G$.

The **converse** also holds.

$\{f_g\}_{g \in G} := \text{a (spectral) representation of } \{X_g\}_{g \in G}.$

Questions or Comments or Concerns?

(Left) stationary $S\alpha S$ random fields

An $S\alpha S$ random field $\{X_g\}_{g \in G}$ is called (left) stationary if for all $h \in G$,

$$\{X_{hg}\}_{g \in G} \stackrel{\mathcal{L}}{=} \{X_g\}_{g \in G},$$

(Left) stationary $S\alpha S$ random fields

An $S\alpha S$ random field $\{X_g\}_{g \in G}$ is called (left) stationary if for all $h \in G$,

$$\{X_{hg}\}_{g \in G} \stackrel{\mathcal{L}}{=} \{X_g\}_{g \in G},$$

i.e., for all $k \geq 1$, for all $g_1, g_2, \dots, g_k, h \in G$ and for all Borel $B \subset \mathbb{R}^k$,

$$P[(X_{hg_1}, X_{hg_2}, \dots, X_{hg_k}) \in B] = P[(X_{g_1}, X_{g_2}, \dots, X_{g_k}) \in B].$$

(Left) stationary $S\alpha S$ random fields

An $S\alpha S$ random field $\{X_g\}_{g \in G}$ is called (left) stationary if for all $h \in G$,

$$\{X_{hg}\}_{g \in G} \stackrel{\mathcal{L}}{=} \{X_g\}_{g \in G},$$

i.e., for all $k \geq 1$, for all $g_1, g_2, \dots, g_k, h \in G$ and for all Borel $B \subset \mathbb{R}^k$,

$$P[(X_{hg_1}, X_{hg_2}, \dots, X_{hg_k}) \in B] = P[(X_{g_1}, X_{g_2}, \dots, X_{g_k}) \in B].$$

In particular, X_g 's are identically distributed.

(Left) stationary $S\alpha S$ random fields

An $S\alpha S$ random field $\{X_g\}_{g \in G}$ is called (left) stationary if for all $h \in G$,

$$\{X_{hg}\}_{g \in G} \stackrel{\mathcal{L}}{=} \{X_g\}_{g \in G},$$

i.e., for all $k \geq 1$, for all $g_1, g_2, \dots, g_k, h \in G$ and for all Borel $B \subset \mathbb{R}^k$,

$$P[(X_{hg_1}, X_{hg_2}, \dots, X_{hg_k}) \in B] = P[(X_{g_1}, X_{g_2}, \dots, X_{g_k}) \in B].$$

In particular, X_g 's are identically distributed.

The law of a (left) stationary $S\alpha S$ random field is invariant under the (left) translation action on its indices.

(Left) stationary $S\alpha S$ random fields

An $S\alpha S$ random field $\{X_g\}_{g \in G}$ is called (left) stationary if for all $h \in G$,

$$\{X_{hg}\}_{g \in G} \stackrel{\mathcal{L}}{=} \{X_g\}_{g \in G},$$

i.e., for all $k \geq 1$, for all $g_1, g_2, \dots, g_k, h \in G$ and for all Borel $B \subset \mathbb{R}^k$,

$$P[(X_{hg_1}, X_{hg_2}, \dots, X_{hg_k}) \in B] = P[(X_{g_1}, X_{g_2}, \dots, X_{g_k}) \in B].$$

In particular, X_g 's are identically distributed.

The law of a (left) stationary $S\alpha S$ random field is invariant under the (left) translation action on its indices.

This invariant G -action, when seen at the level of spectral representations, gives rise to a *quasi-invariant* action.

Quasi-invariant group actions

Suppose $G \curvearrowright (\Lambda, \mathcal{A}, \mu)$ *measurably*,

Quasi-invariant group actions

Suppose $G \curvearrowright (\Lambda, \mathcal{A}, \mu)$ *measurably*, i.e., for all $g \in G$, the map $\phi_g : \Lambda \rightarrow \Lambda$ defined by

$$\phi_g : x \mapsto g^{-1} \cdot x$$

is measurable.

Quasi-invariant group actions

Suppose $G \curvearrowright (\Lambda, \mathcal{A}, \mu)$ *measurably*, i.e., for all $g \in G$, the map $\phi_g : \Lambda \rightarrow \Lambda$ defined by

$$\phi_g : x \mapsto g^{-1} \cdot x$$

is measurable.

Recall: This G -action is called *invariant* (or *measure-preserving*) if for all $g \in G$ and for all $A \in \mathcal{A}$,

$$g_*\mu(A) := \mu \circ \phi_g^{-1}(A) = \mu(A).$$

Quasi-invariant group actions

Suppose $G \curvearrowright (\Lambda, \mathcal{A}, \mu)$ *measurably*, i.e., for all $g \in G$, the map $\phi_g : \Lambda \rightarrow \Lambda$ defined by

$$\phi_g : x \mapsto g^{-1} \cdot x$$

is measurable.

Recall: This G -action is called *invariant* (or *measure-preserving*) if for all $g \in G$ and for all $A \in \mathcal{A}$,

$$g_*\mu(A) := \mu \circ \phi_g^{-1}(A) = \mu(A).$$

Definition: This G -action is called *quasi-invariant* (or *nonsingular*) if for all $g \in G$ and for all $A \in \mathcal{A}$,

$$g_*\mu(A) = 0 \iff \mu(A) = 0.$$

Quasi-invariant group actions

Suppose $G \curvearrowright (\Lambda, \mathcal{A}, \mu)$ *measurably*, i.e., for all $g \in G$, the map $\phi_g : \Lambda \rightarrow \Lambda$ defined by

$$\phi_g : x \mapsto g^{-1} \cdot x$$

is measurable.

Recall: This G -action is called *invariant* (or *measure-preserving*) if for all $g \in G$ and for all $A \in \mathcal{A}$,

$$g_*\mu(A) := \mu \circ \phi_g^{-1}(A) = \mu(A).$$

Definition: This G -action is called *quasi-invariant* (or *nonsingular*) if for all $g \in G$ and for all $A \in \mathcal{A}$,

$$g_*\mu(A) = 0 \iff \mu(A) = 0.$$

Clearly, *invariant* \implies *quasi-invariant* but the converse is not true.

Rosinski (1995, 2000) representation

Suppose $\{X_g\}_{g \in G}$ is a stationary $S\alpha S$ random field.

Rosinski (1995, 2000) representation

Suppose $\{X_g\}_{g \in G}$ is a stationary $S\alpha S$ random field.

Then $\{X_g\}_{g \in G}$ admits a representation of the form

$$f_g = \pm(f_e \circ \phi_g)$$

Rosinski (1995, 2000) representation

Suppose $\{X_g\}_{g \in G}$ is a stationary $S\alpha S$ random field.

Then $\{X_g\}_{g \in G}$ admits a representation of the form

$$f_g = \pm (f_e \circ \phi_g) \left(\frac{d\mu \circ \phi_g}{d\mu} \right)^{1/\alpha}, \quad g \in G \quad (2)$$

for some quasi-invariant G -action $\{\phi_g\}_{g \in G}$ on a σ -finite standard measure space $(\Lambda, \mathcal{A}, \mu)$.

Rosinski (1995, 2000) representation

Suppose $\{X_g\}_{g \in G}$ is a stationary $S\alpha S$ random field.

Then $\{X_g\}_{g \in G}$ admits a representation of the form

$$f_g = \pm (f_e \circ \phi_g) \left(\frac{d\mu \circ \phi_g}{d\mu} \right)^{1/\alpha}, \quad g \in G \quad (2)$$

for some quasi-invariant G -action $\{\phi_g\}_{g \in G}$ on a σ -finite standard measure space $(\Lambda, \mathcal{A}, \mu)$.

The **converse** also holds,

Rosinski (1995, 2000) representation

Suppose $\{X_g\}_{g \in G}$ is a stationary $S\alpha S$ random field.

Then $\{X_g\}_{g \in G}$ admits a representation of the form

$$f_g = \pm(f_e \circ \phi_g) \left(\frac{d\mu \circ \phi_g}{d\mu} \right)^{1/\alpha}, \quad g \in G \quad (2)$$

for some quasi-invariant G -action $\{\phi_g\}_{g \in G}$ on a σ -finite standard measure space $(\Lambda, \mathcal{A}, \mu)$.

The converse also holds, i.e., for any quasi-invariant G -action $\{\phi_g\}_{g \in G}$ on a σ -finite standard measure space $(\Lambda, \mathcal{A}, \mu)$ and any $f_e \in L^\alpha(\Lambda, \mu)$, if we define $\{f_g\}_{g \in G}$ by (2) ($\implies \{f_g\}_{g \in G} \subset L^\alpha(\Lambda, \mu)$), then there exists a stationary $S\alpha S$ random field $\{X_g\}_{g \in G}$ with representation $\{f_g\}_{g \in G}$, i.e., (1) holds.

Beyond Rosinski (1995, 2000) ...

Various probabilistic facets of a stable random field have been connected to the ergodic theoretic properties of the underlying nonsingular action:

Beyond Rosinski (1995, 2000) ...

Various probabilistic facets of a stable random field have been connected to the ergodic theoretic properties of the underlying nonsingular action:

- Mixing features: Rosinski and Samorodnitsky (1996), Samorodnitsky (2005), Roy (2007, 2012), Wang, R. and Stoev (2013)[AoP], R. (2020+)
- Large deviations issues: Mikosch and Samorodnitsky (2000), Fasen and R. (2016)[SPA]
- Growth of maxima: Samorodnitsky (2004), R. and Samorodnitsky (2008)[JTP], Chakrabarty and R. (2013)[JTP], Owada and Samorodnitsky (2015), Athreya, Mj and R. (2022+)[PTRF]
- Extremal point processes: Resnick and Samorodnitsky (2004), R. (2010)[AoP], Sarkar and R. (2018)[AoP]
- Statistical aspects: Bhattacharya and R. (2018)[JAP]
- Uniform Hölder continuity of paths: Panigrahi, R. and Xiao (2021)[SPA]

Beyond Rosinski (1995, 2000) ...

Various probabilistic facets of a stable random field have been connected to the ergodic theoretic properties of the underlying nonsingular action:

- Mixing features: Rosinski and Samorodnitsky (1996), Samorodnitsky (2005), Roy (2007, 2012), Wang, R. and Stoev (2013)[AoP], R. (2020+)
- Large deviations issues: Mikosch and Samorodnitsky (2000), Fasen and R. (2016)[SPA]
- Growth of maxima: Samorodnitsky (2004), R. and Samorodnitsky (2008)[JTP], Chakrabarty and R. (2013)[JTP], Owada and Samorodnitsky (2015), Athreya, Mj and R. (2022+)[PTRF]
- Extremal point processes: Resnick and Samorodnitsky (2004), R. (2010)[AoP], Sarkar and R. (2018)[AoP]
- Statistical aspects: Bhattacharya and R. (2018)[JAP]
- Uniform Hölder continuity of paths: Panigrahi, R. and Xiao (2021)[SPA]

Broad Goal: Carry this link forward to the realms of *Geometric Group Theory* (amenable groups, hyperbolic groups, etc.) and *Operator Algebra* (von Neumann algebras).

**Stationary Symmetric Stable
Random Fields Indexed by G**



Quasi-invariant G -Actions

Weakly Mixing

**Stationary Symmetric Stable
Random Fields Indexed by G**



Quasi-invariant G -Actions

Weak mixing and Rosinski representation

Weakly Mixing



????

**Stationary Symmetric Stable
Random Fields Indexed by G**



Quasi-invariant G -Actions

Weak mixing and Rosinski representation

Weakly Mixing

**Stationary Symmetric Stable
Random Fields Indexed by G**



Null

When G is amenable



Quasi-invariant G -Actions

Weak mixing and Rosinski representation

Weakly Mixing

**Stationary Symmetric Stable
Random Fields Indexed by G**



Null

When G is **amenable**



Quasi-invariant G -Actions

Questions or Comments or Concerns?

Weak Mixing for Random Fields

The induced probability measure on \mathbb{R}^G

Recall that any (left) stationary SoS random field $\mathbf{X} = \{X_g\}_{g \in G}$ induces a measure-preserving (left) shift action (of G) on $(\mathbb{R}^G, \mathbb{P}_{\mathbf{X}})$, where

$$\mathbb{P}_{\mathbf{X}} = \text{law of } \mathbf{X} := \mathbb{P}\left(\{\omega \in \Omega : (X_g(\omega) : g \in G) \in \cdot\}\right).$$

The induced probability measure on \mathbb{R}^G

Recall that any (left) stationary S α S random field $\mathbf{X} = \{X_g\}_{g \in G}$ induces a measure-preserving (left) shift action (of G) on $(\mathbb{R}^G, \mathbb{P}_{\mathbf{X}})$, where

$$\mathbb{P}_{\mathbf{X}} = \text{law of } \mathbf{X} := \mathbb{P}\left(\{\omega \in \Omega : (X_g(\omega) : g \in G) \in \cdot\}\right).$$

Recall that $\mathbb{R}^G := \{x : x \text{ is a map from } G \text{ to } \mathbb{R}\}$

The induced probability measure on \mathbb{R}^G

Recall that any (left) stationary S α S random field $\mathbf{X} = \{X_g\}_{g \in G}$ induces a measure-preserving (left) shift action (of G) on $(\mathbb{R}^G, \mathbb{P}_{\mathbf{X}})$, where

$$\mathbb{P}_{\mathbf{X}} = \text{law of } \mathbf{X} := \mathbb{P}\left(\left\{\omega \in \Omega : (X_g(\omega) : g \in G) \in \cdot\right\}\right).$$

Recall that $\mathbb{R}^G := \{x : x \text{ is a map from } G \text{ to } \mathbb{R}\}$ and a prototypical Borel cylinder subset of \mathbb{R}^G looks like

$$C = \left\{x \in \mathbb{R}^G : x(g_1) \in B_1, x(g_1) \in B_2, \dots, x(g_k) \in B_k\right\}$$

for some $k \in \mathbb{N}$, for some $g_1, g_2, \dots, g_k \in G$ and for some Borel sets $B_1, B_2, \dots, B_k \subseteq \mathbb{R}$.

The induced probability measure on \mathbb{R}^G

Recall that any (left) stationary SoS random field $\mathbf{X} = \{X_g\}_{g \in G}$ induces a measure-preserving (left) shift action (of G) on $(\mathbb{R}^G, \mathbb{P}_{\mathbf{X}})$, where

$$\mathbb{P}_{\mathbf{X}} = \text{law of } \mathbf{X} := \mathbb{P}\left(\{\omega \in \Omega : (X_g(\omega) : g \in G) \in \cdot\}\right).$$

Recall that $\mathbb{R}^G := \{x : x \text{ is a map from } G \text{ to } \mathbb{R}\}$ and a prototypical Borel cylinder subset of \mathbb{R}^G looks like

$$C = \left\{x \in \mathbb{R}^G : x(g_1) \in B_1, x(g_1) \in B_2, \dots, x(g_k) \in B_k\right\}$$

for some $k \in \mathbb{N}$, for some $g_1, g_2, \dots, g_k \in G$ and for some Borel sets $B_1, B_2, \dots, B_k \subseteq \mathbb{R}$.

Clearly

$$\mathbb{P}_{\mathbf{X}}(C) = \mathbb{P}\left(\{\omega \in \Omega : (X_g(\omega) : g \in G) \in C\}\right)$$

The induced probability measure on \mathbb{R}^G

Recall that any (left) stationary S α S random field $\mathbf{X} = \{X_g\}_{g \in G}$ induces a measure-preserving (left) shift action (of G) on $(\mathbb{R}^G, \mathbb{P}_{\mathbf{X}})$, where

$$\mathbb{P}_{\mathbf{X}} = \text{law of } \mathbf{X} := \mathbb{P}\left(\{\omega \in \Omega : (X_g(\omega) : g \in G) \in \cdot\}\right).$$

Recall that $\mathbb{R}^G := \{x : x \text{ is a map from } G \text{ to } \mathbb{R}\}$ and a prototypical Borel cylinder subset of \mathbb{R}^G looks like

$$C = \left\{x \in \mathbb{R}^G : x(g_1) \in B_1, x(g_2) \in B_2, \dots, x(g_k) \in B_k\right\}$$

for some $k \in \mathbb{N}$, for some $g_1, g_2, \dots, g_k \in G$ and for some Borel sets $B_1, B_2, \dots, B_k \subseteq \mathbb{R}$.

Clearly

$$\begin{aligned}\mathbb{P}_{\mathbf{X}}(C) &= \mathbb{P}\left(\{\omega \in \Omega : (X_g(\omega) : g \in G) \in C\}\right) \\ &= \mathbb{P}\left(\{\omega \in \Omega : X_{g_1}(\omega) \in B_1, X_{g_2}(\omega) \in B_2, \dots, X_{g_k}(\omega) \in B_k\}\right)\end{aligned}$$

The induced probability measure on \mathbb{R}^G

Recall that any (left) stationary SoS random field $\mathbf{X} = \{X_g\}_{g \in G}$ induces a measure-preserving (left) shift action (of G) on $(\mathbb{R}^G, \mathbb{P}_{\mathbf{X}})$, where

$$\mathbb{P}_{\mathbf{X}} = \text{law of } \mathbf{X} := \mathbb{P}\left(\{\omega \in \Omega : (X_g(\omega) : g \in G) \in \cdot\}\right).$$

Recall that $\mathbb{R}^G := \{x : x \text{ is a map from } G \text{ to } \mathbb{R}\}$ and a prototypical Borel cylinder subset of \mathbb{R}^G looks like

$$C = \left\{x \in \mathbb{R}^G : x(g_1) \in B_1, x(g_1) \in B_2, \dots, x(g_k) \in B_k\right\}$$

for some $k \in \mathbb{N}$, for some $g_1, g_2, \dots, g_k \in G$ and for some Borel sets $B_1, B_2, \dots, B_k \subseteq \mathbb{R}$.

Clearly

$$\begin{aligned}\mathbb{P}_{\mathbf{X}}(C) &= \mathbb{P}\left(\{\omega \in \Omega : (X_g(\omega) : g \in G) \in C\}\right) \\ &= \mathbb{P}\left(\{\omega \in \Omega : X_{g_1}(\omega) \in B_1, X_{g_2}(\omega) \in B_2, \dots, X_{g_k}(\omega) \in B_k\}\right) \\ &= \mathbb{P}(X_{g_1} \in B_1, X_{g_2} \in B_2, \dots, X_{g_k} \in B_k).\end{aligned}$$

What does weak mixing mean?

Recall that any (left) stationary S α S random field $\mathbf{X} = \{X_g\}_{g \in G}$ induces a measure-preserving (left) shift action (of G) on $(\mathbb{R}^G, \mathbb{P}_{\mathbf{X}} := \text{law of } \mathbf{X})$.

What does weak mixing mean?

Recall that any (left) stationary S α S random field $\mathbf{X} = \{X_g\}_{g \in G}$ induces a measure-preserving (left) shift action (of G) on $(\mathbb{R}^G, \mathbb{P}_{\mathbf{X}} := \text{law of } \mathbf{X})$.

Definition

$\{X_g\}_{g \in G}$ is called weakly mixing if the above shift action is so.

What does weak mixing mean?

Recall that any (left) stationary SoS random field $\mathbf{X} = \{X_g\}_{g \in G}$ induces a measure-preserving (left) shift action (of G) on $(\mathbb{R}^G, \mathbb{P}_{\mathbf{X}} := \text{law of } \mathbf{X})$.

Definition

$\{X_g\}_{g \in G}$ is called weakly mixing if the above shift action is so.

Question: Why is it important?

What does weak mixing mean?

Recall that any (left) stationary SoS random field $\mathbf{X} = \{X_g\}_{g \in G}$ induces a measure-preserving (left) shift action (of G) on $(\mathbb{R}^G, \mathbb{P}_{\mathbf{X}} := \text{law of } \mathbf{X})$.

Definition

$\{X_g\}_{g \in G}$ is called *weakly mixing* if the above shift action is so.

Question: Why is it important?

- Roughly speaking, it's some kind of asymptotic independence: weak mixing of $\{X_t\}_{t \in \mathbb{Z}}$ implies that for all Borel $A, B \subseteq \mathbb{R}$,

$$\frac{1}{n} \sum_{i=1}^n |\mathbb{P}(X_0 \in A, X_i \in B) - \mathbb{P}(X_0 \in A)\mathbb{P}(X_0 \in B)| \rightarrow 0.$$

What does weak mixing mean?

Recall that any (left) stationary SoS random field $\mathbf{X} = \{X_g\}_{g \in G}$ induces a measure-preserving (left) shift action (of G) on $(\mathbb{R}^G, \mathbb{P}_{\mathbf{X}} := \text{law of } \mathbf{X})$.

Definition

$\{X_g\}_{g \in G}$ is called *weakly mixing* if the above shift action is so.

Question: Why is it important?

- Roughly speaking, it's some kind of asymptotic independence: weak mixing of $\{X_t\}_{t \in \mathbb{Z}}$ implies that for all Borel $A, B \subseteq \mathbb{R}$,

$$\frac{1}{n} \sum_{i=1}^n |\mathbb{P}(X_0 \in A, X_i \in B) - \mathbb{P}(X_0 \in A)\mathbb{P}(X_0 \in B)| \rightarrow 0.$$

- When the underlying random field is weakly mixing, then we can apply ergodic theorem to investigate asymptotic properties of estimators/algorithms, large deviation issues, long run behaviour of solutions to SDEs, etc.

Weak mixing of G -indexed stable fields

Recall that any (left) stationary S α S random field $\mathbf{X} = \{X_g\}_{g \in G}$ induces a measure-preserving (left) shift action (of G) on $(\mathbb{R}^G, \mathbb{P}_{\mathbf{X}})$, where

$$\mathbb{P}_{\mathbf{X}} = \text{law of } \mathbf{X} := \mathbb{P}\left(\{\omega \in \Omega : (X_g(\omega) : g \in G) \in \cdot\}\right).$$

Weak mixing of G -indexed stable fields

Recall that any (left) stationary S α S random field $\mathbf{X} = \{X_g\}_{g \in G}$ induces a measure-preserving (left) shift action (of G) on $(\mathbb{R}^G, \mathbb{P}_{\mathbf{X}})$, where

$$\mathbb{P}_{\mathbf{X}} = \text{law of } \mathbf{X} := \mathbb{P}\left(\{\omega \in \Omega : (X_g(\omega) : g \in G) \in \cdot\}\right).$$

Definition

$\{X_g\}_{g \in G}$ is called *weak mixing* if the above shift action is so.

Weak mixing of G -indexed stable fields

Recall that any (left) stationary S α S random field $\mathbf{X} = \{X_g\}_{g \in G}$ induces a measure-preserving (left) shift action (of G) on $(\mathbb{R}^G, \mathbb{P}_{\mathbf{X}})$, where

$$\mathbb{P}_{\mathbf{X}} = \text{law of } \mathbf{X} := \mathbb{P}\left(\{\omega \in \Omega : (X_g(\omega) : g \in G) \in \cdot\}\right).$$

Definition

$\{X_g\}_{g \in G}$ is called *weak mixing* if the above shift action is so.

Question: When is $\{X_g\}_{g \in G}$ weak mixing?

Weak mixing of G -indexed stable fields

Recall that any (left) stationary S α S random field $\mathbf{X} = \{X_g\}_{g \in G}$ induces a measure-preserving (left) shift action (of G) on $(\mathbb{R}^G, \mathbb{P}_{\mathbf{X}})$, where

$$\mathbb{P}_{\mathbf{X}} = \text{law of } \mathbf{X} := \mathbb{P}\left(\{\omega \in \Omega : (X_g(\omega) : g \in G) \in \cdot\}\right).$$

Definition

$\{X_g\}_{g \in G}$ is called *weak mixing* if the above shift action is so.

Question: When is $\{X_g\}_{g \in G}$ weak mixing? [\[Helps in proving limit theorems.\]](#)

Weak mixing of G -indexed stable fields

Recall that any (left) stationary S α S random field $\mathbf{X} = \{X_g\}_{g \in G}$ induces a measure-preserving (left) shift action (of G) on $(\mathbb{R}^G, \mathbb{P}_{\mathbf{X}})$, where

$$\mathbb{P}_{\mathbf{X}} = \text{law of } \mathbf{X} := \mathbb{P}\left(\{\omega \in \Omega : (X_g(\omega) : g \in G) \in \cdot\}\right).$$

Definition

$\{X_g\}_{g \in G}$ is called *weak mixing* if the above shift action is so.

Question: When is $\{X_g\}_{g \in G}$ weak mixing? [\[Helps in proving limit theorems.\]](#)

- $G = \mathbb{Z}$: [Samorodnitsky \(2005\)](#) gave a criterion based on the ergodic theoretic properties of the underlying quasi-invariant action.

Weak mixing of G -indexed stable fields

Recall that any (left) stationary S α S random field $\mathbf{X} = \{X_g\}_{g \in G}$ induces a measure-preserving (left) shift action (of G) on $(\mathbb{R}^G, \mathbb{P}_{\mathbf{X}})$, where

$$\mathbb{P}_{\mathbf{X}} = \text{law of } \mathbf{X} := \mathbb{P}\left(\{\omega \in \Omega : (X_g(\omega) : g \in G) \in \cdot\}\right).$$

Definition

$\{X_g\}_{g \in G}$ is called *weak mixing* if the above shift action is so.

Question: When is $\{X_g\}_{g \in G}$ weak mixing? [\[Helps in proving limit theorems.\]](#)

- $G = \mathbb{Z}$: [Samorodnitsky \(2005\)](#) gave a criterion based on the ergodic theoretic properties of the underlying quasi-invariant action.
- $G = \mathbb{Z}^d$: [Wang, R. and Stoev \(2013\)](#) generalized the above result (for $d = 1$) to any $d \in \mathbb{N}$ using the work of [Takahashi \(1971\)](#).

Stable Random Fields Indexed by Amenable Groups

Weak mixing and Rosinski representation

Weakly Mixing

**Stationary Symmetric Stable
Random Fields Indexed by G**



Null

When G is **amenable**



Quasi-invariant G -Actions

Countable amenable groups

A countable amenable group is a countable group that admits an increasing Følner sequence $F_n \uparrow G$, i.e., an increasing sequence of exhausting finite subsets $F_n \subset G$ such that for all $g \in G$,

$$\lim_{n \rightarrow \infty} \frac{|gF_n \Delta F_n|}{|F_n|} = 0.$$

Countable amenable groups

A countable amenable group is a countable group that admits an increasing Følner sequence $F_n \uparrow G$, i.e., an increasing sequence of exhausting finite subsets $F_n \subset G$ such that for all $g \in G$,

$$\lim_{n \rightarrow \infty} \frac{|gF_n \Delta F_n|}{|F_n|} = 0.$$

The above limiting condition is called the Følner condition, which roughly means that for large n , very small fraction of elements in F_n are moved by a group element.

Countable amenable groups

A countable amenable group is a countable group that admits an increasing Følner sequence $F_n \uparrow G$, i.e., an increasing sequence of exhausting finite subsets $F_n \subset G$ such that for all $g \in G$,

$$\lim_{n \rightarrow \infty} \frac{|gF_n \Delta F_n|}{|F_n|} = 0.$$

The above limiting condition is called the Følner condition, which roughly means that for large n , very small fraction of elements in F_n are moved by a group element.

Examples: finite groups, abelian groups, groups of subexponential growth, solvable groups, lamplighter groups, etc.

Countable amenable groups

A countable amenable group is a countable group that admits an increasing Følner sequence $F_n \uparrow G$, i.e., an increasing sequence of exhausting finite subsets $F_n \subset G$ such that for all $g \in G$,

$$\lim_{n \rightarrow \infty} \frac{|gF_n \Delta F_n|}{|F_n|} = 0.$$

The above limiting condition is called the Følner condition, which roughly means that for large n , very small fraction of elements in F_n are moved by a group element.

Examples: finite groups, abelian groups, groups of subexponential growth, solvable groups, lamplighter groups, etc.

Lindenstrauss (2001):

- 1 Any Følner sequence in an amenable group G admits a “tempered Følner subsequence”.

Countable amenable groups

A countable amenable group is a countable group that admits an increasing Følner sequence $F_n \uparrow G$, i.e., an increasing sequence of exhausting finite subsets $F_n \subset G$ such that for all $g \in G$,

$$\lim_{n \rightarrow \infty} \frac{|gF_n \Delta F_n|}{|F_n|} = 0.$$

The above limiting condition is called the Følner condition, which roughly means that for large n , very small fraction of elements in F_n are moved by a group element.

Examples: finite groups, abelian groups, groups of subexponential growth, solvable groups, lamplighter groups, etc.

Lindenstrauss (2001):

- ① Any Følner sequence in an amenable group G admits a “tempered Følner subsequence”.
- ② Along any tempered Følner sequence, pointwise ergodic theorem holds for any finite measure preserving G -action.

Why stable random fields indexed by amenable groups?

The reason is two-fold:

Why stable random fields indexed by amenable groups?

The reason is two-fold:

- 1 Statistical:

Why stable random fields indexed by amenable groups?

The reason is two-fold:

1 Statistical:

- ▶ Such random fields arise naturally in **machine learning algorithms** for structured and dependent data; see, e.g., [Austern and Orbanz \(2022+\)](#).

Why stable random fields indexed by amenable groups?

The reason is two-fold:

1 Statistical:

- ▶ Such random fields arise naturally in **machine learning algorithms** for structured and dependent data; see, e.g., [Austern and Orbanz \(2022+\)](#).
- ▶ Weak mixing is an important tool for investigating asymptotic properties of any method/algorithm. In the context of **space-time statistical inference** for max-stable fields, see [Davis, Klüppelberg and Steinkohl \(2013\)](#).

Why stable random fields indexed by amenable groups?

The reason is two-fold:

1 Statistical:

- ▶ Such random fields arise naturally in **machine learning algorithms** for structured and dependent data; see, e.g., [Austern and Orbanz \(2022+\)](#).
- ▶ Weak mixing is an important tool for investigating asymptotic properties of any method/algorithm. In the context of **space-time statistical inference** for max-stable fields, see [Davis, Klüppelberg and Steinkohl \(2013\)](#).

2 Mathematical:

Why stable random fields indexed by amenable groups?

The reason is two-fold:

1 Statistical:

- ▶ Such random fields arise naturally in **machine learning algorithms** for structured and dependent data; see, e.g., [Austern and Orbanz \(2022+\)](#).
- ▶ Weak mixing is an important tool for investigating asymptotic properties of any method/algorithm. In the context of **space-time statistical inference** for max-stable fields, see [Davis, Klüppelberg and Steinkohl \(2013\)](#).

2 Mathematical:

- ▶ How about stable random fields indexed by groups of polynomial growth? Even the seminal work of [Gromov \(1981\)](#) could not help :-)

Why stable random fields indexed by amenable groups?

The reason is two-fold:

1 Statistical:

- ▶ Such random fields arise naturally in **machine learning algorithms** for structured and dependent data; see, e.g., [Austern and Orbanz \(2022+\)](#).
- ▶ Weak mixing is an important tool for investigating asymptotic properties of any method/algorithm. In the context of **space-time statistical inference** for max-stable fields, see [Davis, Klüppelberg and Steinkohl \(2013\)](#).

2 Mathematical:

- ▶ How about stable random fields indexed by groups of polynomial growth? Even the seminal work of [Gromov \(1981\)](#) could not help :-)
- ▶ [Mj, R. and Sarkar \(2022+\)](#): applied a probabilistic+analytic truncation argument + [Lindenstrauss \(2001\)](#) + [Tempelman \(2015\)](#)'s ergodic theorem for Lamperti representations.

Why stable random fields indexed by amenable groups?

The reason is two-fold:

1 Statistical:

- ▶ Such random fields arise naturally in **machine learning algorithms** for structured and dependent data; see, e.g., [Austern and Orbanz \(2022+\)](#).
- ▶ Weak mixing is an important tool for investigating asymptotic properties of any method/algorithm. In the context of **space-time statistical inference** for max-stable fields, see [Davis, Klüppelberg and Steinkohl \(2013\)](#).

2 Mathematical:

- ▶ How about stable random fields indexed by groups of polynomial growth? Even the seminal work of [Gromov \(1981\)](#) could not help :-)
- ▶ [Mj, R. and Sarkar \(2022+\)](#): applied a probabilistic+analytic truncation argument + [Lindenstrauss \(2001\)](#) + [Tempelman \(2015\)](#)'s ergodic theorem for Lamperti representations.
- ▶ [Avraham Re'em \(2022+\)](#) [[arXived a few days ago!](#)]: used the machinery of absolutely continuous invariant probability measure + a Krengel-type criterion + [Lindenstrauss \(2001\)](#).

Questions or Comments or Concerns?

Recall: Rosinski (1995, 2000) representation

Given a stationary $S\alpha S$ ($0 < \alpha < 2$) random field $\{X_g\}_{g \in G}$, there exist

- (i) a σ -finite standard measure space $(\Lambda, \mathcal{A}, \mu)$,
- (ii) a function $f_e : \Lambda \rightarrow \mathbb{R}$ such that $\|f_e\|_\alpha := \left(\int |f_e|^\alpha d\mu \right)^{1/\alpha} < \infty$, and
- (iii) a quasi-invariant G -action $\{\phi_g\}_{g \in G}$ on $(\Lambda, \mathcal{A}, \mu)$

such that each real linear combination

$$\sum_{i=1}^k c_i X_{g_i} \sim S\alpha S \left(\left\| \sum_{i=1}^k c_i f_{g_i} \right\|_\alpha \right), \quad (3)$$

where

$$f_g = \pm \left(\frac{d\mu \circ \phi_g}{d\mu} \right)^{1/\alpha} f_e \circ \phi_g, \quad g \in G.$$

Converse also holds: given (i), (ii) and (iii), there exists a stationary $S\alpha S$ random field $\{X_g\}_{g \in G}$ satisfying (3).

$\{f_g\}_{g \in G} = \mathbf{a}$ Rosinski representation of $\{X_g\}_{g \in G}$.

Weak mixing and Rosinski representation

Weakly Mixing

**Stationary Symmetric Stable
Random Fields Indexed by G**



Null

When G is amenable



Quasi-invariant G -Actions

Null quasi-invariant actions

Let $\{\phi_g\}_{g \in G}$ be a quasi-invariant action on $(\Lambda, \mathcal{A}, \mu)$.

Null quasi-invariant actions

Let $\{\phi_g\}_{g \in G}$ be a quasi-invariant action on $(\Lambda, \mathcal{A}, \mu)$.

$W \in \mathcal{A}$ is called *weakly wandering* if there exists a countably infinite set $L \subseteq G$ (not necessarily a subgroup) such that $\{\phi_g(W) : g \in L\}$ is a pairwise disjoint collection.

Null quasi-invariant actions

Let $\{\phi_g\}_{g \in G}$ be a quasi-invariant action on $(\Lambda, \mathcal{A}, \mu)$.

$W \in \mathcal{A}$ is called *weakly wandering* if there exists a countably infinite set $L \subseteq G$ (not necessarily a subgroup) such that $\{\phi_g(W) : g \in L\}$ is a pairwise disjoint collection.

Roughly speaking, elements of a weakly wandering set do not come back to itself very often by the shift of this action.

Null quasi-invariant actions

Let $\{\phi_g\}_{g \in G}$ be a quasi-invariant action on $(\Lambda, \mathcal{A}, \mu)$.

$W \in \mathcal{A}$ is called *weakly wandering* if there exists a countably infinite set $L \subseteq G$ (not necessarily a subgroup) such that $\{\phi_g(W) : g \in L\}$ is a pairwise disjoint collection.

Roughly speaking, elements of a weakly wandering set do not come back to itself very often by the shift of this action.

“Definition”

The action $\{\phi_g\}_{g \in G}$ is called null if Λ is the “measurable union” of all of its weakly wandering subsets.

Null quasi-invariant actions

Let $\{\phi_g\}_{g \in G}$ be a quasi-invariant action on $(\Lambda, \mathcal{A}, \mu)$.

$W \in \mathcal{A}$ is called *weakly wandering* if there exists a countably infinite set $L \subseteq G$ (not necessarily a subgroup) such that $\{\phi_g(W) : g \in L\}$ is a pairwise disjoint collection.

Roughly speaking, elements of a weakly wandering set do not come back to itself very often by the shift of this action.

“Definition”

The action $\{\phi_g\}_{g \in G}$ is called null if Λ is the “measurable union” of all of its weakly wandering subsets.

Intuitively speaking, a null action does not come back very often.

Stable random fields generated by null actions tend to have **weaker dependence** because ...

Stable random fields generated by null actions tend to have **weaker dependence** because ...

a null action does not come back very often, and hence X_g 's tend to become “asymptotically independent” (of X_e , say) as g moves far and far away (in some suitable metric) from the identity element e .

Stable random fields generated by null actions tend to have **weaker dependence** because ...

a null action does not come back very often, and hence X_g 's tend to become “asymptotically independent” (of X_e , say) as g moves far and far away (in some suitable metric) from the identity element e .

This *weaker dependence* / *asymptotic independence* may manifest itself in the form of weak mixing.

Stable random fields generated by null actions tend to have **weaker dependence** because ...

a null action does not come back very often, and hence X_g 's tend to become “asymptotically independent” (of X_e , say) as g moves far and far away (in some suitable metric) from the identity element e .

This *weaker dependence / asymptotic independence* may manifest itself in the form of weak mixing.

We have been able to prove this formally only when G is amenable. We believe that our result may be true for a much bigger class of groups and/or actions.

Weakly mixing random fields indexed by amenable groups

Let G be a countably infinite amenable group and $\{F_n\}_{n \geq 1}$ is any **increasing** Følner sequence.

Weakly mixing random fields indexed by amenable groups

Let G be a countably infinite amenable group and $\{F_n\}_{n \geq 1}$ is any **increasing** Følner sequence.

Dye (1965) [Cor 1, pg 129] or Bergelson and Gorodnik (2004) [Thm 1.6]:
A random field $\{X_g\}_{g \in G}$ is weakly mixing if and only if for all $k \geq 1$, for all $g_1, g_2, \dots, g_k, h \in G$ and for all Borel $A, B \subset \mathbb{R}^k$,

$$\frac{1}{|F_n|} \sum_{h \in F_n} \left| P[(X_{hg_1}, X_{hg_2}, \dots, X_{hg_k}) \in A, (X_{g_1}, X_{g_2}, \dots, X_{g_k}) \in B] \right. \\ \left. - P[(X_{g_1}, X_{g_2}, \dots, X_{g_k}) \in A] P[(X_{g_1}, X_{g_2}, \dots, X_{g_k}) \in B] \right| \rightarrow 0$$

as $n \rightarrow \infty$.

Weakly mixing random fields indexed by amenable groups

Let G be a countably infinite amenable group and $\{F_n\}_{n \geq 1}$ is any **increasing** Følner sequence.

Dye (1965) [Cor 1, pg 129] or Bergelson and Gorodnik (2004) [Thm 1.6]:
A random field $\{X_g\}_{g \in G}$ is weakly mixing if and only if for all $k \geq 1$, for all $g_1, g_2, \dots, g_k, h \in G$ and for all Borel $A, B \subset \mathbb{R}^k$,

$$\frac{1}{|F_n|} \sum_{h \in F_n} \left| P[(X_{hg_1}, X_{hg_2}, \dots, X_{hg_k}) \in A, (X_{g_1}, X_{g_2}, \dots, X_{g_k}) \in B] \right. \\ \left. - P[(X_{g_1}, X_{g_2}, \dots, X_{g_k}) \in A] P[(X_{g_1}, X_{g_2}, \dots, X_{g_k}) \in B] \right| \rightarrow 0$$

as $n \rightarrow \infty$.

Choose the Følner sequence to be **increasing** and **tempered** according to Lindenstrauss (2001).

Weakly mixing random fields indexed by amenable groups

Let G be a countably infinite amenable group and $\{F_n\}_{n \geq 1}$ is any **increasing** Følner sequence.

Dye (1965) [**Cor 1, pg 129**] or **Bergelson and Gorodnik (2004)** [**Thm 1.6**]:
A random field $\{X_g\}_{g \in G}$ is weakly mixing if and only if for all $k \geq 1$, for all $g_1, g_2, \dots, g_k, h \in G$ and for all Borel $A, B \subset \mathbb{R}^k$,

$$\frac{1}{|F_n|} \sum_{h \in F_n} \left| P[(X_{hg_1}, X_{hg_2}, \dots, X_{hg_k}) \in A, (X_{g_1}, X_{g_2}, \dots, X_{g_k}) \in B] \right. \\ \left. - P[(X_{g_1}, X_{g_2}, \dots, X_{g_k}) \in A] P[(X_{g_1}, X_{g_2}, \dots, X_{g_k}) \in B] \right| \rightarrow 0$$

as $n \rightarrow \infty$.

Choose the Følner sequence to be **increasing** and **tempered** according to **Lindenstrauss (2001)**. These will give us **analytic** and **ergodic theoretic** advantages, respectively.

One of our main results

Theorem (Mj, R. and Sarkar (2022+))

Suppose G is a countably infinite amenable group and $\mathbf{X} := \{X_g\}_{g \in G}$ is a left stationary symmetric α -stable ($0 < \alpha < 2$) random field generated by a quasi-invariant G -action $\{\phi_g\}_{g \in G}$ in its Rosinski representation. Then \mathbf{X} is weakly mixing if and only if $\{\phi_g\}_{g \in G}$ is a null action.

One of our main results

Theorem (Mj, R. and Sarkar (2022+))

Suppose G is a countably infinite amenable group and $\mathbf{X} := \{X_g\}_{g \in G}$ is a left stationary symmetric α -stable ($0 < \alpha < 2$) random field generated by a quasi-invariant G -action $\{\phi_g\}_{g \in G}$ in its Rosinski representation. Then \mathbf{X} is weakly mixing if and only if $\{\phi_g\}_{g \in G}$ is a null action.

Avraham Re'em (2022+) [arXived a few days ago!] has also proved this result using different techniques.

One of our main results

Theorem (Mj, R. and Sarkar (2022+))

Suppose G is a countably infinite amenable group and $\mathbf{X} := \{X_g\}_{g \in G}$ is a left stationary symmetric α -stable ($0 < \alpha < 2$) random field generated by a quasi-invariant G -action $\{\phi_g\}_{g \in G}$ in its Rosinski representation. Then \mathbf{X} is weakly mixing if and only if $\{\phi_g\}_{g \in G}$ is a null action.

Avraham Re'em (2022+) [arXived a few days ago!] has also proved this result using different techniques.

Examples:

One of our main results

Theorem (Mj, R. and Sarkar (2022+))

Suppose G is a countably infinite amenable group and $\mathbf{X} := \{X_g\}_{g \in G}$ is a left stationary symmetric α -stable ($0 < \alpha < 2$) random field generated by a quasi-invariant G -action $\{\phi_g\}_{g \in G}$ in its Rosinski representation. Then \mathbf{X} is weakly mixing if and only if $\{\phi_g\}_{g \in G}$ is a null action.

Avraham Re'em (2022+) [arXived a few days ago!] has also proved this result using different techniques.

Examples:

- Many canonical actions of discrete Heisenberg groups are null.

One of our main results

Theorem (Mj, R. and Sarkar (2022+))

Suppose G is a countably infinite amenable group and $\mathbf{X} := \{X_g\}_{g \in G}$ is a left stationary symmetric α -stable ($0 < \alpha < 2$) random field generated by a quasi-invariant G -action $\{\phi_g\}_{g \in G}$ in its Rosinski representation. Then \mathbf{X} is weakly mixing if and only if $\{\phi_g\}_{g \in G}$ is a null action.

Avraham Re'em (2022+) [arXived a few days ago!] has also proved this result using different techniques.

Examples:

- Many canonical actions of discrete Heisenberg groups are null.
- Whenever an amenable group G has a nontrivial Furstenberg-Poisson boundary (e.g., this is the case for lamplighter groups on \mathbb{Z}^d with $d \geq 3$ thanks to the seminal work of Kaimanovich and Vershik (1983)), the boundary action is nontrivial and null.

One of our main results

Theorem (Mj, R. and Sarkar (2022+))

Suppose G is a countably infinite amenable group and $\mathbf{X} := \{X_g\}_{g \in G}$ is a left stationary symmetric α -stable ($0 < \alpha < 2$) random field generated by a quasi-invariant G -action $\{\phi_g\}_{g \in G}$ in its Rosinski representation. Then \mathbf{X} is weakly mixing if and only if $\{\phi_g\}_{g \in G}$ is a null action.

Avraham Re'em (2022+) [arXived a few days ago!] has also proved this result using different techniques.

Examples:

- Many canonical actions of discrete Heisenberg groups are null.
- Whenever an amenable group G has a nontrivial Furstenberg-Poisson boundary (e.g., this is the case for lamplighter groups on \mathbb{Z}^d with $d \geq 3$ thanks to the seminal work of Kaimanovich and Vershik (1983)), the boundary action is nontrivial and null.
- There are many solvable groups (e.g., suitable discrete subgroups of Lie groups) that admit many natural null actions.

How do we prove this result?

- **Main Obstacle:** Unavailability of pointwise ergodic theorem for quasi-invariant actions.

How do we prove this result?

- **Main Obstacle:** Unavailability of pointwise ergodic theorem for quasi-invariant actions.
- [Lindenstrauss \(2001\)](#) proved pointwise ergodic theorem for **finite measure preserving** actions.

How do we prove this result?

- **Main Obstacle:** Unavailability of pointwise ergodic theorem for quasi-invariant actions.
- Lindenstrauss (2001) proved pointwise ergodic theorem for **finite measure preserving** actions.
- Tempelman (2015) observed that the maximal ergodic theorem established by Lindenstrauss (2001) actually holds for any σ -finite measure preserving action,

How do we prove this result?

- **Main Obstacle:** Unavailability of pointwise ergodic theorem for quasi-invariant actions.
- Lindenstrauss (2001) proved pointwise ergodic theorem for finite measure preserving actions.
- Tempelman (2015) observed that the maximal ergodic theorem established by Lindenstrauss (2001) actually holds for any σ -finite measure preserving action, and used this observation to prove a pointwise ergodic theorem (along any tempered Følner sequence) for bounded Lamperti representations of an amenable group G into L^p -spaces for $p > 1$.

How do we prove this result?

- **Main Obstacle:** Unavailability of pointwise ergodic theorem for quasi-invariant actions.
- [Lindenstrauss \(2001\)](#) proved pointwise ergodic theorem for **finite measure preserving** actions.
- [Tempelman \(2015\)](#) observed that the maximal ergodic theorem established by [Lindenstrauss \(2001\)](#) actually holds for any σ -finite measure preserving action, and used this observation to prove a pointwise ergodic theorem (along any tempered Følner sequence) for bounded Lamperti representations of an amenable group G into L^p -spaces for $p > 1$.
- We apply this result + a truncation argument to establish our result for any amenable group G provided the underlying action is (σ -finite) measure preserving.

How do we prove this result?

- **Main Obstacle:** Unavailability of pointwise ergodic theorem for quasi-invariant actions.
- Lindenstrauss (2001) proved pointwise ergodic theorem for finite measure preserving actions.
- Tempelman (2015) observed that the maximal ergodic theorem established by Lindenstrauss (2001) actually holds for any σ -finite measure preserving action, and used this observation to prove a pointwise ergodic theorem (along any tempered Følner sequence) for bounded Lamperti representations of an amenable group G into L^p -spaces for $p > 1$.
- We apply this result + a truncation argument to establish our result for any amenable group G provided the underlying action is (σ -finite) measure preserving.
- Generalization from the measure preserving to the quasi-invariant case is carried out using Maharam (1964) skew-product.

Outline of proof

Need to show, due to an extension of [Gross \(1994\)](#),

Outline of proof

Need to show, due to an extension of [Gross \(1994\)](#), that for all $\delta > 0$ and $\epsilon > 0$,

$$\lim_{n \rightarrow \infty} \frac{1}{F_n} \sum_{g \in F_n} \mu(x \in \Lambda : \delta^{-1} > |f_e(x)| > \delta, |f_g(x)| > \epsilon) = 0, \quad (4)$$

where $f_g = \pm \left(\frac{d\mu \circ \phi_g}{d\mu} \right)^{1/\alpha} f_e \circ \phi_g$, $g \in G$ is a Rosinski representaion of \mathbf{X} .

Outline of proof

Need to show, due to an extension of [Gross \(1994\)](#), that for all $\delta > 0$ and $\epsilon > 0$,

$$\lim_{n \rightarrow \infty} \frac{1}{F_n} \sum_{g \in F_n} \mu(x \in \Lambda : \delta^{-1} > |f_e(x)| > \delta, |f_g(x)| > \epsilon) = 0, \quad (4)$$

where $f_g = \pm \left(\frac{d\mu \circ \phi_g}{d\mu} \right)^{1/\alpha} f_e \circ \phi_g$, $g \in G$ is a Rosinski representaion of \mathbf{X} .

A probabilistic+analytic truncation argument \Rightarrow enough to show (4) when

Outline of proof

Need to show, due to an extension of [Gross \(1994\)](#), that for all $\delta > 0$ and $\epsilon > 0$,

$$\lim_{n \rightarrow \infty} \frac{1}{F_n} \sum_{g \in F_n} \mu(x \in \Lambda : \delta^{-1} > |f_e(x)| > \delta, |f_g(x)| > \epsilon) = 0, \quad (4)$$

where $f_g = \pm \left(\frac{d\mu \circ \phi_g}{d\mu} \right)^{1/\alpha} f_e \circ \phi_g$, $g \in G$ is a Rosinski representaion of \mathbf{X} .

A probabilistic+analytic truncation argument \Rightarrow enough to show (4) when

1 μ is a finite measure,

Outline of proof

Need to show, due to an extension of [Gross \(1994\)](#), that for all $\delta > 0$ and $\epsilon > 0$,

$$\lim_{n \rightarrow \infty} \frac{1}{F_n} \sum_{g \in F_n} \mu(x \in \Lambda : \delta^{-1} > |f_e(x)| > \delta, |f_g(x)| > \epsilon) = 0, \quad (4)$$

where $f_g = \pm \left(\frac{d\mu \circ \phi_g}{d\mu} \right)^{1/\alpha} f_e \circ \phi_g$, $g \in G$ is a Rosinski representaion of \mathbf{X} .

A probabilistic+analytic truncation argument \Rightarrow enough to show (4) when

- ① μ is a finite measure,
- ② f_e is bounded, and

Outline of proof

Need to show, due to an extension of [Gross \(1994\)](#), that for all $\delta > 0$ and $\epsilon > 0$,

$$\lim_{n \rightarrow \infty} \frac{1}{F_n} \sum_{g \in F_n} \mu(x \in \Lambda : \delta^{-1} > |f_e(x)| > \delta, |f_g(x)| > \epsilon) = 0, \quad (4)$$

where $f_g = \pm \left(\frac{d\mu \circ \phi_g}{d\mu} \right)^{1/\alpha} f_e \circ \phi_g$, $g \in G$ is a Rosinski representaion of \mathbf{X} .

A probabilistic+analytic truncation argument \Rightarrow enough to show (4) when

- ① μ is a finite measure,
- ② f_e is bounded, and
- ③ the Radon-Nikodym derivatives $\omega_g := \frac{d\mu \circ \phi_g}{d\mu}$, $g \in G$ are uniformly bounded away from zero and infinity.

Outline of proof

Need to show, due to an extension of [Gross \(1994\)](#), that for all $\delta > 0$ and $\epsilon > 0$,

$$\lim_{n \rightarrow \infty} \frac{1}{F_n} \sum_{g \in F_n} \mu(x \in \Lambda : \delta^{-1} > |f_e(x)| > \delta, |f_g(x)| > \epsilon) = 0, \quad (4)$$

where $f_g = \pm \left(\frac{d\mu \circ \phi_g}{d\mu} \right)^{1/\alpha} f_e \circ \phi_g$, $g \in G$ is a Rosinski representaion of \mathbf{X} .

A probabilistic+analytic truncation argument \Rightarrow enough to show (4) when

- ① μ is a finite measure,
- ② f_e is bounded, and
- ③ the Radon-Nikodym derivatives $\omega_g := \frac{d\mu \circ \phi_g}{d\mu}$, $g \in G$ are uniformly bounded away from zero and infinity.

Finally, establish (4) using [Tempelman \(2015\)](#)'s ergodic theorem for Lamperti representations and [Maharam \(1964\)](#) skew-product:

Outline of proof

Need to show, due to an extension of [Gross \(1994\)](#), that for all $\delta > 0$ and $\epsilon > 0$,

$$\lim_{n \rightarrow \infty} \frac{1}{F_n} \sum_{g \in F_n} \mu(x \in \Lambda : \delta^{-1} > |f_e(x)| > \delta, |f_g(x)| > \epsilon) = 0, \quad (4)$$

where $f_g = \pm \left(\frac{d\mu \circ \phi_g}{d\mu} \right)^{1/\alpha} f_e \circ \phi_g$, $g \in G$ is a Rosinski representaion of \mathbf{X} .

A probabilistic+analytic truncation argument \Rightarrow enough to show (4) when

- ① μ is a finite measure,
- ② f_e is bounded, and
- ③ the Radon-Nikodym derivatives $\omega_g := \frac{d\mu \circ \phi_g}{d\mu}$, $g \in G$ are uniformly bounded away from zero and infinity.

Finally, establish (4) using [Tempelman \(2015\)](#)'s ergodic theorem for Lamperti representations and [Maharam \(1964\)](#) skew-product:

$$G \curvearrowright (\Lambda \times (0, \infty), \mu \otimes Leb) \quad \text{by} \quad g^{-1} \cdot (x, y) = \left(\phi_g(x), \frac{y}{\omega_g(x)} \right).$$

Future directions

- Need to view known weak mixing families of examples through our lens - Cohen and Samorodnitsky (2006), Dombry and Guillin-Plantard (2009), Owada and Samorodnitsky (2015), etc.

Future directions

- Need to view known weak mixing families of examples through our lens - [Cohen and Samorodnitsky \(2006\)](#), [Dombry and Guillin-Plantard \(2009\)](#), [Owada and Samorodnitsky \(2015\)](#), etc.
- When will a stationary $S\alpha S$ random field be mixing? Connection to [Dombry and Kabluchko \(2017\)](#).

Future directions

- Need to view known weak mixing families of examples through our lens - [Cohen and Samorodnitsky \(2006\)](#), [Dombry and Guillin-Plantard \(2009\)](#), [Owada and Samorodnitsky \(2015\)](#), etc.
- When will a stationary $S\alpha S$ random field be mixing? Connection to [Dombry and Kabluchko \(2017\)](#).
- Weak mixing for more general stationary $S\alpha S$ random fields ...

Future directions

- Need to view known weak mixing families of examples through our lens - [Cohen and Samorodnitsky \(2006\)](#), [Dombry and Guillin-Plantard \(2009\)](#), [Owada and Samorodnitsky \(2015\)](#), etc.
- When will a stationary $S\alpha S$ random field be mixing? Connection to [Dombry and Kabluchko \(2017\)](#).
- Weak mixing for more general stationary $S\alpha S$ random fields ...
 - ▶ Unavailability of ergodic theorem for nonsingular actions of groups

Future directions


- Need to view known weak mixing families of examples through our lens - [Cohen and Samorodnitsky \(2006\)](#), [Dombry and Guillin-Plantard \(2009\)](#), [Owada and Samorodnitsky \(2015\)](#), etc.
- When will a stationary $S\alpha S$ random field be mixing? Connection to [Dombry and Kabluchko \(2017\)](#).
- Weak mixing for more general stationary $S\alpha S$ random fields ...
 - ▶ Unavailability of ergodic theorem for nonsingular actions of groups - extension of [Lindenstrauss \(2001\)](#)?

Future directions


- Need to view known weak mixing families of examples through our lens - [Cohen and Samorodnitsky \(2006\)](#), [Dombry and Guillin-Plantard \(2009\)](#), [Owada and Samorodnitsky \(2015\)](#), etc.
- When will a stationary $S\alpha S$ random field be mixing? Connection to [Dombry and Kabluchko \(2017\)](#).
- Weak mixing for more general stationary $S\alpha S$ random fields ...
 - ▶ Unavailability of ergodic theorem for nonsingular actions of groups - extension of [Lindenstrauss \(2001\)](#)?
 - ▶ How about groups having Kazhdan's property (T)?

Thank You Very Much


Stable fields and von Neumann Algebras




Stable fields and von Neumann Algebras




Stable fields and von Neumann Algebras




Stable fields and von Neumann Algebras



Stable fields and von Neumann Algebras



Stable fields and von Neumann Algebras



A W^* -rigidity question

R. (2020+): **Minimal group measure space construction** is an invariant for any stationary $S\alpha S$ random field.

A W^* -rigidity question

R. (2020+): **Minimal group measure space construction** is an invariant for any stationary $S\alpha S$ random field.

Question: How much does this invariant “remember” the random field?

A W^* -rigidity question

R. (2020+): Minimal group measure space construction is an invariant for any stationary $S\alpha S$ random field.

Question: How much does this invariant “remember” the random field?

In other words, if two stationary $S\alpha S$ random fields (not necessarily indexed by the same group) have isomorphic minimal group measure space constructions, then do they have similar probabilistic properties?

A W^* -rigidity question

R. (2020+): [Minimal group measure space construction](#) is an invariant for any stationary $S\alpha S$ random field.

Question: How much does this invariant “remember” the random field?

In other words, if two stationary $S\alpha S$ random fields (not necessarily indexed by the same group) have isomorphic minimal group measure space constructions, then do they have similar probabilistic properties?

This question parallels the theory of W^* -rigidity (a term coined by [Sorin Popa](#) - see the survey of [Ioana \(2018\)](#)).

A W^* -rigidity question

R. (2020+): [Minimal group measure space construction](#) is an invariant for any stationary $S\alpha S$ random field.

Question: How much does this invariant “remember” the random field?

In other words, if two stationary $S\alpha S$ random fields (not necessarily indexed by the same group) have isomorphic minimal group measure space constructions, then do they have similar probabilistic properties?

This question parallels the theory of W^* -rigidity (a term coined by [Sorin Popa](#) - see the survey of [Ioana \(2018\)](#)). See also the ICM 2018 lecture of [Adrian Ioana](#) from YouTube.

A W^* -rigidity question

R. (2020+): **Minimal group measure space construction** is an invariant for any stationary $S\alpha S$ random field.

Question: How much does this invariant “remember” the random field?

In other words, if two stationary $S\alpha S$ random fields (not necessarily indexed by the same group) have isomorphic minimal group measure space constructions, then do they have similar probabilistic properties?

This question parallels the theory of W^* -rigidity (a term coined by Sorin Popa - see the survey of Ioana (2018)). See also the ICM 2018 lecture of Adrian Ioana from YouTube.

Theorem (R. (2020+) for $G = \mathbb{Z}^d$, Mj, R. and Sarkar (2022+))

Weak mixing is a W^ -rigid property for stationary $S\alpha S$ random fields indexed by amenable groups.*

Connection to orbit equivalence

Corollary (R. (2020+) for $G = \mathbb{Z}^d$, Mj, R. and Sarkar (2022+))

If two stationary $S\alpha S$ random fields indexed by (possibly two different) amenable groups are generated by orbit equivalent free quasi-invariant actions, then one is weak mixing if and only if the other one is so.

Connection to orbit equivalence

Corollary (R. (2020+) for $G = \mathbb{Z}^d$, Mj, R. and Sarkar (2022+))

If two stationary $S\alpha S$ random fields indexed by (possibly two different) amenable groups are generated by orbit equivalent free quasi-invariant actions, then one is weak mixing if and only if the other one is so.

The indexing groups being possibly different ones is actually very useful in the context of orbit equivalence.

Connection to orbit equivalence

Corollary (R. (2020+) for $G = \mathbb{Z}^d$, Mj, R. and Sarkar (2022+))

If two stationary $S\alpha S$ random fields indexed by (possibly two different) amenable groups are generated by orbit equivalent free quasi-invariant actions, then one is weak mixing if and only if the other one is so.

The indexing groups being possibly different ones is actually very useful in the context of orbit equivalence.

The seminal result of [Connes, Feldman and Weiss \(1981\)](#) states that any quasi-invariant action of an amenable group (more generally, any “amenable action”) is orbit equivalent to a quasi-invariant \mathbb{Z} -action.

Connection to orbit equivalence

Corollary (R. (2020+) for $G = \mathbb{Z}^d$, Mj, R. and Sarkar (2022+))

If two stationary $S\alpha S$ random fields indexed by (possibly two different) amenable groups are generated by orbit equivalent free quasi-invariant actions, then one is weak mixing if and only if the other one is so.

The indexing groups being possibly different ones is actually very useful in the context of orbit equivalence.

The seminal result of [Connes, Feldman and Weiss \(1981\)](#) states that any quasi-invariant action of an amenable group (more generally, any “amenable action”) is orbit equivalent to a quasi-invariant \mathbb{Z} -action.

Therefore, it is now possible to associate a stationary $S\alpha S$ process to any stationary $S\alpha S$ random field indexed by an amenable group in a weak mixing preserving manner.

Connection to orbit equivalence

Corollary (R. (2020+) for $G = \mathbb{Z}^d$, Mj, R. and Sarkar (2022+))

If two stationary $S\alpha S$ random fields indexed by (possibly two different) amenable groups are generated by orbit equivalent free quasi-invariant actions, then one is weak mixing if and only if the other one is so.

The indexing groups being possibly different ones is actually very useful in the context of orbit equivalence.

The seminal result of [Connes, Feldman and Weiss \(1981\)](#) states that any quasi-invariant action of an amenable group (more generally, any “amenable action”) is orbit equivalent to a quasi-invariant \mathbb{Z} -action.

Therefore, it is now possible to associate a stationary $S\alpha S$ process to any stationary $S\alpha S$ random field indexed by an amenable group in a weak mixing preserving manner. **This may help in classification of such fields.**

Connections to hyperbolic geometry

Suppose a discrete non-elementary hyperbolic group (after fixing the basepoint to be e)

$G \curvearrowright (\Lambda = \partial G = \text{Gromov boundary}, \mu = \text{Patterson-Sullivan (prob) measure}).$

Connections to hyperbolic geometry

Suppose a discrete non-elementary hyperbolic group (after fixing the basepoint to be e)

$G \curvearrowright (\Lambda = \partial G = \text{Gromov boundary}, \mu = \text{Patterson-Sullivan (prob) measure}).$

[Kaimanovich \(2000\)](#): μ is the exit measure of many (transient) random walks on G starting from e .

Connections to hyperbolic geometry

Suppose a discrete non-elementary hyperbolic group (after fixing the basepoint to be e)

$G \curvearrowright (\Lambda = \partial G = \text{Gromov boundary}, \mu = \text{Patterson-Sullivan (prob) measure}).$

[Kaimanovich \(2000\)](#): μ is the exit measure of many (transient) random walks on G starting from e .

In this case, it is well-known that the action is null.

Connections to hyperbolic geometry

Suppose a discrete non-elementary hyperbolic group (after fixing the basepoint to be e)

$G \curvearrowright (\Lambda = \partial G = \text{Gromov boundary}, \mu = \text{Patterson-Sullivan (prob) measure})$.

[Kaimanovich \(2000\)](#): μ is the exit measure of many (transient) random walks on G starting from e .

In this case, it is well-known that the action is null.

Theorem (Mj, R. and Sarkar (2022+))

The stationary S α S random field indexed by G and generated by the above boundary action is ergodic.

Connections to hyperbolic geometry

Suppose a discrete non-elementary hyperbolic group (after fixing the basepoint to be e)

$G \curvearrowright (\Lambda = \partial G = \text{Gromov boundary}, \mu = \text{Patterson-Sullivan (prob) measure})$.

Kaimanovich (2000): μ is the exit measure of many (transient) random walks on G starting from e .

In this case, it is well-known that the action is null.

Theorem (Mj, R. and Sarkar (2022+))

The stationary S α S random field indexed by G and generated by the above boundary action is ergodic.

We have shown the same for many such natural geometric actions on various negatively curved spaces,

Connections to hyperbolic geometry

Suppose a discrete non-elementary hyperbolic group (after fixing the basepoint to be e)

$G \curvearrowright (\Lambda = \partial G = \text{Gromov boundary}, \mu = \text{Patterson-Sullivan (prob) measure})$.

Kaimanovich (2000): μ is the exit measure of many (transient) random walks on G starting from e .

In this case, it is well-known that the action is null.

Theorem (Mj, R. and Sarkar (2022+))

The stationary SaaS random field indexed by G and generated by the above boundary action is ergodic.

We have shown the same for many such natural geometric actions on various negatively curved spaces, e.g., the double boundary action of $G = F_d$, $d \geq 2$:

$F_d \curvearrowright (\Lambda = \partial F_d \times \partial F_d \setminus \text{Diagonal}, \mu = \text{a suitable } (\sigma\text{-finite) measure})$.

Group Measure Space Construction

Koopman representation

G -action $\{\phi_t\}$ lifts to the space of all real-valued measurable functions on S by

$$\sigma_t g = g \circ \phi_t, \quad t \in G.$$

This lifted action preserves the \mathcal{L}^∞ -norm but not other \mathcal{L}^p -norms.

However, for each $t \in G$, $\pi_t : \mathcal{L}^2(S, \mu) \rightarrow \mathcal{L}^2(S, \mu)$ given by

$$(\pi_t g)(s) = g \circ \phi_t(s) \left(\frac{d\mu \circ \phi_t}{d\mu}(s) \right)^{1/2}, \quad s \in S$$

defines an isometry. The unitary representation $\{\pi_t\}_{t \in G}$ of G inside $\mathcal{L}^2(S, \mu)$ is called the Koopman representation.

The crossed product relation

Using the cocycle relationship

$$\frac{d\mu \circ \phi_{uv}}{d\mu} = \frac{d\mu \circ \phi_u}{d\mu} \sigma_u \left(\frac{d\mu \circ \phi_v}{d\mu} \right), \quad u, v \in G,$$

one gets that for all $a \in \mathcal{L}^\infty(S, \mu)$ (thought of as acting on $\mathcal{L}^2(S, \mu)$ by multiplication), for all $t \in G$ and for all $g \in \mathcal{L}^2(S, \mu)$,

$$(\pi_t a \pi_{t^{-1}} g)(s) = ((\sigma_t a)g)(s), \quad s \in S. \quad (5)$$

In other words, the Koopman representation “normalizes” $\mathcal{L}^\infty(S, \mu)$ inside $\mathcal{B}(\mathcal{L}^2(S, \mu))$. The group measure space construction is a space, where the crossed product relation (5) is internalized.

Group measure space construction

Consider the von Neumann algebra

$$\mathcal{B}(l^2(G) \otimes \mathcal{L}^2(S, \mu)) = \overline{\mathcal{B}(l^2(G)) \otimes \mathcal{B}(\mathcal{L}^2(S, \mu))}$$

(with the closure being taken with respect to the weak/strong operator topology). Define a representation of G by $t \mapsto u_t := \lambda_t \otimes \pi_t$, where $\{\lambda_t\}$ is the left regular representation and $\{\pi_t\}$ is the Koopman representation. We also represent $\mathcal{L}^\infty(S, \mu)$ by $a \mapsto 1 \otimes \mathcal{M}_a$, where \mathcal{M}_a is the multiplication (by a) operator on $\mathcal{L}^2(S, \mu)$. It can be checked that the following “internal” crossed product relation holds:

$$u_t(1 \otimes \mathcal{M}_a)u_{t^{-1}} = 1 \otimes \mathcal{M}_{\sigma_t a}.$$

Define the *group measure space construction* (also known as *crossed product construction*) as

$$\mathcal{L}^\infty(S, \mu) \rtimes G := \{u_t, 1 \otimes \mathcal{M}_a : t \in G, a \in \mathcal{L}^\infty(S, \mu)\}''.$$

Connections to ergodic theory

It can be shown that the internal crossed product relation implies that any $x \in \mathcal{L}^\infty(S, \mu) \rtimes G$ can be uniquely written as $x = \sum_{t \in G} a_t u_t$ with $\{a_t : t \in G\} \subseteq \mathcal{L}^\infty(S, \mu)$. Thus, we can view x as a $|G| \times |G|$ matrix with entries coming from $\mathcal{L}^\infty(S, \mu)$ that are the same along each left group-diagonal; see, e.g. [Jones \(2009\)](#).

Theorem (see, e.g. [Peterson \(2013\)](#))

The following results hold for a nonsingular G -action $\{\phi_t\}$ and the corresponding group measure space construction defined above.

- ❶ *If the action $\{\phi_t\}_{t \in G}$ is free and ergodic, then $\mathcal{L}^\infty(S, \mu) \rtimes G$ is a factor.*
- ❷ *If $\mathcal{L}^\infty(S, \mu) \rtimes G$ is a factor, then $\{\phi_t\}_{t \in G}$ is ergodic.*
- ❸ *If $\{\phi_t\}_{t \in G}$ is free and ergodic, then the factor $\mathcal{L}^\infty(S, \mu) \rtimes G$ is of type II_1 if and only if $\{\phi_t\}_{t \in G}$ is a positive action.*

Furthermore, if the two nonsingular actions (not necessarily of the same group) are orbit-equivalent, then the corresponding group measure space constructions are isomorphic as von Neumann algebras

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

References

- J. AARONSON (1997): *An Introduction to Infinite Ergodic Theory*. American Mathematical Society, Providence.
- J. ATHREYA, M. MJ and P. ROY (2022+): Stable Random Fields, Patterson-Sullivan measures and Extremal Cocycle Growth. *Probability Theory and Related Fields*, to appear; *arXiv:1809.08295*.
- C. BÉGUIN, A. VALETTE and A. ZUK (1997): On the spectrum of a random walk on the discrete Heisenberg group and the norm of Harper's operator. *J. Geom. Phys.* 21:337–356.
- A. BHATTACHARYA and P. ROY (2018): A large sample test for the length of memory of stationary symmetric stable random fields via nonsingular \mathbb{Z}^d -actions. *J. Appl. Probab.* 55:179–195.
- O. BRATTELI and D. W. ROBINSON (1987): *Operator algebras and quantum statistical mechanics. 1*. Texts and Monographs in Physics. Springer-Verlag, New York, 2nd edition. C^* - and W^* -algebras, symmetry groups, decomposition of states.
- S. COHEN and G. SAMORODNITSKY (2006): Random rewards, fractional Brownian local times and stable self-similar processes. *Ann. Appl. Probab.* 16:1432–1461.
- A. CONNES (1976): Classification of injective factors. Cases II_1 , II_∞ , III_λ , $\lambda \neq 1$. *Ann. of Math. (2)* 104:73–115.

- A. CONNES, J. FELDMAN and B. WEISS (1981): An amenable equivalence relation is generated by a single transformation. *Ergodic Theory Dynam. Systems* 1:431–450 (1982).
- C. DOMBRY and N. GUILLOTIN-PLANTARD (2009): Discrete approximation of a stable self-similar stationary increments process. *Bernoulli* 15:195–222.
- C. DOMBRY and Z. KABLUCHKO (2017): Ergodic decompositions of stationary max-stable processes in terms of their spectral functions. *Stochastic Process. Appl.* 127:1763–1784.
- V. FASEN and P. ROY (2016): Stable random fields, point processes and large deviations. *Stochastic Processes and their Applications* 126:832 – 856.
- D. GRETETE (2011): Random walk on a discrete Heisenberg group. *Rend. Circ. Mat. Palermo (2)* 60:329–335.
- C. HARDIN JR. (1981): Isometries on subspaces of L^p . *Indiana Univ. Math. J.* 30:449–465.
- C. HARDIN JR. (1982): On the spectral representation of symmetric stable processes. *Journal of Multivariate Analysis* 12:385–401.
- M. HOCHMAN (2010): A ratio ergodic theorem for multiparameter non-singular actions. *J. Eur. Math. Soc. (JEMS)* 12:365–383.
- A. IOANA (2011): W^* -superrigidity for Bernoulli actions of property (T) groups. *J. Amer. Math. Soc.* 24:1175–1226.

- A. IOANA (2018): Rigidity for von Neumann algebras. In *Proceedings of the International Congress of Mathematicians—Rio de Janeiro 2018. Vol. III. Invited lectures*. World Sci. Publ., Hackensack, NJ, pp. 1639–1672.
- K. JARRETT (2019): An ergodic theorem for nonsingular actions of the Heisenberg groups. *Trans. Amer. Math. Soc.* 372:5507–5529.
- V. F. R. JONES (2009): *Von Neumann Algebras*. Lectures Notes, University of California, Berkeley.
<https://math.berkeley.edu/~vfr/MATH20909/VonNeumann2009.pdf>.
- P. JUNG, T. OWADA and G. SAMORODNITSKY (2017): Functional central limit theorem for a class of negatively dependent heavy-tailed stationary infinitely divisible processes generated by conservative flows. *Ann. Probab.* 45:2087–2130.
- S. KNUDBY (2011): *Disintegration theory for von Neumann algebras*. Graduate Project, University of Copenhagen.
- U. KRENGEL (1985): *Ergodic Theorems*. De Gruyter, Berlin, New York.
- E. LINDENSTRAUSS (2001): Pointwise theorems for amenable groups. *Invent. Math.* 146:259–295.
- T. MIKOSCH and G. SAMORODNITSKY (2000): Ruin probability with claims modeled by a stationary ergodic stable process. *Ann. Probab.* 28:1814–1851.
- F. J. MURRAY and J. VON NEUMANN (1936): On rings of operators. *Ann. of Math. (2)* 37:116–229.

- T. OWADA and G. SAMORODNITSKY (2015): Maxima of long memory stationary symmetric α -stable processes, and self-similar processes with stationary max-increments. *Bernoulli* 21:1575–1599.
- S. PANIGRAHI, P. ROY and Y. XIAO (2018): Maximal moments and uniform modulus of continuity for stable random fields. *arXiv:1709.07135*.
- J. PETERSON (2010): Examples of group actions which are virtually W^* -superrigid. *arXiv:1002.1745*.
- J. PETERSON (2013): *Notes on von Neumann algebras*. Lectures Notes, Vanderbilt University.
<https://math.vanderbilt.edu/peters10/teaching/spring2013/vonNeumannAlgebras/>
- S. POPA (2006): Strong rigidity of II_1 factors arising from malleable actions of w -rigid groups. II. *Invent. Math.* 165:409–451.
- S. POPA and S. VAES (2010): Group measure space decomposition of II_1 factors and W^* -superrigidity. *Invent. Math.* 182:371–417.
- S. POPA and S. VAES (2014): Unique Cartan decomposition for II_1 factors arising from arbitrary actions of free groups. *Acta Math.* 212:141–198.
- S. RESNICK (1992): *Adventures in Stochastic Processes*. Birkhäuser, Boston.
- S. RESNICK and G. SAMORODNITSKY (2004): Point processes associated with stationary stable processes. *Stochastic Process. Appl.* 114:191–209.
- J. ROSIŃSKI (1994): On uniqueness of the spectral representation of stable processes. *J. Theoret. Probab.* 7:615–634.

- J. ROSIŃSKI (1995): On the structure of stationary stable processes. *Ann. Probab.* 23:1163–1187.
- J. ROSIŃSKI (2000): Decomposition of stationary α -stable random fields. *Ann. Probab.* 28:1797–1813.
- J. ROSIŃSKI and G. SAMORODNITSKY (1996): Classes of mixing stable processes. *Bernoulli* 2:3655–378.
- E. ROY (2007): Ergodic properties of Poissonian ID processes. *Ann. Probab.* 35:551–576.
- E. ROY (2012): Maharam extension and stationary stable processes. *Ann. Probab.* 40:1357–1374.
- P. ROY (2010): Ergodic theory, abelian groups and point processes induced by stable random fields. *Ann. Probab.* 38:770–793.
- P. ROY (2017): Maxima of stable random fields, nonsingular actions and finitely generated abelian groups: a survey. *Indian J. Pure Appl. Math.* 48:513–540.
- P. ROY and G. SAMORODNITSKY (2008): Stationary symmetric α -stable discrete parameter random fields. *J. Theoret. Probab.* 21:212–233.
- G. SAMORODNITSKY (2004): Extreme value theory, ergodic theory, and the boundary between short memory and long memory for stationary stable processes. *Ann. Probab.* 32:1438–1468.

- G. SAMORODNITSKY (2005): Null flows, positive flows and the structure of stationary symmetric stable processes. *Ann. Probab.* 33:1782–1803.
- G. SAMORODNITSKY and M. S. TAQQU (1994): *Stable non-Gaussian random processes*. Stochastic Modeling. Chapman & Hall, New York. Stochastic models with infinite variance.
- S. SARKAR and P. ROY (2018): Stable random fields indexed by finitely generated free groups. *Ann. Probab.* 46:2680 – 2714.
- K. SCHMIDT (1977): *Cocycles on ergodic transformation groups*. Macmillan Company of India, Ltd., Delhi. Macmillan Lectures in Mathematics, Vol. 1.
- I. M. SINGER (1955): Automorphisms of finite factors. *Amer. J. Math.* 77:117–133.
- S. STOEV and M. S. TAQQU (2005): Extremal stochastic integrals: a parallel between max-stable processes and α -stable processes. *Extremes* 8:237–266.
- V. S. SUNDER (1987): *An invitation to von Neumann algebras*. Universitext. Springer-Verlag, New York.
- D. SURGAILIS, J. ROSIŃSKI, V. MANDREKAR and S. CAMBANIS (1993): Stable mixed moving averages. *Probab. Theory Related Fields* 97:543–558.
- A. A. TEMPEL'MAN (1972): Ergodic theorems for general dynamical systems. *Trudy Moskov. Mat. Obšč.* 26:95–132.
- S. VAES (2014): Normalizers inside amalgamated free product von Neumann algebras. *Publ. Res. Inst. Math. Sci.* 50:695–721.

- V. VARADARAJAN (1970): *Geometry of Quantum Theory*, volume 2. Van Nostrand Reinhold, New York.
- Y. WANG, P. ROY and S. A. STOEV (2013): Ergodic properties of sum- and max-stable stationary random fields via null and positive group actions. *Ann. Probab.* 41:206–228.
- Y. WANG and S. A. STOEV (2009): On the structure and representations of max-stable processes. Technical Report 487, Department of Statistics, University of Michigan. *arXiv:0903.3594*.
- Y. WANG and S. A. STOEV (2010a): On the association of sum-and max-stable processes. *Statistics & Probability Letters* 80:480–488.
- Y. WANG and S. A. STOEV (2010b): On the structure and representations of max-stable processes. *Advances in Applied Probability* 42:855–877.
- R. ZIMMER (1984): *Ergodic Theory and Semisimple Groups*. Birkhäuser, Boston.