CLINICAL STUDY PROTOCOL: CARDIOVASTIN PHASE 3 (CV-301)

Protocol ID: CV-301-PH3 | Investigational Product: CardioVastin (20 mg oral daily) | NCT-04829102

Section 1.0 — Study Rationale & Objectives

CardioVastin is a novel synthetic small-molecule dual PCSK9/HMG-CoA inhibitor designed to lower low-density lipoprotein cholesterol (LDL-C) and attenuate coronary plaque progression in high-risk atherosclerotic cardiovascular disease (ASCVD) patients.

Section 2.0 — Study Population & Randomization

The trial randomized **1,200 adult patients** across 48 clinical investigation centers in a 1:1 double-blind fashion to either CardioVastin 20 mg once daily (N=600) or matching placebo (N=600) for a 24-week treatment period.

Section 3.0 — Primary Efficacy Endpoint

Primary Endpoint Definition: The primary efficacy endpoint was the percentage change in serum LDL-C concentration from baseline to Week 24.

Trial Primary Efficacy Results: CardioVastin achieved a statistically significant **48.4% mean reduction in LDL-C at Week 24** ($p < 0.001$ versus placebo). Placebo demonstrated a 1.2% reduction. The primary efficacy endpoint was formally met across all pre-specified patient subgroups.

Section 4.0 — Safety Tolerability & Adverse Events Profile

Overall incidence of treatment-emergent adverse events (TEAEs) was **3.8%** in the CardioVastin arm versus 3.2% in the placebo group. The most commonly reported events were transient headache (1.4%), mild elevated liver transaminases (ALT/AST > 3x ULN, 0.8%), and myalgia (0.6%). Discontinuation due to adverse events was low at 1.1%.