

Northstar Demo Hospital

Perioperative Medicines Guideline

Document ID	NDH-CLN-PERI-003	Version	3.0
Status	ACTIVE	Effective Date	2026-01-01
Department	Clinical Governance & Anaesthesia	Review Date	2028-01-01
Access Roles	doctor, pharmacist	Authority	Institutional clinical guideline
Source Type	Synthetic internal policy	Data Class	Demo / de-identified

1. Purpose

To provide a demo-only, machine-searchable hospital policy for testing retrieval, version metadata, access control, citation display, conflict detection, and safe refusal in a healthcare RAG prototype.

2. Scope

Applies to adult planned procedures requiring a preoperative medication review. This synthetic guideline is designed to test retrieval of recommendations, evidence comparison, conflict handling, and citation traceability.

3. Definitions

Demo Drug A: a fictional medicine used only for software testing. **Elective surgery:** a planned procedure for which medication interruption can be scheduled in advance. **Interruption:** temporary discontinuation or holding of the medicine before the procedure.

4.3 Preoperative medication interruption - Demo Drug A

For adult patients taking Demo Drug A who are scheduled for elective surgery, discontinue Demo Drug A 3 calendar days before the planned procedure, unless a specialist documents an exception.

The recommendation above applies only to this synthetic demonstration corpus. It must not be interpreted as medical advice or applied to a real patient.

5. Exceptions and escalation

Urgent or emergency procedures, pediatric patients, pregnancy, severe renal or hepatic impairment, concurrent high-risk medicines, or any situation not explicitly covered by this document require specialist review. If another ACTIVE approved source gives a different interruption interval, the system should surface the disagreement and escalate rather than silently combine or choose between recommendations.

6. Retrieval and citation anchors

Anchor	Value
Primary topic	Demo Drug A - preoperative / perioperative interruption
Synonyms	hold; stop; discontinue; interrupt; pre-op; before surgery
Recommended citation	Perioperative Medicines Guideline - 4.3 Preoperative medication interruption - Demo Drug A - page 1
ACL	doctor, pharmacist
Status	ACTIVE

Synthetic provenance: created specifically for a healthcare RAG hackathon. No real patient, drug, hospital policy, or clinical recommendation is represented.