

# CS 105: DIC on Discrete Structures

Instructor : S. Akshay

Sept 26, 2023

Lecture 19 – Counting and Combinatorics

Solving Recurrence relations via generating functions

## Last few weeks

### Basic counting techniques and applications

1. Sum and product, bijection, double counting principles
2. Binomial coefficients and binomial theorem, Pascal's triangle
3. Permutations and combinations with/without repetitions
4. Counting subsets, relations, Handshake lemma
5. Stirling's approximation: Estimating  $n!$
6. Recurrence relations and one method to solve them.

## Last few weeks

### Basic counting techniques and applications

1. Sum and product, bijection, double counting principles
2. Binomial coefficients and binomial theorem, Pascal's triangle
3. Permutations and combinations with/without repetitions
4. Counting subsets, relations, Handshake lemma
5. Stirling's approximation: Estimating  $n!$
6. Recurrence relations and one method to solve them.

### Today

Solving recurrence relations via generating functions.

## Solving general linear recurrence sequences

- (C.W) Find a solution for  $a_n = a_{n-1} + 2a_{n-2}$  with  $a_0 = 2$ ,  $a_1 = 7$ ?

## Solving general linear recurrence sequences

- ▶ (C.W) Find a solution for  $a_n = a_{n-1} + 2a_{n-2}$  with  $a_0 = 2$ ,  $a_1 = 7$ ?
- ▶ Find a solution for  $a_n = 6a_{n-1} - 9a_{n-2}$ , with  $a_0 = 1, a_1 = 6$ ? Can you apply the same method for this?

## Solving general linear recurrence sequences

- ▶ (C.W) Find a solution for  $a_n = a_{n-1} + 2a_{n-2}$  with  $a_0 = 2$ ,  $a_1 = 7$ ?
- ▶ Find a solution for  $a_n = 6a_{n-1} - 9a_{n-2}$ , with  $a_0 = 1, a_1 = 6$ ? Can you apply the same method for this? What went wrong?
- ▶ Recall the recurrence for Catalan Numbers:

$$C(n) = \sum_{i=1}^{n-1} C(i)C(n-i) \text{ for } n > 1, C(0) = C(1) = 1.$$

No. of ways to bracket a sum of  $n$  terms s.t. it can be computed by adding two numbers at a time?

## Solving general linear recurrence sequences

- ▶ (C.W) Find a solution for  $a_n = a_{n-1} + 2a_{n-2}$  with  $a_0 = 2$ ,  $a_1 = 7$ ?
- ▶ Find a solution for  $a_n = 6a_{n-1} - 9a_{n-2}$ , with  $a_0 = 1, a_1 = 6$ ? Can you apply the same method for this? What went wrong?
- ▶ Recall the recurrence for Catalan Numbers:

$$C(n) = \sum_{i=1}^{n-1} C(i)C(n-i) \text{ for } n > 1, C(0) = C(1) = 1.$$

No. of ways to bracket a sum of  $n$  terms s.t. it can be computed by adding two numbers at a time?

This method does not work if we have repeated roots (this can be fixed!) and non-linear recurrences.

# Solving general linear recurrence sequences

- ▶ (C.W) Find a solution for  $a_n = a_{n-1} + 2a_{n-2}$  with  $a_0 = 2$ ,  $a_1 = 7$ ?
- ▶ Find a solution for  $a_n = 6a_{n-1} - 9a_{n-2}$ , with  $a_0 = 1, a_1 = 6$ ? Can you apply the same method for this? What went wrong?
- ▶ Recall the recurrence for Catalan Numbers:

$$C(n) = \sum_{i=1}^{n-1} C(i)C(n-i) \text{ for } n > 1, C(0) = C(1) = 1.$$

No. of ways to bracket a sum of  $n$  terms s.t. it can be computed by adding two numbers at a time?

This method does not work if we have repeated roots (this can be fixed!) and non-linear recurrences.

## Reading assignment

Read examples/generalizations from Sections 6.1 and 6.2 from Rosen's book (6th Edition).



# Solving general linear recurrence sequences

- ▶ (C.W) Find a solution for  $a_n = a_{n-1} + 2a_{n-2}$  with  $a_0 = 2$ ,  $a_1 = 7$ ?
- ▶ Find a solution for  $a_n = 6a_{n-1} - 9a_{n-2}$ , with  $a_0 = 1, a_1 = 6$ ? Can you apply the same method for this? What went wrong?
- ▶ Recall the recurrence for Catalan Numbers:

$$C(n) = \sum_{i=1}^{n-1} C(i)C(n-i) \text{ for } n > 1, C(0) = C(1) = 1.$$

No. of ways to bracket a sum of  $n$  terms s.t. it can be computed by adding two numbers at a time?

This method does not work if we have repeated roots (this can be fixed!) and non-linear recurrences.

## Reading assignment

Read examples/generalizations from Sections 6.1 and 6.2 from Rosen's book (6th Edition).

We next consider a method of much wider applicability...

## Solving recurrence relations

By solving, we mean give a closed-form expression for  $n^{th}$  term.

## Solving recurrence relations

Fibonacci rec  $\forall n \geq 2, F_n = F_{n-1} + F_{n-2}, F_0 = F_1 = 1$

## Solving recurrence relations

Fibonacci rec  $\forall n \geq 2, F_n = F_{n-1} + F_{n-2}, F_0 = F_1 = 1$

Proof method 1 (for linear recurrences: try  $F_n = \alpha^n$ !)

1.  $\alpha^n = \alpha^{n-1} + \alpha^{n-2}$  implies  $\alpha^{n-2}(\alpha^2 - \alpha - 1) = 0$ .
2. So if  $\alpha^2 - \alpha - 1 = 0$ , the recurrence holds for all  $n$ .
3. Solving,  $\alpha = \frac{1+\sqrt{5}}{2}, \beta = \frac{1-\sqrt{5}}{2}$
4. Thus, general solution is  $F_n = a(\frac{1+\sqrt{5}}{2})^n + b(\frac{1-\sqrt{5}}{2})^n$ .
5. Use  $F_0$  and  $F_1$  – initial conditions:  $a = \frac{\sqrt{5}+1}{2\sqrt{5}}, b = \frac{\sqrt{5}-1}{2\sqrt{5}}$

## Solving recurrence relations

Fibonacci rec  $\forall n \geq 2, F_n = F_{n-1} + F_{n-2}, F_0 = F_1 = 1$

Proof method 1 (for linear recurrences: try  $F_n = \alpha^n$ !)

1.  $\alpha^n = \alpha^{n-1} + \alpha^{n-2}$  implies  $\alpha^{n-2}(\alpha^2 - \alpha - 1) = 0$ .
2. So if  $\alpha^2 - \alpha - 1 = 0$ , the recurrence holds for all  $n$ .
3. Solving,  $\alpha = \frac{1+\sqrt{5}}{2}, \beta = \frac{1-\sqrt{5}}{2}$
4. Thus, general solution is  $F_n = a(\frac{1+\sqrt{5}}{2})^n + b(\frac{1-\sqrt{5}}{2})^n$ .
5. Use  $F_0$  and  $F_1$  – initial conditions:  $a = \frac{\sqrt{5}+1}{2\sqrt{5}}, b = \frac{\sqrt{5}-1}{2\sqrt{5}}$

$$\text{Thus, } F_n = \frac{\sqrt{5}+1}{2\sqrt{5}} \left(\frac{1+\sqrt{5}}{2}\right)^n + \frac{\sqrt{5}-1}{2\sqrt{5}} \left(\frac{1-\sqrt{5}}{2}\right)^n.$$

## Solving recurrence relations

Fibonacci rec  $\forall n \geq 2, F_n = F_{n-1} + F_{n-2}, F_0 = F_1 = 1$

Proof method 1 (for linear recurrences: try  $F_n = \alpha^n$ !)

1.  $\alpha^n = \alpha^{n-1} + \alpha^{n-2}$  implies  $\alpha^{n-2}(\alpha^2 - \alpha - 1) = 0$ .
2. So if  $\alpha^2 - \alpha - 1 = 0$ , the recurrence holds for all  $n$ .
3. Solving,  $\alpha = \frac{1+\sqrt{5}}{2}, \beta = \frac{1-\sqrt{5}}{2}$
4. Thus, general solution is  $F_n = a(\frac{1+\sqrt{5}}{2})^n + b(\frac{1-\sqrt{5}}{2})^n$ .
5. Use  $F_0$  and  $F_1$  – initial conditions:  $a = \frac{\sqrt{5}+1}{2\sqrt{5}}, b = \frac{\sqrt{5}-1}{2\sqrt{5}}$

$$\text{Thus, } F_n = \frac{\sqrt{5}+1}{2\sqrt{5}} \left(\frac{1+\sqrt{5}}{2}\right)^n + \frac{\sqrt{5}-1}{2\sqrt{5}} \left(\frac{1-\sqrt{5}}{2}\right)^n.$$

This method fails for repeated roots/non-linear recurrences.

## Solving recurrence relations

Fibonacci rec  $\forall n \geq 2, F_n = F_{n-1} + F_{n-2}, F_0 = F_1 = 1$

Proof method 1 (for linear recurrences: try  $F_n = \alpha^n$ !)

1.  $\alpha^n = \alpha^{n-1} + \alpha^{n-2}$  implies  $\alpha^{n-2}(\alpha^2 - \alpha - 1) = 0$ .
2. So if  $\alpha^2 - \alpha - 1 = 0$ , the recurrence holds for all  $n$ .
3. Solving,  $\alpha = \frac{1+\sqrt{5}}{2}, \beta = \frac{1-\sqrt{5}}{2}$
4. Thus, general solution is  $F_n = a(\frac{1+\sqrt{5}}{2})^n + b(\frac{1-\sqrt{5}}{2})^n$ .
5. Use  $F_0$  and  $F_1$  – initial conditions:  $a = \frac{\sqrt{5}+1}{2\sqrt{5}}, b = \frac{\sqrt{5}-1}{2\sqrt{5}}$

$$\text{Thus, } F_n = \frac{\sqrt{5}+1}{2\sqrt{5}} \left(\frac{1+\sqrt{5}}{2}\right)^n + \frac{\sqrt{5}-1}{2\sqrt{5}} \left(\frac{1-\sqrt{5}}{2}\right)^n.$$

This method fails for repeated roots/non-linear recurrences.

Reading assignment: how to tackle repeated roots case.

Read Sections 6.1 and 6.2 from Rosen's book (6th Edition)

## Solving recurrence relations

Fibonacci rec  $\forall n \geq 2, F_n = F_{n-1} + F_{n-2}, F_0 = F_1 = 1$

Proof method 1 (for linear recurrences: try  $F_n = \alpha^n$ !)

1.  $\alpha^n = \alpha^{n-1} + \alpha^{n-2}$  implies  $\alpha^{n-2}(\alpha^2 - \alpha - 1) = 0$ .
2. So if  $\alpha^2 - \alpha - 1 = 0$ , the recurrence holds for all  $n$ .
3. Solving,  $\alpha = \frac{1+\sqrt{5}}{2}, \beta = \frac{1-\sqrt{5}}{2}$
4. Thus, general solution is  $F_n = a(\frac{1+\sqrt{5}}{2})^n + b(\frac{1-\sqrt{5}}{2})^n$ .
5. Use  $F_0$  and  $F_1$  – initial conditions:  $a = \frac{\sqrt{5}+1}{2\sqrt{5}}, b = \frac{\sqrt{5}-1}{2\sqrt{5}}$

$$\text{Thus, } F_n = \frac{\sqrt{5}+1}{2\sqrt{5}} \left(\frac{1+\sqrt{5}}{2}\right)^n + \frac{\sqrt{5}-1}{2\sqrt{5}} \left(\frac{1-\sqrt{5}}{2}\right)^n.$$

This method fails for repeated roots/non-linear recurrences.

Reading assignment: how to tackle repeated roots case.

Read Sections 6.1 and 6.2 from Rosen's book (6th Edition)

We next consider a method of much wider applicability...



## Proof Method 2: Using generating functions

### Fibonacci recurrence relation

For  $n \geq 2$ ,  $F_n = F_{n-1} + F_{n-2}$ ,  $F_0 = F_1 = 1$ .

Compute  $F_n$  in terms of  $n$ .

- Consider the power series...  $\phi(t) = \sum_{n=0}^{\infty} F(n)t^n$ .

## Proof Method 2: Using generating functions

### Fibonacci recurrence relation

For  $n \geq 2$ ,  $F_n = F_{n-1} + F_{n-2}$ ,  $F_0 = F_1 = 1$ .

Compute  $F_n$  in terms of  $n$ .

► Consider the power series...  $\phi(t) = \sum_{n=0}^{\infty} F(n)t^n$ .

$$t\phi(t) = \sum_{n=0}^{\infty} F(n)t^{n+1} = \sum_{n=1}^{\infty} F(n-1)t^n$$

## Proof Method 2: Using generating functions

### Fibonacci recurrence relation

For  $n \geq 2$ ,  $F_n = F_{n-1} + F_{n-2}$ ,  $F_0 = F_1 = 1$ .

Compute  $F_n$  in terms of  $n$ .

► Consider the power series...  $\phi(t) = \sum_{n=0}^{\infty} F(n)t^n$ .

$$t\phi(t) = \sum_{n=0}^{\infty} F(n)t^{n+1} = \sum_{n=1}^{\infty} F(n-1)t^n$$

$$t^2\phi(t) = \sum_{n=0}^{\infty} F(n)t^{n+2} = \sum_{n=2}^{\infty} F(n-2)t^n$$

# Proof Method 2: Using generating functions

## Fibonacci recurrence relation

For  $n \geq 2$ ,  $F_n = F_{n-1} + F_{n-2}$ ,  $F_0 = F_1 = 1$ .

Compute  $F_n$  in terms of  $n$ .

► Consider the power series...  $\phi(t) = \sum_{n=0}^{\infty} F(n)t^n$ .

$$t\phi(t) = \sum_{n=0}^{\infty} F(n)t^{n+1} = \sum_{n=1}^{\infty} F(n-1)t^n$$

$$t^2\phi(t) = \sum_{n=0}^{\infty} F(n)t^{n+2} = \sum_{n=2}^{\infty} F(n-2)t^n$$

$$(t + t^2)\phi(t) = \sum_{n=0}^{\infty} F(n)t^n - 1$$

$$(t + t^2)\phi(t) = \phi(t) - 1$$

## Proof Method 2: Using generating functions

►  $\phi(t) = \frac{1}{1-t-t^2}$

## Proof Method 2: Using generating functions

$$\blacktriangleright \phi(t) = \frac{1}{1-t-t^2} = \frac{1}{(1-\alpha t)(1-\beta t)}$$

## Proof Method 2: Using generating functions

- ▶  $\phi(t) = \frac{1}{1-t-t^2} = \frac{1}{(1-\alpha t)(1-\beta t)} = \frac{a}{1-\alpha t} + \frac{b}{1-\beta t}$
- ▶ Solving we get  $\alpha = \frac{1+\sqrt{5}}{2}, \beta = \frac{1-\sqrt{5}}{2}, a = \frac{\sqrt{5}+1}{2\sqrt{5}}, b = \frac{\sqrt{5}-1}{2\sqrt{5}}$

## Proof Method 2: Using generating functions

- ▶  $\phi(t) = \frac{1}{1-t-t^2} = \frac{1}{(1-\alpha t)(1-\beta t)} = \frac{a}{1-\alpha t} + \frac{b}{1-\beta t}$
- ▶ Solving we get  $\alpha = \frac{1+\sqrt{5}}{2}, \beta = \frac{1-\sqrt{5}}{2}, a = \frac{\sqrt{5}+1}{2\sqrt{5}}, b = \frac{\sqrt{5}-1}{2\sqrt{5}}$
- ▶ Now  $\phi(t) = a(1 + \alpha t + \alpha^2 t^2 + \dots) + b(1 + \beta t + \beta^2 t^2 + \dots)$



## Proof Method 2: Using generating functions

- ▶  $\phi(t) = \frac{1}{1-t-t^2} = \frac{1}{(1-\alpha t)(1-\beta t)} = \frac{a}{1-\alpha t} + \frac{b}{1-\beta t}$
- ▶ Solving we get  $\alpha = \frac{1+\sqrt{5}}{2}, \beta = \frac{1-\sqrt{5}}{2}, a = \frac{\sqrt{5}+1}{2\sqrt{5}}, b = \frac{\sqrt{5}-1}{2\sqrt{5}}$
- ▶ Now  $\phi(t) = a(1 + \alpha t + \alpha^2 t^2 + \dots) + b(1 + \beta t + \beta^2 t^2 + \dots)$
- ▶ Equating coefficients of  $t^n$  we get  $F_n = a\alpha^n + b\beta^n$ .

## Proof Method 2: Using generating functions

- ▶  $\phi(t) = \frac{1}{1-t-t^2} = \frac{1}{(1-\alpha t)(1-\beta t)} = \frac{a}{1-\alpha t} + \frac{b}{1-\beta t}$
- ▶ Solving we get  $\alpha = \frac{1+\sqrt{5}}{2}, \beta = \frac{1-\sqrt{5}}{2}, a = \frac{\sqrt{5}+1}{2\sqrt{5}}, b = \frac{\sqrt{5}-1}{2\sqrt{5}}$
- ▶ Now  $\phi(t) = a(1 + \alpha t + \alpha^2 t^2 + \dots) + b(1 + \beta t + \beta^2 t^2 + \dots)$
- ▶ Equating coefficients of  $t^n$  we get  $F_n = a\alpha^n + b\beta^n$ .

thus, as before

$$F(n) = \frac{\sqrt{5}+1}{2\sqrt{5}} \left( \frac{1+\sqrt{5}}{2} \right)^n + \frac{\sqrt{5}-1}{2\sqrt{5}} \left( \frac{1-\sqrt{5}}{2} \right)^n$$

# Properties of generating functions

## Definition

The **(ordinary) generating function** for a sequence  $a_0, a_1, \dots \in \mathbb{R}$  is the infinite series  $\phi(x) = \sum_{k=0}^{\infty} a_k x^k$ .

- Let  $f(x) = \sum_{k=0}^{\infty} a_k x^k$ ,  $g(x) = \sum_{k=0}^{\infty} b_k x^k$ . Then
1. If  $f(x) = g(x)$ , then  $a_k = b_k$  for all  $k$ .
  2.  $f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$ ,
  3.  $f(x)g(x) = \sum_{k=0}^{\infty} (\sum_{j=0}^k a_j b_{k-j}) x^k$ ,
  4.  $\frac{d}{dx} (\sum_{k=0}^{\infty} a_k x^k) = \sum_{k=1}^{\infty} (k a_k) x^{k-1}$

# Properties of generating functions

## Definition

The **(ordinary) generating function** for a sequence  $a_0, a_1, \dots \in \mathbb{R}$  is the infinite series  $\phi(x) = \sum_{k=0}^{\infty} a_k x^k$ .

- ▶ Let  $f(x) = \sum_{k=0}^{\infty} a_k x^k$ ,  $g(x) = \sum_{k=0}^{\infty} b_k x^k$ . Then
  1. If  $f(x) = g(x)$ , then  $a_k = b_k$  for all  $k$ .
  2.  $f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$ ,
  3.  $f(x)g(x) = \sum_{k=0}^{\infty} (\sum_{j=0}^k a_j b_{k-j}) x^k$ ,
  4.  $\frac{d}{dx} (\sum_{k=0}^{\infty} a_k x^k) = \sum_{k=1}^{\infty} (k a_k) x^{k-1}$
- ▶ Let  $u \in \mathbb{R}$ ,  $k \in \mathbb{Z}^{\geq 0}$ , Then **extended binomial coefficient**  $\binom{u}{k}$  is defined as  $\binom{u}{k} = \frac{u(u-1)\dots(u-k+1)}{k!}$  if  $k > 0$  and  $= 1$  if  $k = 0$ .
- ▶ What if  $u = -n$  for  $n \in \mathbb{N}$ ?

# Properties of generating functions

## Definition

The **(ordinary) generating function** for a sequence  $a_0, a_1, \dots \in \mathbb{R}$  is the infinite series  $\phi(x) = \sum_{k=0}^{\infty} a_k x^k$ .

- ▶ Let  $f(x) = \sum_{k=0}^{\infty} a_k x^k$ ,  $g(x) = \sum_{k=0}^{\infty} b_k x^k$ . Then
  1. If  $f(x) = g(x)$ , then  $a_k = b_k$  for all  $k$ .
  2.  $f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$ ,
  3.  $f(x)g(x) = \sum_{k=0}^{\infty} (\sum_{j=0}^k a_j b_{k-j}) x^k$ ,
  4.  $\frac{d}{dx} (\sum_{k=0}^{\infty} a_k x^k) = \sum_{k=1}^{\infty} (k a_k) x^{k-1}$
- ▶ Let  $u \in \mathbb{R}$ ,  $k \in \mathbb{Z}^{\geq 0}$ , Then **extended binomial coefficient**  $\binom{u}{k}$  is defined as  $\binom{u}{k} = \frac{u(u-1)\dots(u-k+1)}{k!}$  if  $k > 0$  and  $= 1$  if  $k = 0$ .
- ▶ What if  $u = -n$  for  $n \in \mathbb{N}$ ?

## The extended binomial theorem

Let  $u \in \mathbb{R}$ ,  $(1+x)^u = \sum_{k=0}^{\infty} \binom{u}{k} x^k$ .

# Properties of generating functions

## Definition

The **(ordinary) generating function** for a sequence  $a_0, a_1, \dots \in \mathbb{R}$  is the infinite series  $\phi(x) = \sum_{k=0}^{\infty} a_k x^k$ .

- ▶ Let  $f(x) = \sum_{k=0}^{\infty} a_k x^k$ ,  $g(x) = \sum_{k=0}^{\infty} b_k x^k$ . Then
  1. If  $f(x) = g(x)$ , then  $a_k = b_k$  for all  $k$ .
  2.  $f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$ ,
  3.  $f(x)g(x) = \sum_{k=0}^{\infty} (\sum_{j=0}^k a_j b_{k-j}) x^k$ ,
  4.  $\frac{d}{dx} (\sum_{k=0}^{\infty} a_k x^k) = \sum_{k=1}^{\infty} (k a_k) x^{k-1}$
- ▶ Let  $u \in \mathbb{R}$ ,  $k \in \mathbb{Z}^{\geq 0}$ , Then **extended binomial coefficient**  $\binom{u}{k}$  is defined as  $\binom{u}{k} = \frac{u(u-1)\dots(u-k+1)}{k!}$  if  $k > 0$  and  $= 1$  if  $k = 0$ .
- ▶ What if  $u = -n$  for  $n \in \mathbb{N}$ ?

## The extended binomial theorem

Let  $u \in \mathbb{R}$ ,  $(1+x)^u = \sum_{k=0}^{\infty} \binom{u}{k} x^k$ .

If you don't like this, take  $x \in \mathbb{R}$ ,  $|x| < 1$ .

# Simple examples using generating functions

Standard identities:

$$\blacktriangleright \frac{1}{1-ax} = \sum_{k=0}^{\infty} a^k x^k$$

$$\blacktriangleright \frac{1}{1-x^r} = \sum_{k=0}^{\infty} x^{rk}$$

$$\blacktriangleright e^x = \sum_{k=0}^{\infty} \frac{x^k}{k!}$$

# Simple examples using generating functions

Standard identities:

$$\blacktriangleright \frac{1}{1-ax} = \sum_{k=0}^{\infty} a^k x^k$$

$$\blacktriangleright \frac{1}{1-x^r} = \sum_{k=0}^{\infty} x^{rk}$$

$$\blacktriangleright e^x = \sum_{k=0}^{\infty} \frac{x^k}{k!}$$

Class work:

1. Solve the recurrence  $a_k = 4a_{k-1}$  with  $a_0 = 3$ .



# Simple examples using generating functions

## Standard identities:

- ▶  $\frac{1}{1-ax} = \sum_{k=0}^{\infty} a^k x^k$
- ▶  $\frac{1}{1-x^r} = \sum_{k=0}^{\infty} x^{rk}$
- ▶  $e^x = \sum_{k=0}^{\infty} \frac{x^k}{k!}$

## Class work:

1. Solve the recurrence  $a_k = 4a_{k-1}$  with  $a_0 = 3$ .
2. Solve the recurrence  $a_k = 8a_{k-1} + 10^{k-1}$  with  $a_0 = 1, a_1 = 9$ .

## Other applications of generating functions

- ▶ What is the number of ways  $a_k$  of selecting  $k$  elements from an  $n$  element set if repetitions are allowed?

## Other applications of generating functions

- ▶ What is the number of ways  $a_k$  of selecting  $k$  elements from an  $n$  element set if repetitions are allowed?
  - ▶ Let  $\phi(x) = \sum_{k=0}^{\infty} a_k x^k$ .

## Other applications of generating functions

- ▶ What is the number of ways  $a_k$  of selecting  $k$  elements from an  $n$  element set if repetitions are allowed?
  - ▶ Let  $\phi(x) = \sum_{k=0}^{\infty} a_k x^k$ .
  - ▶ Observe that  $\phi(x) = (1 + x + x^2 + \dots)^n = (1 - x)^{-n}$

## Other applications of generating functions

- ▶ What is the number of ways  $a_k$  of selecting  $k$  elements from an  $n$  element set if repetitions are allowed?
  - ▶ Let  $\phi(x) = \sum_{k=0}^{\infty} a_k x^k$ .
  - ▶ Observe that  $\phi(x) = (1 + x + x^2 + \dots)^n = (1 - x)^{-n}$
  - ▶ Expand this by the extended binomial theorem and compare coefficients of  $x^k$ .

## Other applications of generating functions

- ▶ What is the number of ways  $a_k$  of selecting  $k$  elements from an  $n$  element set if repetitions are allowed?
  - ▶ Let  $\phi(x) = \sum_{k=0}^{\infty} a_k x^k$ .
  - ▶ Observe that  $\phi(x) = (1 + x + x^2 + \dots)^n = (1 - x)^{-n}$
  - ▶ Expand this by the extended binomial theorem and compare coefficients of  $x^k$ .
  - ▶  $a_k = \binom{-n}{k}(-1)^k = (-1)^k \binom{n+k-1}{k}(-1)^k = \binom{n+k-1}{k}$ .

## Other applications of generating functions

- ▶ What is the number of ways  $a_k$  of selecting  $k$  elements from an  $n$  element set if repetitions are allowed?
  - ▶ Let  $\phi(x) = \sum_{k=0}^{\infty} a_k x^k$ .
  - ▶ Observe that  $\phi(x) = (1 + x + x^2 + \dots)^n = (1 - x)^{-n}$
  - ▶ Expand this by the extended binomial theorem and compare coefficients of  $x^k$ .
  - ▶  $a_k = \binom{-n}{k}(-1)^k = (-1)^k \binom{n+k-1}{k}(-1)^k = \binom{n+k-1}{k}$ .
  - ▶ (H.W) What if there must be  $\geq 1$  element of each type?

## Other applications of generating functions

- ▶ What is the number of ways  $a_k$  of selecting  $k$  elements from an  $n$  element set if repetitions are allowed?
  - ▶ Let  $\phi(x) = \sum_{k=0}^{\infty} a_k x^k$ .
  - ▶ Observe that  $\phi(x) = (1 + x + x^2 + \dots)^n = (1 - x)^{-n}$
  - ▶ Expand this by the extended binomial theorem and compare coefficients of  $x^k$ .
  - ▶  $a_k = \binom{-n}{k}(-1)^k = (-1)^k \binom{n+k-1}{k}(-1)^k = \binom{n+k-1}{k}$ .
  - ▶ (H.W) What if there must be  $\geq 1$  element of each type?
- ▶ Proving binomial identities: Show that  $\sum_{k=0}^n \binom{n}{k}^2 = \binom{2n}{n}$ .



## Other applications of generating functions

- ▶ What is the number of ways  $a_k$  of selecting  $k$  elements from an  $n$  element set if repetitions are allowed?
  - ▶ Let  $\phi(x) = \sum_{k=0}^{\infty} a_k x^k$ .
  - ▶ Observe that  $\phi(x) = (1 + x + x^2 + \dots)^n = (1 - x)^{-n}$
  - ▶ Expand this by the extended binomial theorem and compare coefficients of  $x^k$ .
  - ▶  $a_k = \binom{-n}{k}(-1)^k = (-1)^k \binom{n+k-1}{k}(-1)^k = \binom{n+k-1}{k}$ .
  - ▶ (H.W) What if there must be  $\geq 1$  element of each type?
- ▶ Proving binomial identities: Show that  $\sum_{k=0}^n \binom{n}{k}^2 = \binom{2n}{n}$ .
  - ▶ Compare coefficients of  $x^n$  in  $(1+x)^{2n} = ((1+x)^n)^2$ .

## Other applications of generating functions

- ▶ What is the number of ways  $a_k$  of selecting  $k$  elements from an  $n$  element set if repetitions are allowed?
  - ▶ Let  $\phi(x) = \sum_{k=0}^{\infty} a_k x^k$ .
  - ▶ Observe that  $\phi(x) = (1 + x + x^2 + \dots)^n = (1 - x)^{-n}$
  - ▶ Expand this by the extended binomial theorem and compare coefficients of  $x^k$ .
  - ▶  $a_k = \binom{-n}{k}(-1)^k = (-1)^k \binom{n+k-1}{k}(-1)^k = \binom{n+k-1}{k}$ .
  - ▶ (H.W) What if there must be  $\geq 1$  element of each type?
- ▶ Proving binomial identities: Show that  $\sum_{k=0}^n \binom{n}{k}^2 = \binom{2n}{n}$ .
  - ▶ Compare coefficients of  $x^n$  in  $(1+x)^{2n} = ((1+x)^n)^2$ .
- ▶ (H.W) Write a recurrence for the number of derrangements. That is, no. of ways to arrange  $n$  letters into  $n$  addressed envelopes such that no letter goes to the correct envelope.

## Other applications of generating functions

- ▶ What is the number of ways  $a_k$  of selecting  $k$  elements from an  $n$  element set if repetitions are allowed?
  - ▶ Let  $\phi(x) = \sum_{k=0}^{\infty} a_k x^k$ .
  - ▶ Observe that  $\phi(x) = (1 + x + x^2 + \dots)^n = (1 - x)^{-n}$
  - ▶ Expand this by the extended binomial theorem and compare coefficients of  $x^k$ .
  - ▶  $a_k = \binom{-n}{k}(-1)^k = (-1)^k \binom{n+k-1}{k}(-1)^k = \binom{n+k-1}{k}$ .
  - ▶ (H.W) What if there must be  $\geq 1$  element of each type?
- ▶ Proving binomial identities: Show that  $\sum_{k=0}^n \binom{n}{k}^2 = \binom{2n}{n}$ .
  - ▶ Compare coefficients of  $x^n$  in  $(1+x)^{2n} = ((1+x)^n)^2$ .
- ▶ (H.W) Write a recurrence for the number of derrangements. That is, no. of ways to arrange  $n$  letters into  $n$  addressed envelopes such that no letter goes to the correct envelope.
- ▶ (H.W) How many ways can a convex  $n$ -sided polygon be cut into triangles by adding non-intersecting diagonals (i.e., connecting vertices with non-crossing lines)? Write a recurrence and solve it!