

# MA 105: D3 Lecture 17

Ravi Raghunathan

Department of Mathematics

September 14, 2023



# Partial Derivatives

Exercise 1. Let  $z = \ln(e^x + e^y)$ . Express  $\frac{\partial z}{\partial x}$  in terms of  $\frac{\partial z}{\partial y}$ .

# Partial Derivatives

Exercise 1. Let  $z = \ln(e^x + e^y)$ . Express  $\frac{\partial z}{\partial x}$  in terms of  $\frac{\partial z}{\partial y}$ .

# Partial Derivatives

**Exercise 1.** Let  $z = \ln(e^x + e^y)$ . Express  $\frac{\partial z}{\partial x}$  in terms of  $\frac{\partial z}{\partial y}$ .

**Solution:**  $\frac{\partial z}{\partial x} = 1 - \frac{\partial z}{\partial y}$ .

A relation between the different partial derivatives of a function (including the zeroth partial derivative which by convention is the function itself) is called a partial differential equation.

# Partial Derivatives

**Exercise 1.** Let  $z = \ln(e^x + e^y)$ . Express  $\frac{\partial z}{\partial x}$  in terms of  $\frac{\partial z}{\partial y}$ .

**Solution:**  $\frac{\partial z}{\partial x} = 1 - \frac{\partial z}{\partial y}$ .

A relation between the different partial derivatives of a function (including the zeroth partial derivative which by convention is the function itself) is called a partial differential equation.

**Exercise 2.** Find a partial differential equation satisfied by the Cobb-Douglas production function  $P = bL^\alpha K^\beta$ .

# Partial Derivatives

**Exercise 1.** Let  $z = \ln(e^x + e^y)$ . Express  $\frac{\partial z}{\partial x}$  in terms of  $\frac{\partial z}{\partial y}$ .

**Solution:**  $\frac{\partial z}{\partial x} = 1 - \frac{\partial z}{\partial y}$ .

A relation between the different partial derivatives of a function (including the zeroth partial derivative which by convention is the function itself) is called a partial differential equation.

**Exercise 2.** Find a partial differential equation satisfied by the Cobb-Douglas production function  $P = bL^\alpha K^\beta$ .

# Partial Derivatives

**Exercise 1.** Let  $z = \ln(e^x + e^y)$ . Express  $\frac{\partial z}{\partial x}$  in terms of  $\frac{\partial z}{\partial y}$ .

**Solution:**  $\frac{\partial z}{\partial x} = 1 - \frac{\partial z}{\partial y}$ .

A relation between the different partial derivatives of a function (including the zeroth partial derivative which by convention is the function itself) is called a partial differential equation.

**Exercise 2.** Find a partial differential equation satisfied by the Cobb-Douglas production function  $P = bL^\alpha K^\beta$ .

**Solution:**



# Partial Derivatives

**Exercise 1.** Let  $z = \ln(e^x + e^y)$ . Express  $\frac{\partial z}{\partial x}$  in terms of  $\frac{\partial z}{\partial y}$ .

**Solution:**  $\frac{\partial z}{\partial x} = 1 - \frac{\partial z}{\partial y}$ .

A relation between the different partial derivatives of a function (including the zeroth partial derivative which by convention is the function itself) is called a partial differential equation.

**Exercise 2.** Find a partial differential equation satisfied by the Cobb-Douglas production function  $P = bL^\alpha K^\beta$ .

**Solution:**

$$L \frac{\partial P}{\partial L} + K \frac{\partial P}{\partial K} = (\alpha + \beta)P.$$

Note that in many situations  $\beta = 1 - \alpha$ . The factor “ $b$ ” is sometimes called the “total factor productivity”.

# Partial Derivatives

**Exercise 1.** Let  $z = \ln(e^x + e^y)$ . Express  $\frac{\partial z}{\partial x}$  in terms of  $\frac{\partial z}{\partial y}$ .

**Solution:**  $\frac{\partial z}{\partial x} = 1 - \frac{\partial z}{\partial y}$ .

A relation between the different partial derivatives of a function (including the zeroth partial derivative which by convention is the function itself) is called a partial differential equation.

**Exercise 2.** Find a partial differential equation satisfied by the Cobb-Douglas production function  $P = bL^\alpha K^\beta$ .

**Solution:**

$$L \frac{\partial P}{\partial L} + K \frac{\partial P}{\partial K} = (\alpha + \beta)P.$$

Note that in many situations  $\beta = 1 - \alpha$ . The factor “ $b$ ” is sometimes called the “total factor productivity”.

**Exercise 3.** Show that for an ideal gas of fixed mass,

$$\frac{\partial P}{\partial V} \frac{\partial V}{\partial T} \frac{\partial T}{\partial P} = -1.$$

# Partial Derivatives

**Exercise 1.** Let  $z = \ln(e^x + e^y)$ . Express  $\frac{\partial z}{\partial x}$  in terms of  $\frac{\partial z}{\partial y}$ .

**Solution:**  $\frac{\partial z}{\partial x} = 1 - \frac{\partial z}{\partial y}$ .

A relation between the different partial derivatives of a function (including the zeroth partial derivative which by convention is the function itself) is called a partial differential equation.

**Exercise 2.** Find a partial differential equation satisfied by the Cobb-Douglas production function  $P = bL^\alpha K^\beta$ .

**Solution:**

$$L \frac{\partial P}{\partial L} + K \frac{\partial P}{\partial K} = (\alpha + \beta)P.$$

Note that in many situations  $\beta = 1 - \alpha$ . The factor “ $b$ ” is sometimes called the “total factor productivity”.

**Exercise 3.** Show that for an ideal gas of fixed mass,

$$\frac{\partial P}{\partial V} \frac{\partial V}{\partial T} \frac{\partial T}{\partial P} = -1.$$

( $PV = mrT$ ).

# Differentiability

**Exercise 4.** Show that the function  $f(x, y) = x^2 + y^2$  is differentiable at every point in  $\mathbb{R}^2$  from first principles, that is from the definition of differentiability (how will you do it if we omit “from first principles” from the question)?

# Differentiability

**Exercise 4.** Show that the function  $f(x, y) = x^2 + y^2$  is differentiable at every point in  $\mathbb{R}^2$  from first principles, that is from the definition of differentiability (how will you do it if we omit “from first principles” from the question)?

# Differentiability

**Exercise 4.** Show that the function  $f(x, y) = x^2 + y^2$  is differentiable at every point in  $\mathbb{R}^2$  from first principles, that is from the definition of differentiability (how will you do it if we omit “from first principles” from the question)?

**Solution:** The partial derivatives at  $(x_0, y_0)$  are  $2x_0$  and  $2y_0$ . We have

$$(x_0 + h)^2 + (y_0 + k)^2 - 2x_0h - 2y_0h = h^2 + k^2 = \|(h, k)\|^2.$$

Obviously,

$$\lim_{(h,k) \rightarrow 0} \frac{(x_0 + h)^2 + (y_0 + k)^2 - 2x_0h - 2y_0h}{\|(h, k)\|} = \lim_{(h,k) \rightarrow 0} \|(h, k)\| = 0.$$

If we are not required to use the definition, we can observe that the two partial derivatives ( $2x$  and  $2y$  respectively) are continuous functions everywhere, and hence, also in any disc around any point  $(x_0, y_0)$ . Hence  $f(x, y)$  is differentiable at every point in  $\mathbb{R}^2$ .

# The derivative

**Exercise 5.** Find the tangent plane at the given point for the surface  $z = f(x, y)$  for the following functions (why are these functions differentiable?).

# The derivative

**Exercise 5.** Find the tangent plane at the given point for the surface  $z = f(x, y)$  for the following functions (why are these functions differentiable?).

(a)  $z = \ln(x - 2y)$  at  $(3, 1, 0)$ , (b)  $z = xe^{xy}$  at  $(2, 0, 2)$ .



# The derivative

**Exercise 5.** Find the tangent plane at the given point for the surface  $z = f(x, y)$  for the following functions (why are these functions differentiable?).

(a)  $z = \ln(x - 2y)$  at  $(3, 1, 0)$ , (b)  $z = xe^{xy}$  at  $(2, 0, 2)$ .

**Exercise 6.** Find the derivative matrix/Jacobian matrix/total derivative for the following functions

# The derivative

**Exercise 5.** Find the tangent plane at the given point for the surface  $z = f(x, y)$  for the following functions (why are these functions differentiable?).

(a)  $z = \ln(x - 2y)$  at  $(3, 1, 0)$ , (b)  $z = xe^{xy}$  at  $(2, 0, 2)$ .

**Exercise 6.** Find the derivative matrix/Jacobian matrix/total derivative for the following functions

(a)  $f(x, t) = e^{-2x} \cos(2\pi t)$ , (b)  $L(x, y, z) = xze^{-y^2 - z^2}$

# The gradient and the Chain rule

**Exercise 7.** Find the maximum rate of change of  $f(p, q, r) = \arctan(pqr)$  at the point  $(1, 2, 1)$  and the direction in which it occurs.

# The gradient and the Chain rule

**Exercise 7.** Find the maximum rate of change of  $f(p, q, r) = \arctan(pqr)$  at the point  $(1, 2, 1)$  and the direction in which it occurs.

**Exercise 8.** At what point on the paraboloid  $y = x^2 + z^2$  is the tangent plane parallel to the plane  $x + 2y + 3z = 1$ ?

# The gradient and the Chain rule

**Exercise 7.** Find the maximum rate of change of  $f(p, q, r) = \arctan(pqr)$  at the point  $(1, 2, 1)$  and the direction in which it occurs.

**Exercise 8.** At what point on the paraboloid  $y = x^2 + z^2$  is the tangent plane parallel to the plane  $x + 2y + 3z = 1$ ?

**Exercise 9.** Show that the sum of the  $x$ -,  $y$ -, and  $z$ - intercepts of any tangent plane to the surface  $\sqrt{x} + \sqrt{y} + \sqrt{z} = \sqrt{c}$  is a constant.

# The gradient and the Chain rule

**Exercise 7.** Find the maximum rate of change of  $f(p, q, r) = \arctan(pqr)$  at the point  $(1, 2, 1)$  and the direction in which it occurs.

**Exercise 8.** At what point on the paraboloid  $y = x^2 + z^2$  is the tangent plane parallel to the plane  $x + 2y + 3z = 1$ ?

**Exercise 9.** Show that the sum of the  $x$ -,  $y$ -, and  $z$ - intercepts of any tangent plane to the surface  $\sqrt{x} + \sqrt{y} + \sqrt{z} = \sqrt{c}$  is a constant.

**Exercise 10.** Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  be a function satisfying  $f(tx, ty) = t^n f(x, y)$  for all  $t, x$  and  $y$ , and some positive integer  $n$ . Show that

$$x_0 \frac{\partial f}{\partial x}(x_0, y_0) + y_0 \frac{\partial f}{\partial y}(x_0, y_0) = nf(x_0, y_0).$$