

# MA 110 - Ordinary Differential Equations

Santanu Dey

Department of Mathematics,  
Indian Institute of Technology Bombay,  
Powai, Mumbai 76  
*santanudey@iitb.ac.in*

March 12, 2024

# Outline of the lecture


- Orthogonal Trajectories
- Lipschitz continuity
- Existence & uniqueness

# Orthogonal Trajectories

If two families of curves always intersect each other at right angles, then they are said to be orthogonal trajectories of each other.

# Orthogonal Trajectories

If two families of curves always intersect each other at right angles, then they are said to be orthogonal trajectories of each other.



# Working Rule

To find the OT of a family of curves

$$F(x, y, c) = 0.$$

# Working Rule

To find the OT of a family of curves

$$F(x, y, c) = 0.$$

- Find the DE  $\frac{dy}{dx} = f(x, y)$ .

# Working Rule

To find the OT of a family of curves

$$F(x, y, c) = 0.$$

- Find the DE  $\frac{dy}{dx} = f(x, y)$ .
- Slopes of the OT's are given by

$$\frac{dy}{dx} = -\frac{1}{f(x, y)}.$$

To find the OT of a family of curves

$$F(x, y, c) = 0.$$

- Find the DE  $\frac{dy}{dx} = f(x, y)$ .
- Slopes of the OT's are given by

$$\frac{dy}{dx} = -\frac{1}{f(x, y)}.$$

- Obtain a one parameter family of curves  $G(x, y, c) = 0$  as solutions of the above DE.



To find the OT of a family of curves

$$F(x, y, c) = 0.$$

- Find the DE  $\frac{dy}{dx} = f(x, y)$ .
- Slopes of the OT's are given by

$$\frac{dy}{dx} = -\frac{1}{f(x, y)}.$$

- Obtain a one parameter family of curves  $G(x, y, c) = 0$  as solutions of the above DE.

( Leaving a part certain trajectories that are vertical lines!)

# Example

Find the set of OT's of the family of circles  $x^2 + y^2 = c^2$ .

# Example

Find the set of OT's of the family of circles  $x^2 + y^2 = c^2$ .

$$x + y \frac{dy}{dx} = 0$$

# Example

Find the set of OT's of the family of circles  $x^2 + y^2 = c^2$ .

$$x + y \frac{dy}{dx} = 0 \implies \frac{dy}{dx} = -\frac{x}{y}$$

## Example

Find the set of OT's of the family of circles  $x^2 + y^2 = c^2$ .

$$x + y \frac{dy}{dx} = 0 \implies \frac{dy}{dx} = -\frac{x}{y}$$

The slope of OT's are

## Example

Find the set of OT's of the family of circles  $x^2 + y^2 = c^2$ .

$$x + y \frac{dy}{dx} = 0 \implies \frac{dy}{dx} = -\frac{x}{y}$$

The slope of OT's are  $\frac{dy}{dx} = \frac{y}{x}$

## Example

Find the set of OT's of the family of circles  $x^2 + y^2 = c^2$ .

$$x + y \frac{dy}{dx} = 0 \implies \frac{dy}{dx} = -\frac{x}{y}$$

The slope of OT's are  $\frac{dy}{dx} = \frac{y}{x} \implies y = kx$ .

## Example

Find the set of OT's of the family of circles  $x^2 + y^2 = c^2$ .

$$x + y \frac{dy}{dx} = 0 \implies \frac{dy}{dx} = -\frac{x}{y}$$

The slope of OT's are  $\frac{dy}{dx} = \frac{y}{x} \implies y = kx$ .

Hence the orthogonal trajectories are given by  $y = kx$ .



- 1 Let  $f$  be a real function defined on  $D$ , where  $D$  is either a domain or a closed domain of the  $xy$  plane. The function  $f$  is said to be **bounded** in  $D$  if there exists a positive number  $M$  such that

$$|f(x, y)| \leq M$$

for all  $(x, y)$  in  $D$ .

# Definitions

- 1 Let  $f$  be a real function defined on  $D$ , where  $D$  is either a domain or a closed domain of the  $xy$  plane. The function  $f$  is said to be **bounded** in  $D$  if there exists a positive number  $M$  such that

$$|f(x, y)| \leq M$$

for all  $(x, y)$  in  $D$ .

- 2 Let  $f$  be defined and continuous on a closed rectangle  $R : a \leq x \leq b, c \leq y \leq d$ .

# Definitions

- 1 Let  $f$  be a real function defined on  $D$ , where  $D$  is either a domain or a closed domain of the  $xy$  plane. The function  $f$  is said to be **bounded** in  $D$  if there exists a positive number  $M$  such that

$$|f(x, y)| \leq M$$

for all  $(x, y)$  in  $D$ .

- 2 Let  $f$  be defined and continuous on a closed rectangle  $R : a \leq x \leq b, c \leq y \leq d$ . Then,  $f$  is bounded in  $R$ .

# Definitions

- 1 Let  $f$  be a real function defined on  $D$ , where  $D$  is either a domain or a closed domain of the  $xy$  plane. The function  $f$  is said to be **bounded** in  $D$  if there exists a positive number  $M$  such that

$$|f(x, y)| \leq M$$

for all  $(x, y)$  in  $D$ .

- 2 Let  $f$  be defined and continuous on a closed rectangle  $R : a \leq x \leq b, c \leq y \leq d$ . Then,  $f$  is bounded in  $R$ .
- 3 Let  $f$  be defined on  $D$ , where  $D$  is either a domain or a closed domain of the  $xy$ - plane. The function  $f$  is said to satisfy **Lipschitz condition** (with respect to  $y$ ) in  $D$  if  $\exists$  a constant  $M > 0$  such that

# Definitions

- 1 Let  $f$  be a real function defined on  $D$ , where  $D$  is either a domain or a closed domain of the  $xy$  plane. The function  $f$  is said to be **bounded** in  $D$  if there exists a positive number  $M$  such that

$$|f(x, y)| \leq M$$

for all  $(x, y)$  in  $D$ .

- 2 Let  $f$  be defined and continuous on a closed rectangle  $R : a \leq x \leq b, c \leq y \leq d$ . Then,  $f$  is bounded in  $R$ .
- 3 Let  $f$  be defined on  $D$ , where  $D$  is either a domain or a closed domain of the  $xy$ - plane. The function  $f$  is said to satisfy **Lipschitz condition** (with respect to  $y$ ) in  $D$  if  $\exists$  a constant  $M > 0$  such that

$$|f(x, y_1) - f(x, y_2)| \leq M|y_1 - y_2|$$

# Definitions

- 1 Let  $f$  be a real function defined on  $D$ , where  $D$  is either a domain or a closed domain of the  $xy$  plane. The function  $f$  is said to be **bounded** in  $D$  if there exists a positive number  $M$  such that

$$|f(x, y)| \leq M$$

for all  $(x, y)$  in  $D$ .

- 2 Let  $f$  be defined and continuous on a closed rectangle  $R : a \leq x \leq b, c \leq y \leq d$ . Then,  $f$  is bounded in  $R$ .
- 3 Let  $f$  be defined on  $D$ , where  $D$  is either a domain or a closed domain of the  $xy$ - plane. The function  $f$  is said to satisfy **Lipschitz condition** (with respect to  $y$ ) in  $D$  if  $\exists$  a constant  $M > 0$  such that

$$|f(x, y_1) - f(x, y_2)| \leq M|y_1 - y_2|$$

for every  $(x, y_1), (x, y_2)$  which belong to  $D$ . The constant  $M$  is called the **Lipschitz constant**.

# Definitions

- 1 Let  $f$  be a real function defined on  $D$ , where  $D$  is either a domain or a closed domain of the  $xy$  plane. The function  $f$  is said to be **bounded** in  $D$  if there exists a positive number  $M$  such that

$$|f(x, y)| \leq M$$

for all  $(x, y)$  in  $D$ .

- 2 Let  $f$  be defined and continuous on a closed rectangle  $R : a \leq x \leq b, c \leq y \leq d$ . Then,  $f$  is bounded in  $R$ .
- 3 Let  $f$  be defined on  $D$ , where  $D$  is either a domain or a closed domain of the  $xy$ - plane. The function  $f$  is said to satisfy **Lipschitz condition** (with respect to  $y$ ) in  $D$  if  $\exists$  a constant  $M > 0$  such that

$$|f(x, y_1) - f(x, y_2)| \leq M|y_1 - y_2|$$

for every  $(x, y_1), (x, y_2)$  which belong to  $D$ . The constant  $M$  is called the **Lipschitz constant**.

# Understanding the Lipschitz condition - $y = g(x)$

Consider

$$|g(x_2) - g(x_1)| \leq M|x_2 - x_1| \quad \forall x_1, x_2 \text{ in the domain of } g.$$



# Understanding the Lipschitz condition - $y = g(x)$

Consider

$$|g(x_2) - g(x_1)| \leq M|x_2 - x_1| \quad \forall x_1, x_2 \text{ in the domain of } g.$$

This condition in the form  $\frac{|g(x_2) - g(x_1)|}{|x_2 - x_1|} \leq M$  can be interpreted as follows :

# Understanding the Lipschitz condition - $y = g(x)$

Consider

$$|g(x_2) - g(x_1)| \leq M|x_2 - x_1| \quad \forall x_1, x_2 \text{ in the domain of } g.$$

This condition in the form  $\frac{|g(x_2) - g(x_1)|}{|x_2 - x_1|} \leq M$  can be interpreted as follows :

At each point  $(a, g(a))$ , the entire graph of  $g$  lies between the lines

$$y = g(a) - M(x - a) \text{ \& } y = g(a) + M(x - a).$$

# Understanding the Lipschitz condition - $y = g(x)$


Consider

$$|g(x_2) - g(x_1)| \leq M|x_2 - x_1| \quad \forall x_1, x_2 \text{ in the domain of } g.$$

This condition in the form  $\frac{|g(x_2) - g(x_1)|}{|x_2 - x_1|} \leq M$  can be interpreted as follows :

At each point  $(a, g(a))$ , the entire graph of  $g$  lies between the lines

$$y = g(a) - M(x - a) \text{ \& } y = g(a) + M(x - a).$$



Example :  $x^2$  is Lipschitz in  $[1, 2]$ .

# Understanding Lipschitz condition - $z = f(x, y)$

- Let  $(x, y_1)$  and  $(x, y_2)$  be any two points in  $D$  having the same abscissa  $x$ .

# Understanding Lipschitz condition - $z = f(x, y)$

- Let  $(x, y_1)$  and  $(x, y_2)$  be any two points in  $D$  having the same abscissa  $x$ .
- Consider the corresponding points

$$P_1(x, y_1, f(x, y_1)) \text{ \& } P_2(x, y_2, f(x, y_2))$$

on the surface  $z = f(x, y)$ , and let  $\alpha$  ( $0 \leq \alpha \leq \pi/2$ ) denote the angle that the chord joining  $P_1$  and  $P_2$  makes with the  $xy$ - plane.

# Understanding Lipschitz condition - $z = f(x, y)$

- Let  $(x, y_1)$  and  $(x, y_2)$  be any two points in  $D$  having the same abscissa  $x$ .
- Consider the corresponding points

$$P_1(x, y_1, f(x, y_1)) \text{ \& } P_2(x, y_2, f(x, y_2))$$

on the surface  $z = f(x, y)$ , and let  $\alpha$  ( $0 \leq \alpha \leq \pi/2$ ) denote the angle that the chord joining  $P_1$  and  $P_2$  makes with the  $xy$ - plane.

- Then if the condition

$$|f(x, y_1) - f(x, y_2)| \leq M|y_1 - y_2|$$

holds in  $D$ , then  $\tan \alpha$  is bounded in absolute value.

# Understanding Lipschitz condition - $z = f(x, y)$

- Let  $(x, y_1)$  and  $(x, y_2)$  be any two points in  $D$  having the same abscissa  $x$ .
- Consider the corresponding points

$$P_1(x, y_1, f(x, y_1)) \text{ \& } P_2(x, y_2, f(x, y_2))$$

on the surface  $z = f(x, y)$ , and let  $\alpha$  ( $0 \leq \alpha \leq \pi/2$ ) denote the angle that the chord joining  $P_1$  and  $P_2$  makes with the  $xy$ - plane.

- Then if the condition

$$|f(x, y_1) - f(x, y_2)| \leq M|y_1 - y_2|$$

holds in  $D$ , then  $\tan \alpha$  is bounded in absolute value.

- That is, the chord joining  $P_1$  and  $P_2$  is bounded away from being perpendicular to the  $xy$ - plane.

# Understanding Lipschitz condition - $z = f(x, y)$

- Let  $(x, y_1)$  and  $(x, y_2)$  be any two points in  $D$  having the same abscissa  $x$ .
- Consider the corresponding points

$$P_1(x, y_1, f(x, y_1)) \text{ \& } P_2(x, y_2, f(x, y_2))$$

on the surface  $z = f(x, y)$ , and let  $\alpha$  ( $0 \leq \alpha \leq \pi/2$ ) denote the angle that the chord joining  $P_1$  and  $P_2$  makes with the  $xy$ - plane.

- Then if the condition

$$|f(x, y_1) - f(x, y_2)| \leq M|y_1 - y_2|$$

holds in  $D$ , then  $\tan \alpha$  is bounded in absolute value.

- That is, the chord joining  $P_1$  and  $P_2$  is bounded away from being perpendicular to the  $xy$ - plane.
- Further, this bound is independent of the points  $(x, y_1)$  and  $(x, y_2)$  belonging to  $D$ .



# Lipschitz condition $\implies$ Continuity ?

If  $f$  satisfies Lipschitz condition with respect to  $y$  in  $D$ , then for each fixed  $x$ , the resulting function of  $y$  is a continuous function of  $y$ , for all  $(x, y)$  in  $D$ .

# Lipschitz condition $\implies$ Continuity ?

If  $f$  satisfies Lipschitz condition with respect to  $y$  in  $D$ , then for each fixed  $x$ , the resulting function of  $y$  is a continuous function of  $y$ , for all  $(x, y)$  in  $D$ .

**Example :** Let  $f(x, y) = y + [x]$  where  $g(x) = [x]$  is the greatest integer function.

# Lipschitz condition $\implies$ Continuity ?

If  $f$  satisfies Lipschitz condition with respect to  $y$  in  $D$ , then for each fixed  $x$ , the resulting function of  $y$  is a continuous function of  $y$ , for all  $(x, y)$  in  $D$ .

**Example :** Let  $f(x, y) = y + [x]$  where  $g(x) = [x]$  is the greatest integer function. For fixed  $x$ ,

# Lipschitz condition $\implies$ Continuity ?

If  $f$  satisfies Lipschitz condition with respect to  $y$  in  $D$ , then for each fixed  $x$ , the resulting function of  $y$  is a continuous function of  $y$ , for all  $(x, y)$  in  $D$ .

**Example :** Let  $f(x, y) = y + [x]$  where  $g(x) = [x]$  is the greatest integer function. For fixed  $x$ ,

$$f(x, y_1) - f(x, y_2)$$

# Lipschitz condition $\implies$ Continuity ?

If  $f$  satisfies Lipschitz condition with respect to  $y$  in  $D$ , then for each fixed  $x$ , the resulting function of  $y$  is a continuous function of  $y$ , for all  $(x, y)$  in  $D$ .

**Example :** Let  $f(x, y) = y + [x]$  where  $g(x) = [x]$  is the greatest integer function. For fixed  $x$ ,

$$f(x, y_1) - f(x, y_2) = y_1 + [x] - y_2 - [x]$$

# Lipschitz condition $\implies$ Continuity ?

If  $f$  satisfies Lipschitz condition with respect to  $y$  in  $D$ , then for each fixed  $x$ , the resulting function of  $y$  is a continuous function of  $y$ , for all  $(x, y)$  in  $D$ .

**Example :** Let  $f(x, y) = y + [x]$  where  $g(x) = [x]$  is the greatest integer function. For fixed  $x$ ,

$$\begin{aligned} f(x, y_1) - f(x, y_2) &= y_1 + [x] - y_2 - [x] \\ &= y_1 - y_2 \end{aligned}$$

# Lipschitz condition $\implies$ Continuity ?

If  $f$  satisfies Lipschitz condition with respect to  $y$  in  $D$ , then for each fixed  $x$ , the resulting function of  $y$  is a continuous function of  $y$ , for all  $(x, y)$  in  $D$ .

**Example :** Let  $f(x, y) = y + [x]$  where  $g(x) = [x]$  is the greatest integer function. For fixed  $x$ ,

$$\begin{aligned} f(x, y_1) - f(x, y_2) &= y_1 + [x] - y_2 - [x] \\ &= y_1 - y_2 \end{aligned}$$

That is,  $|f(x, y_1) - f(x, y_2)| = |y_1 - y_2|$

# Lipschitz condition $\implies$ Continuity ?

If  $f$  satisfies Lipschitz condition with respect to  $y$  in  $D$ , then for each fixed  $x$ , the resulting function of  $y$  is a continuous function of  $y$ , for all  $(x, y)$  in  $D$ .

**Example :** Let  $f(x, y) = y + [x]$  where  $g(x) = [x]$  is the greatest integer function. For fixed  $x$ ,

$$\begin{aligned} f(x, y_1) - f(x, y_2) &= y_1 + [x] - y_2 - [x] \\ &= y_1 - y_2 \end{aligned}$$

That is,  $|f(x, y_1) - f(x, y_2)| = |y_1 - y_2| \leq 1 \cdot |y_1 - y_2|$



# Lipschitz condition $\implies$ Continuity ?

If  $f$  satisfies Lipschitz condition with respect to  $y$  in  $D$ , then for each fixed  $x$ , the resulting function of  $y$  is a continuous function of  $y$ , for all  $(x, y)$  in  $D$ .

**Example :** Let  $f(x, y) = y + [x]$  where  $g(x) = [x]$  is the greatest integer function. For fixed  $x$ ,

$$\begin{aligned} f(x, y_1) - f(x, y_2) &= y_1 + [x] - y_2 - [x] \\ &= y_1 - y_2 \end{aligned}$$

That is,  $|f(x, y_1) - f(x, y_2)| = |y_1 - y_2| \leq 1 \cdot |y_1 - y_2|$

But we know that  $f$  is **discontinuous** w.r.t.  $x$  for every integral value of  $x$ .

# Lipschitz condition $\implies$ Continuity ?

If  $f$  satisfies Lipschitz condition with respect to  $y$  in  $D$ , then for each fixed  $x$ , the resulting function of  $y$  is a continuous function of  $y$ , for all  $(x, y)$  in  $D$ .

**Example :** Let  $f(x, y) = y + [x]$  where  $g(x) = [x]$  is the greatest integer function. For fixed  $x$ ,

$$\begin{aligned} f(x, y_1) - f(x, y_2) &= y_1 + [x] - y_2 - [x] \\ &= y_1 - y_2 \end{aligned}$$

That is,  $|f(x, y_1) - f(x, y_2)| = |y_1 - y_2| \leq 1 \cdot |y_1 - y_2|$

But we know that  $f$  is **discontinuous** w.r.t.  $x$  for every integral value of  $x$ .

Note that the condition of Lipschitz continuity implies **nothing** concerning the continuity of  $f$  with respect to  $x$ .

Does Continuity w.r.t. second variable  $\implies$  Lipschitz condtn. w.r.t. second variable?

Continuity w.r.t. second variable DOES NOT imply Lipschitz condtn. w.r.t. second variable.

# Does Continuity w.r.t. second variable $\implies$ Lipschitz condtn. w.r.t. second variable?

Continuity w.r.t. second variable DOES NOT imply Lipschitz condtn. w.r.t. second variable.

Example : Consider  $f(x, y) = \sqrt{|y|}$ .

# Does Continuity w.r.t. second variable $\implies$ Lipschitz condtn. w.r.t. second variable?

Continuity w.r.t. second variable DOES NOT imply Lipschitz condtn. w.r.t. second variable.

Example : Consider  $f(x, y) = \sqrt{|y|}$ .

$f$  is continuous for all  $y$ .

# Does Continuity w.r.t. second variable $\implies$ Lipschitz condtn. w.r.t. second variable?

Continuity w.r.t. second variable DOES NOT imply Lipschitz condtn. w.r.t. second variable.

Example : Consider  $f(x, y) = \sqrt{|y|}$ .

$f$  is continuous for all  $y$ .

Note that  $f$  doesn't satisfy Lipschitz condition in any region that includes  $y = 0$  as for  $y_1 = 0$ ,  $y_2 > 0$ , we have

# Does Continuity w.r.t. second variable $\implies$ Lipschitz condtn. w.r.t. second variable?

Continuity w.r.t. second variable DOES NOT imply Lipschitz condtn. w.r.t. second variable.

Example : Consider  $f(x, y) = \sqrt{|y|}$ .

$f$  is continuous for all  $y$ .

Note that  $f$  doesn't satisfy Lipschitz condition in any region that includes  $y = 0$  as for  $y_1 = 0$ ,  $y_2 > 0$ , we have

$$\frac{|f(x, y_1) - f(x, y_2)|}{|y_1 - y_2|} =$$

# Does Continuity w.r.t. second variable $\implies$ Lipschitz condtn. w.r.t. second variable?

Continuity w.r.t. second variable DOES NOT imply Lipschitz condtn. w.r.t. second variable.

Example : Consider  $f(x, y) = \sqrt{|y|}$ .

$f$  is continuous for all  $y$ .

Note that  $f$  doesn't satisfy Lipschitz condition in any region that includes  $y = 0$  as for  $y_1 = 0$ ,  $y_2 > 0$ , we have

$$\frac{|f(x, y_1) - f(x, y_2)|}{|y_1 - y_2|} = \frac{\sqrt{y_2}}{|y_2|}$$



# Does Continuity w.r.t. second variable $\implies$ Lipschitz condtn. w.r.t. second variable?

Continuity w.r.t. second variable DOES NOT imply Lipschitz condtn. w.r.t. second variable.

Example : Consider  $f(x, y) = \sqrt{|y|}$ .

$f$  is continuous for all  $y$ .

Note that  $f$  doesn't satisfy Lipschitz condition in any region that includes  $y = 0$  as for  $y_1 = 0$ ,  $y_2 > 0$ , we have

$$\frac{|f(x, y_1) - f(x, y_2)|}{|y_1 - y_2|} = \frac{\sqrt{y_2}}{|y_2|} = \frac{1}{\sqrt{y_2}}$$

# Does Continuity w.r.t. second variable $\implies$ Lipschitz condtn. w.r.t. second variable?

Continuity w.r.t. second variable DOES NOT imply Lipschitz condtn. w.r.t. second variable.

Example : Consider  $f(x, y) = \sqrt{|y|}$ .

$f$  is continuous for all  $y$ .

Note that  $f$  doesn't satisfy Lipschitz condition in any region that includes  $y = 0$  as for  $y_1 = 0$ ,  $y_2 > 0$ , we have

$$\frac{|f(x, y_1) - f(x, y_2)|}{|y_1 - y_2|} = \frac{\sqrt{y_2}}{|y_2|} = \frac{1}{\sqrt{y_2}}$$

which can be made as large as we want by making  $y_2$  smaller.

# Does Continuity w.r.t. second variable $\implies$ Lipschitz condtn. w.r.t. second variable?

Continuity w.r.t. second variable DOES NOT imply Lipschitz condtn. w.r.t. second variable.

Example : Consider  $f(x, y) = \sqrt{|y|}$ .

$f$  is continuous for all  $y$ .

Note that  $f$  doesn't satisfy Lipschitz condition in any region that includes  $y = 0$  as for  $y_1 = 0$ ,  $y_2 > 0$ , we have

$$\frac{|f(x, y_1) - f(x, y_2)|}{|y_1 - y_2|} = \frac{\sqrt{y_2}}{|y_2|} = \frac{1}{\sqrt{y_2}}$$

which can be made as large as we want by making  $y_2$  smaller. The Lipschitz condition requires that the quotient should be bounded by a fixed constant  $K$ .

Does Continuity w.r.t. second variable  $\implies$  Lipschitz condtn. w.r.t. second variable?

Continuity w.r.t. second variable  $\not\Rightarrow$  Lipschitz condtn. w.r.t. second variable.

# Does Continuity w.r.t. second variable $\implies$ Lipschitz condtn. w.r.t. second variable?

Continuity w.r.t. second variable  $\not\Rightarrow$  Lipschitz condtn. w.r.t. second variable.

Example : Consider  $f(x, y) = \sqrt{|y|}$ .

# Does Continuity w.r.t. second variable $\implies$ Lipschitz condtn. w.r.t. second variable?

Continuity w.r.t. second variable  $\not\Rightarrow$  Lipschitz condtn. w.r.t. second variable.

**Example :** Consider  $f(x, y) = \sqrt{|y|}$ .

$f$  is continuous for all  $y$ .

# Does Continuity w.r.t. second variable $\implies$ Lipschitz condtn. w.r.t. second variable?

Continuity w.r.t. second variable  $\nRightarrow$  Lipschitz condtn. w.r.t. second variable.

**Example :** Consider  $f(x, y) = \sqrt{|y|}$ .

$f$  is continuous for all  $y$ .

Note that  $f$  doesn't satisfy Lipschitz condition in any region that includes  $y = 0$  as for  $y_1 = 0$ ,  $y_2 > 0$ , we have

# Does Continuity w.r.t. second variable $\implies$ Lipschitz condtn. w.r.t. second variable?

Continuity w.r.t. second variable  $\not\Rightarrow$  Lipschitz condtn. w.r.t. second variable.

**Example :** Consider  $f(x, y) = \sqrt{|y|}$ .

$f$  is continuous for all  $y$ .

Note that  $f$  doesn't satisfy Lipschitz condition in any region that includes  $y = 0$  as for  $y_1 = 0$ ,  $y_2 > 0$ , we have

$$\frac{|f(x, y_1) - f(x, y_2)|}{|y_1 - y_2|} =$$



# Does Continuity w.r.t. second variable $\implies$ Lipschitz condtn. w.r.t. second variable?

Continuity w.r.t. second variable  $\not\Rightarrow$  Lipschitz condtn. w.r.t. second variable.

**Example :** Consider  $f(x, y) = \sqrt{|y|}$ .

$f$  is continuous for all  $y$ .

Note that  $f$  doesn't satisfy Lipschitz condition in any region that includes  $y = 0$  as for  $y_1 = 0$ ,  $y_2 > 0$ , we have

$$\frac{|f(x, y_1) - f(x, y_2)|}{|y_1 - y_2|} = \frac{\sqrt{y_2}}{|y_2|}$$

# Does Continuity w.r.t. second variable $\implies$ Lipschitz condtn. w.r.t. second variable?

Continuity w.r.t. second variable  $\not\Rightarrow$  Lipschitz condtn. w.r.t. second variable.

**Example :** Consider  $f(x, y) = \sqrt{|y|}$ .

$f$  is continuous for all  $y$ .

Note that  $f$  doesn't satisfy Lipschitz condition in any region that includes  $y = 0$  as for  $y_1 = 0$ ,  $y_2 > 0$ , we have

$$\frac{|f(x, y_1) - f(x, y_2)|}{|y_1 - y_2|} = \frac{\sqrt{y_2}}{|y_2|} = \frac{1}{\sqrt{y_2}}$$

# Does Continuity w.r.t. second variable $\implies$ Lipschitz condtn. w.r.t. second variable?

Continuity w.r.t. second variable  $\not\Rightarrow$  Lipschitz condtn. w.r.t. second variable.

**Example :** Consider  $f(x, y) = \sqrt{|y|}$ .

$f$  is continuous for all  $y$ .

Note that  $f$  doesn't satisfy Lipschitz condition in any region that includes  $y = 0$  as for  $y_1 = 0$ ,  $y_2 > 0$ , we have

$$\frac{|f(x, y_1) - f(x, y_2)|}{|y_1 - y_2|} = \frac{\sqrt{y_2}}{|y_2|} = \frac{1}{\sqrt{y_2}}$$

which can be made as large as we want by making  $y_2$  smaller.

# Does Continuity w.r.t. second variable $\implies$ Lipschitz condtn. w.r.t. second variable?

Continuity w.r.t. second variable  $\not\Rightarrow$  Lipschitz condtn. w.r.t. second variable.

**Example :** Consider  $f(x, y) = \sqrt{|y|}$ .

$f$  is continuous for all  $y$ .

Note that  $f$  doesn't satisfy Lipschitz condition in any region that includes  $y = 0$  as for  $y_1 = 0$ ,  $y_2 > 0$ , we have

$$\frac{|f(x, y_1) - f(x, y_2)|}{|y_1 - y_2|} = \frac{\sqrt{y_2}}{|y_2|} = \frac{1}{\sqrt{y_2}}$$

which can be made as large as we want by making  $y_2$  smaller. The Lipschitz condition requires that the quotient should be bounded by a fixed constant  $K$ .

Result : If  $f$  is such that  $\frac{\partial f}{\partial y}$  exists and is bounded for all  $(x, y) \in D$ , then  $f$  satisfies Lipschitz condition w.r.t.  $y$  in  $D$ , where the Lipschitz constant

$$M = l.u.b._{(x,y) \in D} \left| \frac{\partial f}{\partial y}(x, y) \right|.$$

Result : If  $f$  is such that  $\frac{\partial f}{\partial y}$  exists and is bounded for all  $(x, y) \in D$ , then  $f$  satisfies Lipschitz condition w.r.t.  $y$  in  $D$ , where the Lipschitz constant

$$M = l.u.b._{(x,y) \in D} \left| \frac{\partial f}{\partial y}(x, y) \right|.$$

Proof : Mean value theorem

$\implies$

# Sufficiency

Result : If  $f$  is such that  $\frac{\partial f}{\partial y}$  exists and is bounded for all  $(x, y) \in D$ , then  $f$  satisfies Lipschitz condition w.r.t.  $y$  in  $D$ , where the Lipschitz constant

$$M = l.u.b._{(x,y) \in D} \left| \frac{\partial f}{\partial y}(x, y) \right|.$$

Proof : Mean value theorem

$$\implies f(x, y_1) - f(x, y_2) = (y_1 - y_2) \frac{\partial f}{\partial y}(x, \xi),$$

Result : If  $f$  is such that  $\frac{\partial f}{\partial y}$  exists and is bounded for all  $(x, y) \in D$ , then  $f$  satisfies Lipschitz condition w.r.t.  $y$  in  $D$ , where the Lipschitz constant

$$M = l.u.b._{(x,y) \in D} \left| \frac{\partial f}{\partial y}(x, y) \right|.$$

Proof : Mean value theorem

$$\implies f(x, y_1) - f(x, y_2) = (y_1 - y_2) \frac{\partial f}{\partial y}(x, \xi), \quad \xi \in (y_1, y_2).$$



Result : If  $f$  is such that  $\frac{\partial f}{\partial y}$  exists and is bounded for all  $(x, y) \in D$ , then  $f$  satisfies Lipschitz condition w.r.t.  $y$  in  $D$ , where the Lipschitz constant

$$M = l.u.b._{(x,y) \in D} \left| \frac{\partial f}{\partial y}(x, y) \right|.$$

Proof : Mean value theorem

$$\implies f(x, y_1) - f(x, y_2) = (y_1 - y_2) \frac{\partial f}{\partial y}(x, \xi), \quad \xi \in (y_1, y_2).$$

$$|f(x, y_1) - f(x, y_2)|$$

Result : If  $f$  is such that  $\frac{\partial f}{\partial y}$  exists and is bounded for all  $(x, y) \in D$ , then  $f$  satisfies Lipschitz condition w.r.t.  $y$  in  $D$ , where the Lipschitz constant

$$M = l.u.b._{(x,y) \in D} \left| \frac{\partial f}{\partial y}(x, y) \right|.$$

Proof : Mean value theorem

$$\implies f(x, y_1) - f(x, y_2) = (y_1 - y_2) \frac{\partial f}{\partial y}(x, \xi), \quad \xi \in (y_1, y_2).$$

$$|f(x, y_1) - f(x, y_2)| = |y_1 - y_2| \left| \frac{\partial f}{\partial y}(x, \xi) \right|$$

Result : If  $f$  is such that  $\frac{\partial f}{\partial y}$  exists and is bounded for all  $(x, y) \in D$ , then  $f$  satisfies Lipschitz condition w.r.t.  $y$  in  $D$ , where the Lipschitz constant

$$M = l.u.b._{(x,y) \in D} \left| \frac{\partial f}{\partial y}(x, y) \right|.$$

Proof : Mean value theorem

$$\implies f(x, y_1) - f(x, y_2) = (y_1 - y_2) \frac{\partial f}{\partial y}(x, \xi), \quad \xi \in (y_1, y_2).$$

$$\begin{aligned} |f(x, y_1) - f(x, y_2)| &= |y_1 - y_2| \left| \frac{\partial f}{\partial y}(x, \xi) \right| \\ &\leq |y_1 - y_2| \, l.u.b._{(x,y) \in D} \left| \frac{\partial f}{\partial y}(x, y) \right|. \end{aligned}$$

Result : If  $f$  is such that  $\frac{\partial f}{\partial y}$  exists and is bounded for all  $(x, y) \in D$ , then  $f$  satisfies Lipschitz condition w.r.t.  $y$  in  $D$ , where the Lipschitz constant

$$M = l.u.b._{(x,y) \in D} \left| \frac{\partial f}{\partial y}(x, y) \right|.$$

Proof : Mean value theorem

$$\implies f(x, y_1) - f(x, y_2) = (y_1 - y_2) \frac{\partial f}{\partial y}(x, \xi), \quad \xi \in (y_1, y_2).$$

$$\begin{aligned} |f(x, y_1) - f(x, y_2)| &= |y_1 - y_2| \left| \frac{\partial f}{\partial y}(x, \xi) \right| \\ &\leq |y_1 - y_2| \, l.u.b._{(x,y) \in D} \left| \frac{\partial f}{\partial y}(x, y) \right|. \end{aligned}$$

That is,  $f$  satisfies Lipschitz condition.

# Example

Consider

$$f(x, y) = y^2 \text{ defined in } D : |x| \leq a, |y| \leq b.$$

# Example

Consider

$$f(x, y) = y^2 \text{ defined in } D : |x| \leq a, |y| \leq b.$$

$f_y = 2y$  is bounded in  $D$ .

# Example

Consider

$$f(x, y) = y^2 \text{ defined in } D : |x| \leq a, |y| \leq b.$$

$f_y = 2y$  is bounded in  $D$ . The Lipschitz constant is

$$M =$$

# Example

Consider

$$f(x, y) = y^2 \text{ defined in } D : |x| \leq a, |y| \leq b.$$

$f_y = 2y$  is bounded in  $D$ . The Lipschitz constant is

$$M = \text{l.u.b.}_{(x,y) \in D} \left| \frac{\partial f}{\partial y}(x, y) \right| =$$



# Example

Consider

$$f(x, y) = y^2 \text{ defined in } D : |x| \leq a, |y| \leq b.$$

$f_y = 2y$  is bounded in  $D$ . The Lipschitz constant is

$$M = \text{l.u.b.}_{(x,y) \in D} \left| \frac{\partial f}{\partial y}(x, y) \right| = \text{l.u.b.}_{(x,y) \in D} |2y|$$

# Example

Consider

$$f(x, y) = y^2 \text{ defined in } D : |x| \leq a, |y| \leq b.$$

$f_y = 2y$  is bounded in  $D$ . The Lipschitz constant is

$$M = \text{l.u.b.}_{(x,y) \in D} \left| \frac{\partial f}{\partial y}(x, y) \right| = \text{l.u.b.}_{(x,y) \in D} |2y| = 2b.$$

# Example

Consider

$$f(x, y) = y^2 \text{ defined in } D : |x| \leq a, |y| \leq b.$$

$f_y = 2y$  is bounded in  $D$ . The Lipschitz constant is

$$M = \text{l.u.b.}_{(x,y) \in D} \left| \frac{\partial f}{\partial y}(x, y) \right| = \text{l.u.b.}_{(x,y) \in D} |2y| = 2b.$$

(Verify Lipschitz condition directly! )

# Bounded derivative - sufficient condition

Consider

$$f(x, y) = x|y| \quad \text{defined in } D : |x| \leq a, |y| \leq b.$$

# Bounded derivative - sufficient condition

Consider

$$f(x, y) = x|y| \quad \text{defined in } D : |x| \leq a, |y| \leq b.$$

$\frac{\partial f}{\partial y}$  doesn't exist for any point  $(x, 0) \in D$ .

# Bounded derivative - sufficient condition

Consider

$$f(x, y) = x|y| \quad \text{defined in } D : |x| \leq a, |y| \leq b.$$

$\frac{\partial f}{\partial y}$  doesn't exist for any point  $(x, 0) \in D$ . (Why?)

Now  $f$  satisfies Lipschitz condition :

# Bounded derivative - sufficient condition

Consider

$$f(x, y) = x|y| \quad \text{defined in } D : |x| \leq a, |y| \leq b.$$

$\frac{\partial f}{\partial y}$  doesn't exist for any point  $(x, 0) \in D$ . (Why?)

Now  $f$  satisfies Lipschitz condition :

$$|f(x, y_1) - f(x, y_2)| =$$

# Bounded derivative - sufficient condition

Consider

$$f(x, y) = x|y| \quad \text{defined in } D : |x| \leq a, |y| \leq b.$$

$\frac{\partial f}{\partial y}$  doesn't exist for any point  $(x, 0) \in D$ . (Why?)

Now  $f$  satisfies Lipschitz condition :

$$|f(x, y_1) - f(x, y_2)| = |x|y_1| - x|y_2||$$



# Bounded derivative - sufficient condition

Consider

$$f(x, y) = x|y| \quad \text{defined in } D : |x| \leq a, |y| \leq b.$$

$\frac{\partial f}{\partial y}$  doesn't exist for any point  $(x, 0) \in D$ . (Why?)

Now  $f$  satisfies Lipschitz condition :

$$\begin{aligned} |f(x, y_1) - f(x, y_2)| &= |x|y_1| - x|y_2|| \\ &= |x| \left| |y_1| - |y_2| \right| \end{aligned}$$

# Bounded derivative - sufficient condition

Consider

$$f(x, y) = x|y| \quad \text{defined in } D : |x| \leq a, |y| \leq b.$$

$\frac{\partial f}{\partial y}$  doesn't exist for any point  $(x, 0) \in D$ . (Why?)

Now  $f$  satisfies Lipschitz condition :

$$\begin{aligned} |f(x, y_1) - f(x, y_2)| &= |x|y_1| - x|y_2|| \\ &= |x| \left| |y_1| - |y_2| \right| \\ &\leq |x| |y_1 - y_2| \end{aligned}$$

# Bounded derivative - sufficient condition

Consider

$$f(x, y) = x|y| \quad \text{defined in } D : |x| \leq a, |y| \leq b.$$

$\frac{\partial f}{\partial y}$  doesn't exist for any point  $(x, 0) \in D$ . (Why?)

Now  $f$  satisfies Lipschitz condition :

$$\begin{aligned} |f(x, y_1) - f(x, y_2)| &= |x|y_1| - x|y_2|| \\ &= |x| \left| |y_1| - |y_2| \right| \\ &\leq |x| |y_1 - y_2| \\ &\leq a|y_1 - y_2| \end{aligned}$$

# Bounded derivative - sufficient condition

Consider

$$f(x, y) = x|y| \text{ defined in } D : |x| \leq a, |y| \leq b.$$

$\frac{\partial f}{\partial y}$  doesn't exist for any point  $(x, 0) \in D$ . (Why?)

Now  $f$  satisfies Lipschitz condition :

$$\begin{aligned} |f(x, y_1) - f(x, y_2)| &= |x|y_1| - x|y_2|| \\ &= |x| \left| |y_1| - |y_2| \right| \\ &\leq |x| |y_1 - y_2| \\ &\leq a|y_1 - y_2| \end{aligned}$$

Existence of bounded derivative  $f_y$  is a sufficient condition for Lipschitz condition to hold true (not necessary).

# Existence - Uniqueness Theorem

Let  $R$  be a rectangle containing  $(x_0, y_0)$  in the domain  $D$ ,

- $f(x, y)$  be **continuous** at all points  $(x, y)$  in  $R : |x - x_0| < a, |y - y_0| < b$  and

# Existence - Uniqueness Theorem

Let  $R$  be a rectangle containing  $(x_0, y_0)$  in the domain  $D$ ,

- $f(x, y)$  be **continuous** at all points  $(x, y)$  in  $R : |x - x_0| < a, |y - y_0| < b$  and
- **bounded** in  $R$ , that is,  $|f(x, y)| \leq K \quad \forall (x, y) \in R$ .

# Existence - Uniqueness Theorem

Let  $R$  be a rectangle containing  $(x_0, y_0)$  in the domain  $D$ ,

- $f(x, y)$  be **continuous** at all points  $(x, y)$  in  $R : |x - x_0| < a, |y - y_0| < b$  and
- **bounded** in  $R$ , that is,  $|f(x, y)| \leq K \quad \forall (x, y) \in R$ .

Then, the IVP  $y' = f(x, y), y(x_0) = y_0$  has **at least one solution**  $y(x)$  defined for all  $x$  in the interval  $|x - x_0| < \alpha$ , where

$$\alpha = \min \left\{ a, \frac{b}{K} \right\}.$$

# Existence - Uniqueness Theorem

Let  $R$  be a rectangle containing  $(x_0, y_0)$  in the domain  $D$ ,

- $f(x, y)$  be **continuous** at all points  $(x, y)$  in  $R : |x - x_0| < a, |y - y_0| < b$  and
- **bounded** in  $R$ , that is,  $|f(x, y)| \leq K \quad \forall (x, y) \in R$ .

Then, the IVP  $y' = f(x, y), y(x_0) = y_0$  has **at least one solution**  $y(x)$  defined for all  $x$  in the interval  $|x - x_0| < \alpha$ , where

$$\alpha = \min \left\{ a, \frac{b}{K} \right\}.$$

In addition to the above conditions, if  $f$  satisfies the **Lipschitz condition** with respect to  $y$  in  $R$ , that is,



# Existence - Uniqueness Theorem

Let  $R$  be a rectangle containing  $(x_0, y_0)$  in the domain  $D$ ,

- $f(x, y)$  be **continuous** at all points  $(x, y)$  in  $R : |x - x_0| < a, |y - y_0| < b$  and
- **bounded** in  $R$ , that is,  $|f(x, y)| \leq K \quad \forall (x, y) \in R$ .

Then, the IVP  $y' = f(x, y), y(x_0) = y_0$  has **at least one solution**  $y(x)$  defined for all  $x$  in the interval  $|x - x_0| < \alpha$ , where

$$\alpha = \min \left\{ a, \frac{b}{K} \right\}.$$

In addition to the above conditions, if  $f$  satisfies the **Lipschitz condition** with respect to  $y$  in  $R$ , that is,

$$|f(x, y_1) - f(x, y_2)| \leq M|y_1 - y_2| \quad \forall (x, y_1), (x, y_2) \text{ in } R,$$

# Existence - Uniqueness Theorem

Let  $R$  be a rectangle containing  $(x_0, y_0)$  in the domain  $D$ ,

- $f(x, y)$  be **continuous** at all points  $(x, y)$  in  $R : |x - x_0| < a, |y - y_0| < b$  and
- **bounded** in  $R$ , that is,  $|f(x, y)| \leq K \quad \forall (x, y) \in R$ .

Then, the IVP  $y' = f(x, y), y(x_0) = y_0$  has **at least one solution**  $y(x)$  defined for all  $x$  in the interval  $|x - x_0| < \alpha$ , where

$$\alpha = \min \left\{ a, \frac{b}{K} \right\}.$$

In addition to the above conditions, if  $f$  satisfies the **Lipschitz condition** with respect to  $y$  in  $R$ , that is,

$$|f(x, y_1) - f(x, y_2)| \leq M|y_1 - y_2| \quad \forall (x, y_1), (x, y_2) \text{ in } R,$$

then, the solution  $y(x)$  defined at least for all  $x$  in the interval  $|x - x_0| < \alpha$ , with  $\alpha$  defined above is **unique**<sup>1</sup>.

<sup>1</sup>Existence - Peano, Existence & uniqueness -Picard

# Existence - Uniqueness Theorem

Let  $R$  be a rectangle containing  $(x_0, y_0)$  in the domain  $D$ ,

- $f(x, y)$  be **continuous** at all points  $(x, y)$  in  $R : |x - x_0| < a, |y - y_0| < b$  and
- **bounded** in  $R$ , that is,  $|f(x, y)| \leq K \quad \forall (x, y) \in R$ .

Then, the IVP  $y' = f(x, y), y(x_0) = y_0$  has **at least one solution**  $y(x)$  defined for all  $x$  in the interval  $|x - x_0| < \alpha$ , where

$$\alpha = \min \left\{ a, \frac{b}{K} \right\}.$$

In addition to the above conditions, if  $f$  satisfies the **Lipschitz condition** with respect to  $y$  in  $R$ , that is,

$$|f(x, y_1) - f(x, y_2)| \leq M|y_1 - y_2| \quad \forall (x, y_1), (x, y_2) \text{ in } R,$$

then, the solution  $y(x)$  defined at least for all  $x$  in the interval  $|x - x_0| < \alpha$ , with  $\alpha$  defined above is **unique**<sup>1</sup>.

<sup>1</sup>Existence - Peano, Existence & uniqueness -Picard