

Assignment 6

Group 25

September 19, 2018

1.1

$$A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 3 \\ 6 & -6 & -6 \end{bmatrix}$$

compute eigen values using $|A - \lambda I| = 0$

Eigen values(λ) = 2, 0, -3

Algebraic multiplicity corresponding to $\lambda=(2, 0, -3)$ are 1, 1, 1

Finding Eigen vector for $\lambda_1 = 2$:

$$(A - \lambda_1 I)X_1 = 0$$

$$\begin{bmatrix} 2-2 & 0 & 0 \\ -1 & 3-2 & 3 \\ 6 & -6 & -6-2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = 0$$

reducing to row echelon form

$$\begin{bmatrix} 0 & 0 & 0 \\ -1 & 1 & 3 \\ 6 & -6 & -8 \end{bmatrix}$$

$$R_3 \rightarrow R_3 + 6R_2$$

$$\begin{bmatrix} 0 & 0 & 0 \\ -1 & 1 & 3 \\ 0 & 0 & 10 \end{bmatrix}$$

$$R_1 \leftrightarrow R_3$$

$$\begin{bmatrix} 0 & 0 & 10 \\ -1 & 1 & 3 \\ 0 & 0 & 0 \end{bmatrix}$$

$$R_1 \leftrightarrow R_2$$

$$\begin{bmatrix} -1 & 1 & 3 \\ 0 & 0 & 10 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\text{Eigen vector } v_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$$

Geometric multiplicity for $(\lambda=2) = 1$

Finding Eigen vector for $\lambda_2 = 2$:

$$(A - \lambda_2 I)X_2 = 0$$

$$\begin{bmatrix} 2-0 & 0 & 0 \\ -1 & 3-0 & 3 \\ 6 & -6 & -6-0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = 0$$

reducing to row echelon form

$$\begin{bmatrix} 2 & 0 & 0 \\ 0 & 3 & 3 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\text{Eigen vector } v_2 = \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix}$$

Geometric multiplicity for $(\lambda=0) = 1$

Finding Eigen vector for $\lambda_3 = -3$:

$$(A - \lambda_3 I)X_3 = 0$$

$$\begin{bmatrix} 2+3 & 0 & 0 \\ -1 & 3+3 & 3 \\ 6 & -6 & -6+3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = 0$$

reducing to row echelon form

$$\begin{bmatrix} 5 & 0 & 0 \\ 0 & 6 & 3 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\text{Eigen vector } v_3 = \begin{bmatrix} 0 \\ 1 \\ -2 \end{bmatrix}$$

Geometric multiplicity for $(\lambda=-3) = 1$

for every value of λ , $AM = GM$, so matrix A is diagonalizable.

$$X = \begin{bmatrix} 1 & 0 & 0 \\ 1 & -1 & 1 \\ 0 & 1 & -2 \end{bmatrix}$$

$$X^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 2 & -2 & -1 \\ 1 & -1 & 1 \end{bmatrix}$$

$$\Lambda = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -3 \end{bmatrix}$$

$$AX = X\Lambda$$

$$A = X\Lambda X^{-1}$$

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 1 & -1 & 1 \\ 0 & 1 & -2 \end{bmatrix} \begin{bmatrix} 2 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -3 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 2 & -2 & -1 \\ 1 & -1 & 1 \end{bmatrix}$$

1.2

$$A = \begin{bmatrix} 1 & 4 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 5 & 3 & 1 & 1 \\ 4 & 7 & 2 & 2 \end{bmatrix}$$

compute eigen values using $|A - \lambda I| = 0$

Eigen values(λ) = 0, 1, 2, 3

Algebraic multiplicity corresponding to $\lambda = (0, 1, 2, 3)$ are 1, 1, 1, 1

Finding Eigen vector for $\lambda_1 = 0$:

$$(A - \lambda_1 I)X_1 = 0$$

$$\begin{bmatrix} 1-1 & 4 & 0 & 0 \\ 0 & 2-1 & 0 & 0 \\ 5 & 3 & 1-1 & 1 \\ 4 & 7 & 2 & 2-1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = 0$$

reducing to row echelon form

$$\begin{bmatrix} 1 & 4 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\text{Eigen vector } v_1 = \begin{bmatrix} 0 \\ 0 \\ -1 \\ 1 \end{bmatrix}$$

Geometric multiplicity for ($\lambda=0$) = 1

Finding Eigen vector for $\lambda_2 = 1$:

$$(A - \lambda_2 I)X_2 = 0$$

$$\begin{bmatrix} 1-2 & 4 & 0 & 0 \\ 0 & 2-2 & 0 & 0 \\ 5 & 3 & 1-2 & 1 \\ 4 & 7 & 2 & 2-2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = 0$$

reducing to row echelon form

$$\begin{bmatrix} 4 & 7 & 2 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 10 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\text{Eigen vector } v_2 = \begin{bmatrix} 2 \\ 0 \\ 1 \\ -10 \end{bmatrix}$$

Geometric multiplicity for $(\lambda=1) = 1$

Finding Eigen vector for $\lambda_3 = 2$:

$$(A - \lambda_3 I)X_3 = 0$$

$$\begin{bmatrix} 1-3 & 4 & 0 & 0 \\ 0 & 2-3 & 0 & 0 \\ 5 & 3 & 1-3 & 1 \\ 4 & 7 & 2 & 2-3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = 0$$

reducing to row echelon form

$$\begin{bmatrix} -1 & 4 & 0 & 0 \\ 0 & 23 & 2 & 0 \\ 0 & 0 & -3 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\text{Eigen vector } v_3 = \begin{bmatrix} 8 \\ 2 \\ -23 \\ -69 \end{bmatrix}$$

Geometric multiplicity for $(\lambda=2) = 1$

Finding Eigen vector for $\lambda_4 = 3$:

$$(A - \lambda_4 I)X_4 = 0$$

$$\begin{bmatrix} 1-0 & 4 & 0 & 0 \\ 0 & 2-0 & 0 & 0 \\ 5 & 3 & 1-0 & 1 \\ 4 & 7 & 2 & 2-0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = 0$$

reducing to row echelon form

$$\begin{bmatrix} -2 & 4 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -2 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Eigen vector $v_4 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 2 \end{bmatrix}$

Geometric multicity for $(\lambda=3) = 1$

for every value of λ , AM = GM, so matrix A is diagonalizable.

$$X = \begin{bmatrix} 0 & 2 & 8 & 0 \\ 0 & 0 & 2 & 0 \\ -1 & 1 & -23 & 1 \\ 1 & -10 & -69 & 2 \end{bmatrix}$$

$$X^{-1} = \begin{bmatrix} 2 & -25/6 & -2/3 & 1/3 \\ 1/2 & -2 & 0 & 0 \\ 0 & 1/2 & 0 & 0 \\ 3/2 & 28/3 & 1/3 & 1/3 \end{bmatrix}$$

$$\Lambda = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \end{bmatrix}$$

$$AX = X\Lambda$$

$$A = X\Lambda X^{-1}$$

$$A = \begin{bmatrix} 0 & 2 & 8 & 0 \\ 0 & 0 & 2 & 0 \\ -1 & 1 & -23 & 1 \\ 1 & -10 & -69 & 2 \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \end{bmatrix} \begin{bmatrix} 2 & -25/6 & -2/3 & 1/3 \\ 1/2 & -2 & 0 & 0 \\ 0 & 1/2 & 0 & 0 \\ 3/2 & 28/3 & 1/3 & 1/3 \end{bmatrix}$$

1.3

$$A = \begin{bmatrix} -2 & 0 & 0 & 0 \\ 0 & -2 & 0 & 0 \\ 24 & -12 & 2 & 0 \\ 0 & 0 & 0 & 2 \end{bmatrix}$$

compute eigen values using $|A - \lambda I| = 0$

Eigen values(λ) = 2, 2, -2, -2

AM($\lambda=2$) = 2

AM($\lambda=-2$) = 2

Finding Eigen vector for $\lambda_1 = 2$:

$$(A - \lambda_1 I)X_1 = 0$$

$$A = \begin{bmatrix} -4 & 0 & 0 & 0 \\ 0 & -4 & 0 & 0 \\ 24 & -12 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = 0$$

reducing to row echelon form

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\text{Eigen vector } v_1 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}$$

Geometric multiplicity for ($\lambda=2$) = 2

Finding Eigen vector for $\lambda_1 = -2$:

$$(A - \lambda_2 I)X_2 = 0$$

$$A = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 24 & -12 & 4 & 0 \\ 0 & 0 & 0 & 4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = 0$$

reducing to row echelon form

$$\begin{bmatrix} 6 & -3 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\text{Eigen vector } v_2 = \begin{bmatrix} 2 \\ 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ -1 \\ 0 \end{bmatrix}$$

Geometric multiplicity for $(\lambda=-2) = 2$

for every value of λ , AM = GM, so matrix A is diagonalizable.

$$X = \begin{bmatrix} 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 6 \\ 1 & 0 & 1 & -1 \\ 0 & 1 & 0 & 0 \end{bmatrix}$$

$$X^{-1} = \begin{bmatrix} -1/2 & 1/6 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1/2 & 0 & 0 & 0 \\ 0 & 1/6 & 0 & 0 \end{bmatrix}$$

$$\Lambda = \begin{bmatrix} 2 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & -2 \end{bmatrix}$$

$$AX = X\Lambda$$

$$A = X\Lambda X^{-1}$$

$$A = \begin{bmatrix} 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 6 \\ 1 & 0 & 1 & -1 \\ 0 & 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} 2 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & -2 \end{bmatrix} \begin{bmatrix} -1/2 & 1/6 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1/2 & 0 & 0 & 0 \\ 0 & 1/6 & 0 & 0 \end{bmatrix}$$

2.1

$$A = \begin{bmatrix} 7 & 1 \\ 5 & 5 \\ 0 & 0 \end{bmatrix}, A = U_{m \times m} \Sigma_{m \times n} V_{n \times n}$$

$$AA^T = \begin{bmatrix} 7 & 1 \\ 5 & 5 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 7 & 5 & 0 \\ 1 & 5 & 0 \end{bmatrix} = \begin{bmatrix} 50 & 40 & 0 \\ 40 & 50 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Eigen value for AA^T ,

$\lambda = 0, 10, 90$

Singular values, $\sigma_1 = \sqrt{10}, \sigma_2 = \sqrt{90}$

$$\Sigma_{3 \times 2} = \begin{bmatrix} \sqrt{10} & 0 \\ 0 & \sqrt{90} \\ 0 & 0 \end{bmatrix}$$

$$(AA^T - \lambda I) = 0$$

for $\lambda = 0$,

Reduced echelon form is

$$\begin{bmatrix} 5 & 4 & 0 \\ 0 & 9/5 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\text{Eigen vector } v_1 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

$$\text{Orthogonal vector } q_1 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

for $\lambda = 10$,

Reduced echelon form is

$$\begin{bmatrix} 40 & 40 & 0 \\ 0 & 0 & -10 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\text{Eigen vector } v_2 = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$$

$$\text{Orthogonal vector } q_2 = 1/\sqrt{2} \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$$

for $\lambda = 90$,

Reduced echelon form is

$$\begin{bmatrix} -40 & 40 & 0 \\ 0 & 0 & -90 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\text{Eigen vector } v_3 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$$

$$\text{Orthogonal vector } q_3 = 1/\sqrt{2} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$$

$$U_{3 \times 3} = \begin{bmatrix} -1/\sqrt{2} & 1/\sqrt{2} & 0 \\ 1/\sqrt{2} & 1/\sqrt{2} & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$A^T A = \begin{bmatrix} 74 & 32 \\ 32 & 26 \end{bmatrix}$$

Eigen value for $A^T A$,

$\lambda = 10, 90$

Singular values, $\sigma_1 = \sqrt{10}$, $\sigma_2 = \sqrt{90}$

for $\lambda = 10$,

$$\text{Eigen vector } v_1 = \begin{bmatrix} -1 \\ 2 \end{bmatrix}$$

$$\text{Orthogonal vector } q_1 = 1/\sqrt{5} \begin{bmatrix} -1 \\ 2 \end{bmatrix}$$

for $\lambda = 90$,

$$\text{Eigen vector } v_2 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

$$\text{Orthogonal vector } q_2 = 1/\sqrt{5} \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

$$V_{2 \times 2} = \begin{bmatrix} -1/\sqrt{5} & 2/\sqrt{5} \\ 2/\sqrt{5} & 1/\sqrt{5} \end{bmatrix}$$

$$V_{2 \times 2}^T = \begin{bmatrix} -1/\sqrt{5} & 2/\sqrt{5} \\ 2/\sqrt{5} & 1/\sqrt{5} \end{bmatrix}$$

$$A = \begin{bmatrix} -1/\sqrt{2} & 1/\sqrt{2} & 0 \\ 1/\sqrt{2} & 1/\sqrt{2} & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \sqrt{10} & 0 \\ 0 & \sqrt{90} \\ 0 & 0 \end{bmatrix} \begin{bmatrix} -1/\sqrt{5} & 2/\sqrt{5} \\ 2/\sqrt{5} & 1/\sqrt{5} \end{bmatrix}$$

2.2

$$A = \begin{bmatrix} 3 & 2 & 2 \\ 2 & 3 & -2 \end{bmatrix}, A = U_{2 \times 2} \Sigma_{2 \times 3} V_{3 \times 3}$$

$$AA^T = \begin{bmatrix} 17 & 8 \\ 8 & 17 \end{bmatrix}$$

Eigen value for AA^T ,

$\lambda = 9, 25$

Singular values, $\sigma_1 = 3, \sigma_2 = 5$

$$\Sigma_{2 \times 3} = \begin{bmatrix} 3 & 0 & 0 \\ 0 & 5 & 0 \end{bmatrix}$$

$$(AA^T - \lambda I) = 0$$

for $\lambda = 9$,

Reduced echelon form is

$$\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$$

$$\text{Eigen vector } v_1 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

$$\text{Orthogonal vector } q_1 = 1/\sqrt{2} \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

for $\lambda = 25$,

Reduced echelon form is

$$\begin{bmatrix} -1 & 1 \\ 0 & 0 \end{bmatrix}$$

$$\text{Eigen vector } v_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$\text{Orthogonal vector } q_2 = 1/\sqrt{2} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$U_{2 \times 2} = \begin{bmatrix} -1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & 1/\sqrt{2} \end{bmatrix}$$

calculating V matrix,

$$v_i = 1/\sigma_i A^T u_i$$

$$v_1 = 1/3 \begin{bmatrix} 3 & 2 \\ 2 & 3 \\ 2 & -2 \end{bmatrix} \begin{bmatrix} -1/\sqrt{2} \\ 1/\sqrt{2} \end{bmatrix}$$

$$v_1 = \begin{bmatrix} -1/3\sqrt{2} \\ 1/3\sqrt{2} \\ -4/3\sqrt{2} \end{bmatrix}$$

$$v_2 = 1/5 \begin{bmatrix} 3 & 2 \\ 2 & 3 \\ 2 & -2 \end{bmatrix} \begin{bmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \end{bmatrix}$$

$$v_2 = \begin{bmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \\ 0 \end{bmatrix}$$

taking v_3 as perpendicular to both v_1 and v_2 ,

$$v_3 = \begin{bmatrix} -2/3 \\ 2/3 \\ 1/3 \end{bmatrix}$$

$$V_{3 \times 3} = \begin{bmatrix} -1/3\sqrt{2} & 1/\sqrt{2} & -2/3 \\ 1/3\sqrt{2} & 1/\sqrt{2} & 2/3 \\ -4/3\sqrt{2} & 0 & 1/3 \end{bmatrix}$$

$$V_{3 \times 3}^T = \begin{bmatrix} -1/3\sqrt{2} & 1/3\sqrt{2} & -4/3\sqrt{2} \\ 1/\sqrt{2} & 1/\sqrt{2} & 0 \\ -2/3 & 2/3 & 1/3 \end{bmatrix}$$

$$A = \begin{bmatrix} -1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & 1/\sqrt{2} \end{bmatrix} \begin{bmatrix} 3 & 0 & 0 \\ 0 & 5 & 0 \end{bmatrix} \begin{bmatrix} -1/3\sqrt{2} & 1/3\sqrt{2} & -4/3\sqrt{2} \\ 1/\sqrt{2} & 1/\sqrt{2} & 0 \\ -2/3 & 2/3 & 1/3 \end{bmatrix}$$

$$5.1 \quad A = \begin{bmatrix} -6 & 2 & 0 \\ 2 & -6 & 2 \\ 0 & 2 & -6 \end{bmatrix}$$

A is square matrix and transpose is:

$$A^T = \begin{bmatrix} -6 & 2 & 0 \\ 2 & -6 & 2 \\ 0 & 2 & -6 \end{bmatrix}$$

$A = A^T$, so A is symmetric matrix.

$$5.2 \quad A = \begin{bmatrix} 1 & 2 & 2 & 1 \\ 2 & 2 & 2 & 1 \\ 2 & 2 & 1 & 2 \end{bmatrix}$$

A is not a square matrix so A is not symmetric.

7.

$$B = C^T A C$$

$$X^T B X = X^T C^T A C X$$

assume, $CX = Y$ then $Y^T = X^T C^T$

$$X^T B X = Y^T A Y,$$

A is symmetric positive definite and C is nonsingular

so, $Y \neq 0$,

hence,

$$X^T B X > 0.$$

B is also symmetric positive definite