

MATHEMATICS-I (MATH F111)

Dr. Krishnendra Shekhawat

BITS PILANI
Department of Mathematics



Section 11.4

Graphing in Polar Coordinates



Symmetry Tests

- **Symmetry about x -axis:** If (r, θ) lies on the graph, then $(r, -\theta)$ or $(-r, \pi - \theta)$ also lies on the graph.



Symmetry Tests

- **Symmetry about x -axis:** If (r, θ) lies on the graph, then $(r, -\theta)$ or $(-r, \pi - \theta)$ also lies on the graph.
- **Symmetry about y -axis:** If (r, θ) lies on the graph, then $(-r, -\theta)$ or $(r, \pi - \theta)$ also lies on the graph.



Symmetry Tests

- **Symmetry about x -axis:** If (r, θ) lies on the graph, then $(r, -\theta)$ or $(-r, \pi - \theta)$ also lies on the graph.
- **Symmetry about y -axis:** If (r, θ) lies on the graph, then $(-r, -\theta)$ or $(r, \pi - \theta)$ also lies on the graph.
- **Symmetry about origin:** If (r, θ) lies on the graph, then $(-r, \theta)$ or $(r, \pi + \theta)$ also lies on the graph.



Symmetry Tests

- **Symmetry about x -axis:** If (r, θ) lies on the graph, then $(r, -\theta)$ or $(-r, \pi - \theta)$ also lies on the graph.
- **Symmetry about y -axis:** If (r, θ) lies on the graph, then $(-r, -\theta)$ or $(r, \pi - \theta)$ also lies on the graph.
- **Symmetry about origin:** If (r, θ) lies on the graph, then $(-r, \theta)$ or $(r, \pi + \theta)$ also lies on the graph.
- **Symmetry about line $y = x$:** If (r, θ) lies on the graph, then $(r, \frac{\pi}{2} - \theta)$ or $(-r, -\frac{\pi}{2} - \theta)$ also lies on the graph.



Slope of a Polar Curve



Slope of a Polar Curve

- The parametric equations of $r = f(\theta)$ are $x = r \cos \theta$, $y = r \sin \theta$.



Slope of a Polar Curve

- The parametric equations of $r = f(\theta)$ are $x = r \cos \theta$, $y = r \sin \theta$.
- The slope of the curve $r = f(\theta)$ at any point (r, θ) is given by

$$\left. \frac{dy}{dx} \right|_{(r, \theta)} = \frac{\left. \frac{dy}{d\theta} \right|_{(r, \theta)}}{\left. \frac{dx}{d\theta} \right|_{(r, \theta)}} = \frac{f'(\theta) \sin \theta + f(\theta) \cos \theta}{f'(\theta) \cos \theta - f(\theta) \sin \theta}$$

provided $\frac{dx}{d\theta} \neq 0$ at any point (r, θ) .



Tracing the Curve in Polar Coordinate

Q.: Trace the curve $r = 1 + \cos \theta$.



Tracing the Curve in Polar Coordinate

Q:. Trace the curve $r = 1 + \cos \theta$.

Step 1. Check for symmetries (it will reduce the work for tracing).



Tracing the Curve in Polar Coordinate

Q:. Trace the curve $r = 1 + \cos \theta$.

Step 1. Check for symmetries (it will reduce the work for tracing).

- Since $(r, -\theta)$ lies on the curve, it is symmetric about x -axis.



Tracing the Curve in Polar Coordinate

Q:. Trace the curve $r = 1 + \cos \theta$.

Step 1. Check for symmetries (it will reduce the work for tracing).

- Since $(r, -\theta)$ lies on the curve, it is symmetric about x -axis. Hence, it is enough to consider the steps for $0 \leq \theta \leq \pi$.



Step 2. Solve the equation $r = 0$ for θ .



Step 2. Solve the equation $r = 0$ for θ . If $\theta = \theta_0$ satisfies $r = 0$, then the line $\theta = \theta_0$ will be a tangent to the curve at pole.



Step 2. Solve the equation $r = 0$ for θ . If $\theta = \theta_0$ satisfies $r = 0$, then the line $\theta = \theta_0$ will be a tangent to the curve at pole.

□ $r = 0$ gives $\cos \theta = -1 \implies \theta = \pi$. Thus $\theta = \pi$ is a tangent to the curve at pole.



Step 3 Find $\frac{dr}{d\theta}$.



Step 3 Find $\frac{dr}{d\theta}$.

$$\square \frac{dr}{d\theta} = -\sin \theta.$$



Step 3 Find $\frac{dr}{d\theta}$.

$$\square \frac{dr}{d\theta} = -\sin \theta.$$

Step 3.1 Find the values of θ for which $\frac{dr}{d\theta} > 0$.



Step 3 Find $\frac{dr}{d\theta}$.

$$\square \frac{dr}{d\theta} = -\sin \theta.$$

Step 3.1 Find the values of θ for which $\frac{dr}{d\theta} > 0$. This would be an interval (or no value of θ). In this interval r is an increasing function.



Step 3 Find $\frac{dr}{d\theta}$.

$$\square \frac{dr}{d\theta} = -\sin \theta.$$

Step 3.1 Find the values of θ for which $\frac{dr}{d\theta} > 0$. This would be an interval (or no value of θ). In this interval r is an increasing function. Find max/min values of r and the associated θ values in this interval.



Step 3 Find $\frac{dr}{d\theta}$.

$$\square \frac{dr}{d\theta} = -\sin \theta.$$

Step 3.1 Find the values of θ for which $\frac{dr}{d\theta} > 0$. This would be an interval (or no value of θ). In this interval r is an increasing function. Find max/min values of r and the associated θ values in this interval.

$$\square \frac{dr}{d\theta} > 0 \Rightarrow \sin \theta < 0,$$



Step 3 Find $\frac{dr}{d\theta}$.

$$\square \frac{dr}{d\theta} = -\sin \theta.$$

Step 3.1 Find the values of θ for which $\frac{dr}{d\theta} > 0$. This would be an interval (or no value of θ). In this interval r is an increasing function. Find max/min values of r and the associated θ values in this interval.

$$\square \frac{dr}{d\theta} > 0 \Rightarrow \sin \theta < 0, \text{ thus no value of } \theta \text{ in between } 0 \text{ and } \pi.$$



Step 3.2 Similarly find the values of θ for which $\frac{dr}{d\theta} < 0$. This would be an interval (or no value of θ). In this interval r is a decreasing function. Find max/min values of r and the associated θ values in this interval.



Step 3.2 Similarly find the values of θ for which $\frac{dr}{d\theta} < 0$. This would be an interval (or no value of θ). In this interval r is a decreasing function. Find max/min values of r and the associated θ values in this interval.

- $\frac{dr}{d\theta} < 0 \Rightarrow \sin \theta > 0 \Rightarrow 0 < \theta < \pi$, thus r decreases in the interval $[0, \pi]$.



Step 3.2 Similarly find the values of θ for which $\frac{dr}{d\theta} < 0$. This would be an interval (or no value of θ). In this interval r is a decreasing function. Find max/min values of r and the associated θ values in this interval.

□ $\frac{dr}{d\theta} < 0 \Rightarrow \sin \theta > 0 \Rightarrow 0 < \theta < \pi$, thus r decreases in the interval $[0, \pi]$.

Clearly $\max r = 2$ at $\theta = 0$ and $\min r = 0$ at $\theta = \pi$.



Step 4. Find slopes at some special points (like at end points of the intervals obtained in step 3).



Step 4. Find slopes at some special points (like at end points of the intervals obtained in step 3).

$$\left. \frac{dy}{dx} \right|_{\theta=0} = \infty, \quad \left. \frac{dy}{dx} \right|_{\theta=\frac{\pi}{2}} = 1.$$



Step 4. Find slopes at some special points (like at end points of the intervals obtained in step 3).

$$\left. \frac{dy}{dx} \right|_{\theta=0} = \infty, \quad \left. \frac{dy}{dx} \right|_{\theta=\frac{\pi}{2}} = 1.$$

At $\theta = 0$, $r = 2$ and $\tan \theta_1 = \infty \implies \theta_1 = \frac{\pi}{2}$.



Step 4. Find slopes at some special points (like at end points of the intervals obtained in step 3).

$$\left. \frac{dy}{dx} \right|_{\theta=0} = \infty, \quad \left. \frac{dy}{dx} \right|_{\theta=\frac{\pi}{2}} = 1.$$

At $\theta = 0$, $r = 2$ and $\tan \theta_1 = \infty \implies \theta_1 = \frac{\pi}{2}$.

At $\theta = \frac{\pi}{2}$, $r = 1$ and $\tan \theta_1 = 1 \implies \theta_1 = \frac{\pi}{4}$.



Step 5. Make a table θ vs r for different values of θ .



Step 5. Make a table θ vs r for different values of θ .

θ	0	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	π
r	2	$1 + \frac{1}{\sqrt{2}}$	$\frac{3}{2}$	1	$\frac{1}{2}$	$1 - \frac{1}{\sqrt{2}}$	0



Step 6. Plot the curve while considering the above steps.






Figure: $r = 1 + \cos \theta$: The cardioid



Q: Trace the curve $r = 1 - \sin \theta$.



Q:. Trace the curve $r = 1 - \sin \theta$.

Step 1. Symmetry



Q:. Trace the curve $r = 1 - \sin \theta$.

Step 1. Symmetry

□ It is symmetric about y -axis.



Q:. Trace the curve $r = 1 - \sin \theta$.

Step 1. Symmetry

- It is symmetric about y-axis. Hence, it is enough to consider the steps for $\pi/2 \leq \theta \leq 3\pi/2$.



Q:. Trace the curve $r = 1 - \sin \theta$.

Step 1. Symmetry

- It is symmetric about y-axis. Hence, it is enough to consider the steps for $\pi/2 \leq \theta \leq 3\pi/2$.

Step 2. Solve the equation $r = 0$ for θ .



Q.: Trace the curve $r = 1 - \sin \theta$.

Step 1. Symmetry

- It is symmetric about y-axis. Hence, it is enough to consider the steps for $\pi/2 \leq \theta \leq 3\pi/2$.

Step 2. Solve the equation $r = 0$ for θ .

- $r = 0$ gives $\sin \theta = 1 \implies \theta = \pi/2$. Thus $\theta = \pi/2$ is a tangent to the curve at pole.



Step 3 Find $\frac{dr}{d\theta}$.



Step 3 Find $\frac{dr}{d\theta}$.

$$\square \frac{dr}{d\theta} = -\cos \theta.$$



Step 3 Find $\frac{dr}{d\theta}$.

$$\square \frac{dr}{d\theta} = -\cos \theta.$$

Step 3.1 and 3.2 Find the values of θ for which $\frac{dr}{d\theta} > 0$ and $\frac{dr}{d\theta} < 0$.



Step 3 Find $\frac{dr}{d\theta}$.

$$\square \frac{dr}{d\theta} = -\cos \theta.$$

Step 3.1 and 3.2 Find the values of θ for which $\frac{dr}{d\theta} > 0$ and $\frac{dr}{d\theta} < 0$.



Step 3 Find $\frac{dr}{d\theta}$.

$$\square \frac{dr}{d\theta} = -\cos \theta.$$

Step 3.1 and 3.2 Find the values of θ for which $\frac{dr}{d\theta} > 0$ and $\frac{dr}{d\theta} < 0$.

$$\square \frac{dr}{d\theta} > 0 \Rightarrow \cos \theta < 0 \Rightarrow \pi/2 < \theta < 3\pi/2,$$



Step 3 Find $\frac{dr}{d\theta}$.

$$\square \frac{dr}{d\theta} = -\cos \theta.$$

Step 3.1 and 3.2 Find the values of θ for which $\frac{dr}{d\theta} > 0$ and $\frac{dr}{d\theta} < 0$.

$$\square \frac{dr}{d\theta} > 0 \Rightarrow \cos \theta < 0 \Rightarrow \pi/2 < \theta < 3\pi/2, \text{ thus } r \text{ increases in the interval } [\pi/2, 3\pi/2].$$



Step 3 Find $\frac{dr}{d\theta}$.

$$\square \frac{dr}{d\theta} = -\cos \theta.$$

Step 3.1 and 3.2 Find the values of θ for which $\frac{dr}{d\theta} > 0$ and $\frac{dr}{d\theta} < 0$.

$$\square \frac{dr}{d\theta} > 0 \Rightarrow \cos \theta < 0 \Rightarrow \pi/2 < \theta < 3\pi/2, \text{ thus } r \text{ increases in the interval } [\pi/2, 3\pi/2].$$

Clearly $\max r = 2$ at $\theta = 3\pi/2$ and $\min r = 0$ at $\theta = \pi/2$.



Step 3 Find $\frac{dr}{d\theta}$.

$$\square \frac{dr}{d\theta} = -\cos \theta.$$

Step 3.1 and 3.2 Find the values of θ for which $\frac{dr}{d\theta} > 0$ and $\frac{dr}{d\theta} < 0$.

$$\square \frac{dr}{d\theta} > 0 \Rightarrow \cos \theta < 0 \Rightarrow \pi/2 < \theta < 3\pi/2, \text{ thus } r \text{ increases in the interval } [\pi/2, 3\pi/2].$$

Clearly $\max r = 2$ at $\theta = 3\pi/2$ and $\min r = 0$ at $\theta = \pi/2$.

$$\square \frac{dr}{d\theta} < 0 \Rightarrow \cos \theta > 0,$$



Step 3 Find $\frac{dr}{d\theta}$.

$$\square \frac{dr}{d\theta} = -\cos \theta.$$

Step 3.1 and 3.2 Find the values of θ for which $\frac{dr}{d\theta} > 0$ and $\frac{dr}{d\theta} < 0$.

$$\square \frac{dr}{d\theta} > 0 \Rightarrow \cos \theta < 0 \Rightarrow \pi/2 < \theta < 3\pi/2, \text{ thus } r \text{ increases in the interval } [\pi/2, 3\pi/2].$$

Clearly $\max r = 2$ at $\theta = 3\pi/2$ and $\min r = 0$ at $\theta = \pi/2$.

$$\square \frac{dr}{d\theta} < 0 \Rightarrow \cos \theta > 0, \text{ thus no value of } \theta.$$



Step 4. Find slopes at some special points.



Step 4. Find slopes at some special points.

$$\left. \frac{dy}{dx} \right|_{\theta=\pi} = 1, \quad \left. \frac{dy}{dx} \right|_{\theta=\frac{3\pi}{2}} = 0.$$



Step 4. Find slopes at some special points.

$$\left. \frac{dy}{dx} \right|_{\theta=\pi} = 1, \quad \left. \frac{dy}{dx} \right|_{\theta=\frac{3\pi}{2}} = 0.$$

At $\theta = \pi$, $r = 1$ and $\tan \theta_1 = 1 \implies \theta_1 = \frac{\pi}{4}$.



Step 4. Find slopes at some special points.

$$\left. \frac{dy}{dx} \right|_{\theta=\pi} = 1, \quad \left. \frac{dy}{dx} \right|_{\theta=\frac{3\pi}{2}} = 0.$$

At $\theta = \pi$, $r = 1$ and $\tan \theta_1 = 1 \implies \theta_1 = \frac{\pi}{4}$.

At $\theta = \frac{3\pi}{2}$, $r = 2$ and $\tan \theta_1 = 0 \implies \theta_1 = 0$.



Step 4. Find slopes at some special points.

$$\left. \frac{dy}{dx} \right|_{\theta=\pi} = 1, \quad \left. \frac{dy}{dx} \right|_{\theta=\frac{3\pi}{2}} = 0.$$

At $\theta = \pi$, $r = 1$ and $\tan \theta_1 = 1 \implies \theta_1 = \frac{\pi}{4}$.

At $\theta = \frac{3\pi}{2}$, $r = 2$ and $\tan \theta_1 = 0 \implies \theta_1 = 0$.

Step 5. Make a table θ vs r for different values of θ .






Figure: $r = 1 - \sin \theta$: The cardioid



Q:.. Trace the curve $r^2 = \sin 2\theta$.



Q:. Trace the curve $r^2 = \sin 2\theta$.

Sol. Since $r^2 = \sin 2\theta \geq 0 \implies \theta \in [0, \pi/2] \cup [\pi, 3\pi/2]$.
Hence, the graph would be in first and third quadrant only.



Q:. Trace the curve $r^2 = \sin 2\theta$.

Sol. Since $r^2 = \sin 2\theta \geq 0 \implies \theta \in [0, \pi/2] \cup [\pi, 3\pi/2]$.
Hence, the graph would be in first and third quadrant only.

Remark

Note that it is enough to consider either $r = \sqrt{\sin 2\theta}$ or $r = -\sqrt{\sin 2\theta}$. Symmetry will give the graph corresponding to other.



Step 1. $(-r, \theta)$ and $(r, \frac{\pi}{2} - \theta)$ lies on the curve, therefore it is symmetric about the pole and about the line $y = x$ respectively.



Step 1. $(-r, \theta)$ and $(r, \frac{\pi}{2} - \theta)$ lies on the curve, therefore it is symmetric about the pole and about the line $y = x$ respectively. Hence, it is enough to consider the steps for $0 \leq \theta \leq \frac{\pi}{4}$.



Step 1. $(-r, \theta)$ and $(r, \frac{\pi}{2} - \theta)$ lies on the curve, therefore it is symmetric about the pole and about the line $y = x$ respectively. Hence, it is enough to consider the steps for $0 \leq \theta \leq \frac{\pi}{4}$.

Step 2. $r = 0$ gives $\sin 2\theta = 0 \implies \theta = 0$ (only $\theta = 0$ lies in $[0, \frac{\pi}{4}]$).



Step 1. $(-r, \theta)$ and $(r, \frac{\pi}{2} - \theta)$ lies on the curve, therefore it is symmetric about the pole and about the line $y = x$ respectively. Hence, it is enough to consider the steps for $0 \leq \theta \leq \frac{\pi}{4}$.

Step 2. $r = 0$ gives $\sin 2\theta = 0 \implies \theta = 0$ (only $\theta = 0$ lies in $[0, \frac{\pi}{4}]$). Thus $\theta = 0$ is tangent to the curve at pole.



Step 3. $\frac{dr}{d\theta} = \frac{\cos 2\theta}{\sqrt{\sin 2\theta}}.$



Step 3. $\frac{dr}{d\theta} = \frac{\cos 2\theta}{\sqrt{\sin 2\theta}}.$

□ $\frac{dr}{d\theta} > 0$ only when $\cos 2\theta$ and $\sin 2\theta$ both are positive or negative,



Step 3. $\frac{dr}{d\theta} = \frac{\cos 2\theta}{\sqrt{\sin 2\theta}}.$

□ $\frac{dr}{d\theta} > 0$ only when $\cos 2\theta$ and $\sin 2\theta$ both are positive or negative, i.e., $2\theta \in [0, \pi/2]$ or $\theta \in [0, \pi/4]$.



Step 3. $\frac{dr}{d\theta} = \frac{\cos 2\theta}{\sqrt{\sin 2\theta}}.$

□ $\frac{dr}{d\theta} > 0$ only when $\cos 2\theta$ and $\sin 2\theta$ both are positive or negative, i.e., $2\theta \in [0, \pi/2]$ or $\theta \in [0, \pi/4]$. Thus r increases in the interval $[0, \pi/4]$.



Step 3. $\frac{dr}{d\theta} = \frac{\cos 2\theta}{\sqrt{\sin 2\theta}}.$

- $\frac{dr}{d\theta} > 0$ only when $\cos 2\theta$ and $\sin 2\theta$ both are positive or negative, i.e., $2\theta \in [0, \pi/2]$ or $\theta \in [0, \pi/4]$. Thus r increases in the interval $[0, \pi/4]$.
- $\frac{dr}{d\theta} < 0$ when $\cos 2\theta$ and $\sin 2\theta$ are of opposite sign, which is not possible for any value of θ .



Step 3. $\frac{dr}{d\theta} = \frac{\cos 2\theta}{\sqrt{\sin 2\theta}}.$

- $\frac{dr}{d\theta} > 0$ only when $\cos 2\theta$ and $\sin 2\theta$ both are positive or negative, i.e., $2\theta \in [0, \pi/2]$ or $\theta \in [0, \pi/4]$. Thus r increases in the interval $[0, \pi/4]$.
- $\frac{dr}{d\theta} < 0$ when $\cos 2\theta$ and $\sin 2\theta$ are of opposite sign, which is not possible for any value of θ .

Clearly max value of r is 1 at $\theta = \pi/4$ and min value of r is 0 at $\theta = 0$.



Step 4.

$$\begin{aligned}\frac{dy}{dx} &= \frac{f'(\theta) \sin \theta + f(\theta) \cos \theta}{f'(\theta) \cos \theta - f(\theta) \sin \theta} \\&= \frac{\frac{\cos 2\theta}{\sqrt{\sin 2\theta}} \sin \theta + \sqrt{\sin 2\theta} \cos \theta}{\frac{\cos 2\theta}{\sqrt{\sin 2\theta}} \cos \theta - \sqrt{\sin 2\theta} \sin \theta} \\&= \frac{\cos 2\theta \sin \theta + \sin 2\theta \cos \theta}{\cos 2\theta \cos \theta - \sin 2\theta \sin \theta} \\&= \frac{\sin 3\theta}{\cos 3\theta} = \tan 3\theta\end{aligned}$$



Thus $\left. \frac{dy}{dx} \right|_{\theta=\pi/4} = \tan(3\pi/4)$. Thus slope of the tangent at $\theta = \pi/4$ is $3\pi/4$.



Thus $\left. \frac{dy}{dx} \right|_{\theta=\pi/4} = \tan(3\pi/4)$. Thus slope of the tangent at $\theta = \pi/4$ is $3\pi/4$.

Step 5. Table θ vs r : Not required.






Figure: Lemniscate: Two leaved rose



Q: Trace the curve $r = \cos 2\theta$.



Q:. Trace the curve $r = \cos 2\theta$.

Sol.

Step 1. ① Since $(r, -\theta)$ lies on the curve, so symmetric about x -axis.



Q:. Trace the curve $r = \cos 2\theta$.

Sol.

Step 1. ① Since $(r, -\theta)$ lies on the curve, so symmetric about x -axis.

② Since $(r, \pi - \theta)$ lies on the curve, so symmetric about y -axis.



Q:. Trace the curve $r = \cos 2\theta$.

Sol.

- Step 1.** ① Since $(r, -\theta)$ lies on the curve, so symmetric about x -axis.
- ② Since $(r, \pi - \theta)$ lies on the curve, so symmetric about y -axis.
- ③ Since $(-r, -\frac{\pi}{2} - \theta)$ lies on the curve, so symmetric about the line $y = x$.



Q:. Trace the curve $r = \cos 2\theta$.

Sol.

- Step 1.** ① Since $(r, -\theta)$ lies on the curve, so symmetric about x -axis.
- ② Since $(r, \pi - \theta)$ lies on the curve, so symmetric about y -axis.
- ③ Since $(-r, -\frac{\pi}{2} - \theta)$ lies on the curve, so symmetric about the line $y = x$.

Hence, it is enough to consider the region $0 \leq \theta \leq \frac{\pi}{4}$.



Step 2. $r = 0$ gives $\cos 2\theta = 0 \implies \theta = \frac{\pi}{4}$. Thus, $\theta = \frac{\pi}{4}$ is a tangent to the curve at pole.



Step 2. $r = 0$ gives $\cos 2\theta = 0 \implies \theta = \frac{\pi}{4}$. Thus, $\theta = \frac{\pi}{4}$ is a tangent to the curve at pole.

Step 3. $\frac{dr}{d\theta} = -2 \sin 2\theta$.



Step 2. $r = 0$ gives $\cos 2\theta = 0 \implies \theta = \frac{\pi}{4}$. Thus, $\theta = \frac{\pi}{4}$ is a tangent to the curve at pole.

Step 3. $\frac{dr}{d\theta} = -2 \sin 2\theta$.

□ $\frac{dr}{d\theta} > 0 \implies \sin 2\theta < 0$, thus no value of θ .



Step 2. $r = 0$ gives $\cos 2\theta = 0 \implies \theta = \frac{\pi}{4}$. Thus, $\theta = \frac{\pi}{4}$ is a tangent to the curve at pole.

Step 3. $\frac{dr}{d\theta} = -2 \sin 2\theta$.

- $\frac{dr}{d\theta} > 0 \implies \sin 2\theta < 0$, thus no value of θ .
- $\frac{dr}{d\theta} < 0 \implies \sin 2\theta > 0 \implies 0 < 2\theta < \frac{\pi}{2}$, thus r decreases in the interval $\left[0, \frac{\pi}{4}\right]$.



Step 2. $r = 0$ gives $\cos 2\theta = 0 \implies \theta = \frac{\pi}{4}$. Thus, $\theta = \frac{\pi}{4}$ is a tangent to the curve at pole.

Step 3. $\frac{dr}{d\theta} = -2 \sin 2\theta$.

□ $\frac{dr}{d\theta} > 0 \implies \sin 2\theta < 0$, thus no value of θ .

□ $\frac{dr}{d\theta} < 0 \implies \sin 2\theta > 0 \implies 0 < 2\theta < \frac{\pi}{2}$, thus r decreases in the interval $\left[0, \frac{\pi}{4}\right]$.

Clearly Max $r = 1$ at $\theta = 0$ and min $r = 0$ at $\theta = \frac{\pi}{4}$.



Step 4.

$$\left. \frac{dy}{dx} \right|_{\theta=0} = \infty.$$

Thus the tangent to the curve at $\theta = 0$ is perpendicular to x -axis.



Step 4.

$$\left. \frac{dy}{dx} \right|_{\theta=0} = \infty.$$

Thus the tangent to the curve at $\theta = 0$ is perpendicular to x -axis.

Step 5. Table

θ	0	$\frac{\pi}{12}$	$\frac{\pi}{6}$	$\frac{\pi}{4}$
r	1	0.86	0.5	0






Figure: $r = \cos 2\theta$: Four leaved rose



Q: Trace the curve $r = 1 + 2 \sin \theta$.



Q:. Trace the curve $r = 1 + 2 \sin \theta$.

Sol.

Step 1. Since $(r, \pi - \theta)$ lies on the curve, so symmetric about y-axis.



Q:. Trace the curve $r = 1 + 2 \sin \theta$.

Sol.

Step 1. Since $(r, \pi - \theta)$ lies on the curve, so symmetric about y -axis. Hence, it is enough to consider the region $\frac{\pi}{2} \leq \theta \leq \frac{3\pi}{2}$.



Q.: Trace the curve $r = 1 + 2 \sin \theta$.

Sol.

Step 1. Since $(r, \pi - \theta)$ lies on the curve, so symmetric about y -axis. Hence, it is enough to consider the region $\frac{\pi}{2} \leq \theta \leq \frac{3\pi}{2}$.

Step 2. $r = 0$ gives $\sin \theta = -1/2 \implies \theta = \frac{7\pi}{6}, \frac{11\pi}{6}$. Thus, $\theta = \frac{7\pi}{6}$ and $\theta = \frac{11\pi}{6}$ are tangents to the curve at pole.



Step 3. $\frac{dr}{d\theta} = 2 \cos \theta.$



Step 3. $\frac{dr}{d\theta} = 2\cos\theta$.

□ $\frac{dr}{d\theta} > 0 \Rightarrow \cos\theta > 0$, thus no value of θ .



Step 3. $\frac{dr}{d\theta} = 2\cos\theta$.

□ $\frac{dr}{d\theta} > 0 \Rightarrow \cos\theta > 0$, thus no value of θ .

□ $\frac{dr}{d\theta} < 0 \Rightarrow \cos\theta < 0 \Rightarrow \frac{\pi}{2} < \theta < \frac{3\pi}{2}$, thus r decreases in the interval $\left[\frac{\pi}{2}, \frac{3\pi}{2}\right]$.



Step 3. $\frac{dr}{d\theta} = 2\cos\theta$.

□ $\frac{dr}{d\theta} > 0 \Rightarrow \cos\theta > 0$, thus no value of θ .

□ $\frac{dr}{d\theta} < 0 \Rightarrow \cos\theta < 0 \Rightarrow \frac{\pi}{2} < \theta < \frac{3\pi}{2}$, thus r decreases in the interval $\left[\frac{\pi}{2}, \frac{3\pi}{2}\right]$.

Clearly Max $r = 3$ at $\theta = \frac{\pi}{2}$ and min $r = -1$ at $\theta = \frac{3\pi}{2}$.



Step 4.

$$\left. \frac{dy}{dx} \right|_{\theta=\frac{\pi}{2}} = 0, \quad \left. \frac{dy}{dx} \right|_{\theta=\pi} = -1/2, \quad \left. \frac{dy}{dx} \right|_{\theta=\frac{3\pi}{2}} = 0.$$



Step 4.

$$\left. \frac{dy}{dx} \right|_{\theta=\frac{\pi}{2}} = 0, \quad \left. \frac{dy}{dx} \right|_{\theta=\pi} = -1/2, \quad \left. \frac{dy}{dx} \right|_{\theta=\frac{3\pi}{2}} = 0.$$

At $\theta = \frac{\pi}{2}$, $r = 3$ and $\tan \theta_1 = 0 \implies \theta_1 = 0$.



Step 4.

$$\left. \frac{dy}{dx} \right|_{\theta=\frac{\pi}{2}} = 0, \quad \left. \frac{dy}{dx} \right|_{\theta=\pi} = -1/2, \quad \left. \frac{dy}{dx} \right|_{\theta=\frac{3\pi}{2}} = 0.$$

At $\theta = \frac{\pi}{2}$, $r = 3$ and $\tan \theta_1 = 0 \implies \theta_1 = 0$.

At $\theta = \pi$, $r = 1$ and

$\tan \theta_1 = -1/2 \implies \theta_1 = -26.57$.



Step 4.

$$\left. \frac{dy}{dx} \right|_{\theta=\frac{\pi}{2}} = 0, \quad \left. \frac{dy}{dx} \right|_{\theta=\pi} = -1/2, \quad \left. \frac{dy}{dx} \right|_{\theta=\frac{3\pi}{2}} = 0.$$

At $\theta = \frac{\pi}{2}$, $r = 3$ and $\tan \theta_1 = 0 \implies \theta_1 = 0$.

At $\theta = \pi$, $r = 1$ and
 $\tan \theta_1 = -1/2 \implies \theta_1 = -26.57$.

At $\theta = \frac{3\pi}{2}$, $r = -1$ and $\tan \theta_1 = 0 \implies \theta_1 = 0$.



Step 4.

$$\left. \frac{dy}{dx} \right|_{\theta=\frac{\pi}{2}} = 0, \quad \left. \frac{dy}{dx} \right|_{\theta=\pi} = -1/2, \quad \left. \frac{dy}{dx} \right|_{\theta=\frac{3\pi}{2}} = 0.$$


At $\theta = \frac{\pi}{2}$, $r = 3$ and $\tan \theta_1 = 0 \implies \theta_1 = 0$.

At $\theta = \pi$, $r = 1$ and
 $\tan \theta_1 = -1/2 \implies \theta_1 = -26.57$.


At $\theta = \frac{3\pi}{2}$, $r = -1$ and $\tan \theta_1 = 0 \implies \theta_1 = 0$.

Step 5. Table





Trace the curve $r^2 = \cos 2\theta$



Q:.. Trace the curve $r = 1 - \cos \theta$.



Q:. Trace the curve $r = 1 - \cos \theta$.




Figure: $r = 1 - \cos \theta$: The cardioid



Sol. We can write $r = 1 - \cos \theta = 1 + \cos(\theta - \pi)$.



Sol. We can write $r = 1 - \cos \theta = 1 + \cos(\theta - \pi)$.

Thus the curve $r = 1 - \cos \theta$ is obtained from $r = 1 + \cos \theta$ by replacing θ by $\theta + \pi$. Therefore, to obtain the curve of $r = 1 - \cos \theta$, we just need to rotate the curve of $r = 1 + \cos \theta$ by an angle π .



Faster Graphing



Steps for Faster Graphing

The polar equation can quickly be captured by plotting $r = f(\theta)$ in cartesian θr -plane.



Steps for Faster Graphing

The polar equation can quickly be captured by plotting $r = f(\theta)$ in cartesian θr -plane. How?

Step 1: Compute $\theta - r$ table for θ in the interval $[0, 2\pi]$.



Steps for Faster Graphing

The polar equation can quickly be captured by plotting $r = f(\theta)$ in cartesian θr -plane. **How?**

Step 1: Compute $\theta - r$ table for θ in the interval $[0, 2\pi]$.

□ Consider the curve $r = \cos 2\theta$




θ	0	$\frac{\pi}{4}$	$\frac{\pi}{2}$	$\frac{3\pi}{4}$	π	$\frac{5\pi}{4}$	$\frac{3\pi}{2}$	$\frac{7\pi}{4}$	2π
r	1	0	-1	0	1	0	-1	0	1



Step 2: Graph $r = f(\theta)$ in the cartesian θr -plane (that is, x -axis for θ and y -axis for r) in the interval $[0, 2\pi]$.



Step 2: Graph $r = f(\theta)$ in the cartesian θr -plane (that is, x -axis for θ and y -axis for r) in the interval $[0, 2\pi]$.



Remark

The value of θ where the curve crosses (or touches) θ -axis is a tangent to the curve at pole.



Remark

The value of θ where the curve crosses (or touches) θ -axis is a tangent to the curve at pole. For example, $\theta = \frac{\pi}{4}$ is a tangent to the curve $r = \cos 2\theta$.







Figure: $r = \cos 2\theta$: Four leaved rose



Q: Trace the curve $r = \sin 3\theta$ using faster graphing.



Q:. Trace the curve $r = \sin 3\theta$ using faster graphing.






Figure: $r = \sin 3\theta$



$$r^2 = f(\theta)$$


For the equation of the form $r^2 = f(\theta)$ first trace $r^2 = f(\theta)$ in the cartesian θr^2 -plane (that is, x -axis for θ and y -axis for r^2) in the interval $[0, 2\pi]$ and then graph $r = f(\theta)$ in the cartesian θr -plane (that is, x -axis for θ and y -axis for r) in the interval $[0, 2\pi]$.




Q: Trace the curve $r^2 = 4 \cos 2\theta$ using faster graphing.



Q:. Trace the curve $r^2 = 4 \cos 2\theta$ using faster graphing.





Remarks

- The only drawback of faster graphing is that we can not find the slopes at particular points. But the slope is not important in the case when we evaluate the areas or lengths of curves.



Remarks

- The only drawback of faster graphing is that we can not find the slopes at particular points. But the slope is not important in the case when we evaluate the areas or lengths of curves.
- More examples over Fasting Graphing will be covered in Next Section.

