

MATHEMATICS-II (MATH F112)

Dr. Krishnendra Shekhawat

BITS PILANI
Department of Mathematics



Section 3.4

Eigenvalues and Eigenvectors



Eigenvalue and Eigenvector



Eigenvalue and Eigenvector

Let A be $n \times n$ matrix.



Eigenvalue and Eigenvector

Let A be $n \times n$ matrix. A real number λ is an **eigenvalue** of A if and only if



Eigenvalue and Eigenvector

Let A be $n \times n$ matrix. A real number λ is an **eigenvalue** of A if and only if there is a nonzero n -vector X such that $AX = \lambda X$.



Eigenvalue and Eigenvector

Let A be $n \times n$ matrix. A real number λ is an **eigenvalue** of A if and only if there is a nonzero n -vector X such that $AX = \lambda X$.

Also, any nonzero vector X for which $AX = \lambda X$,



Eigenvalue and Eigenvector

Let A be $n \times n$ matrix. A real number λ is an **eigenvalue** of A if and only if there is a nonzero n -vector X such that $AX = \lambda X$.

Also, any nonzero vector X for which $AX = \lambda X$, is an **eigenvector** corresponding to the eigenvalue λ .



Example 1



Example 1

Consider the matrix $A = \begin{bmatrix} -4 & 8 & -12 \\ 6 & -6 & 12 \\ 6 & -8 & 14 \end{bmatrix}$.



Example 1

Consider the matrix $A = \begin{bmatrix} -4 & 8 & -12 \\ 6 & -6 & 12 \\ 6 & -8 & 14 \end{bmatrix}$.

We can see that $A \begin{bmatrix} 4 \\ 3 \\ 0 \end{bmatrix} = 2 \begin{bmatrix} 4 \\ 3 \\ 0 \end{bmatrix} \Rightarrow$



Example 1

Consider the matrix $A = \begin{bmatrix} -4 & 8 & -12 \\ 6 & -6 & 12 \\ 6 & -8 & 14 \end{bmatrix}$.

We can see that $A \begin{bmatrix} 4 \\ 3 \\ 0 \end{bmatrix} = 2 \begin{bmatrix} 4 \\ 3 \\ 0 \end{bmatrix} \implies \lambda = 2$ is an eigenvalue for A and $X = [4, 3, 0]$ is the corresponding eigenvector.



If X is an eigenvector of A corresponding to an eigenvalue λ i.e.

$$AX = \lambda X,$$



If X is an eigenvector of A corresponding to an eigenvalue λ i.e.

$$AX = \lambda X,$$

then for $c \in \mathbb{R}$, we have

$$A(cX) = c(AX) = c(\lambda X) = \lambda(cX).$$



If X is an eigenvector of A corresponding to an eigenvalue λ i.e.

$$AX = \lambda X,$$

then for $c \in \mathbb{R}$, we have

$$A(cX) = c(AX) = c(\lambda X) = \lambda(cX).$$

Thus, if X is an eigenvector of A corresponding to an eigenvalue λ then, for $c \in \mathbb{R}, c \neq 0$, cX is also an eigenvector corresponding to λ .



If X is an eigenvector of A corresponding to an eigenvalue λ i.e.

$$AX = \lambda X,$$

then for $c \in \mathbb{R}$, we have

$$A(cX) = c(AX) = c(\lambda X) = \lambda(cX).$$

Thus, if X is an eigenvector of A corresponding to an eigenvalue λ then, for $c \in \mathbb{R}, c \neq 0$, cX is also an eigenvector corresponding to λ . Hence, there are infinitely many eigenvectors corresponding to an eigenvalue.



Q:. How to compute all the eigenvalues and eigenvectors of an $n \times n$ matrix.



Q:. How to compute all the eigenvalues and eigenvectors of an $n \times n$ matrix.

Sol. $AX = \lambda X = \lambda I_n X \implies$



Q:. How to compute all the eigenvalues and eigenvectors of an $n \times n$ matrix.

Sol. $AX = \lambda X = \lambda I_n X \implies (\lambda I_n - A)X = \mathbf{0}.$



Q:. How to compute all the eigenvalues and eigenvectors of an $n \times n$ matrix.

Sol. $AX = \lambda X = \lambda I_n X \implies (\lambda I_n - A)X = \mathbf{0}$. Now X is a nontrivial solution to the homogeneous system whose coefficient matrix is $\lambda I_n - A$.



Q:. How to compute all the eigenvalues and eigenvectors of an $n \times n$ matrix.

Sol. $AX = \lambda X = \lambda I_n X \implies (\lambda I_n - A)X = \mathbf{0}$. Now X is a nontrivial solution to the homogeneous system whose coefficient matrix is $\lambda I_n - A$. Therefore, $|\lambda I_n - A| = 0$.



Q:. How to compute all the eigenvalues and eigenvectors of an $n \times n$ matrix.

Sol. $AX = \lambda X = \lambda I_n X \implies (\lambda I_n - A)X = \mathbf{0}$. Now X is a nontrivial solution to the homogeneous system whose coefficient matrix is $\lambda I_n - A$. Therefore, $|\lambda I_n - A| = 0$.

Theorem: Let A be $n \times n$ matrix and λ be a real number. Then λ is an eigenvalue of A if and only if $|\lambda I_n - A| = 0$.



Q:. How to compute all the eigenvalues and eigenvectors of an $n \times n$ matrix.

Sol. $AX = \lambda X = \lambda I_n X \implies (\lambda I_n - A)X = \mathbf{0}$. Now X is a nontrivial solution to the homogeneous system whose coefficient matrix is $\lambda I_n - A$. Therefore, $|\lambda I_n - A| = 0$.

Theorem: Let A be $n \times n$ matrix and λ be a real number. Then λ is an eigenvalue of A if and only if $|\lambda I_n - A| = 0$. The eigenvectors are the nontrivial solutions of the homogeneous system $(\lambda I_n - A)X = \mathbf{0}$.



The Characteristic Polynomial of a Matrix



The Characteristic Polynomial of a Matrix

Let A is a $n \times n$ matrix, then the characteristic polynomial of A



The Characteristic Polynomial of a Matrix

Let A is a $n \times n$ matrix, then the characteristic polynomial of A is the polynomial $p_A(x) = |xI_n - A|$.



The Characteristic Polynomial of a Matrix

Let A is a $n \times n$ matrix, then the **characteristic polynomial of A** is the polynomial $p_A(x) = |xI_n - A|$.

Clearly, $p_A(x)$ is a polynomial of degree $n \implies$



The Characteristic Polynomial of a Matrix

Let A is a $n \times n$ matrix, then the **characteristic polynomial of A** is the polynomial $p_A(x) = |xI_n - A|$.

Clearly, $p_A(x)$ is a polynomial of degree $n \implies$ it has at most n real roots. Hence, from above Theorem we can say that



The Characteristic Polynomial of a Matrix

Let A is a $n \times n$ matrix, then the **characteristic polynomial of A** is the polynomial $p_A(x) = |xI_n - A|$.

Clearly, $p_A(x)$ is a polynomial of degree $n \implies$ it has at most n real roots. Hence, from above Theorem we can say that

The eigenvalues of an $n \times n$ matrix A are precisely the real roots of the characteristic polynomial $p_A(x)$.



Example 2



Example 2

Q:. Find the characteristic polynomial and eigenvalues of the matrix.



Example 2

Q:. Find the characteristic polynomial and eigenvalues of the matrix.

$$\begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & -3 \\ 0 & 0 & -5 \end{bmatrix}$$



Sol. For $A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & -3 \\ 0 & 0 & -5 \end{bmatrix}$, characteristic polynomial



Sol. For $A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & -3 \\ 0 & 0 & -5 \end{bmatrix}$, characteristic polynomial

$$p_A(x) = |xI_3 - A| =$$


Sol. For $A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & -3 \\ 0 & 0 & -5 \end{bmatrix}$, characteristic polynomial

$$p_A(x) = |xI_3 - A| =$$

$$\begin{vmatrix} x-1 & 0 & -1 \\ 0 & x-2 & 3 \\ 0 & 0 & x+5 \end{vmatrix} = (x-1)(x-2)(x+5)$$



Sol. For $A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & -3 \\ 0 & 0 & -5 \end{bmatrix}$, characteristic polynomial

$$p_A(x) = |xI_3 - A| =$$

$$\begin{vmatrix} x-1 & 0 & -1 \\ 0 & x-2 & 3 \\ 0 & 0 & x+5 \end{vmatrix} = (x-1)(x-2)(x+5)$$

Now, eigenvalues of A are the roots of $p_A(x)$.



Sol. For $A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & -3 \\ 0 & 0 & -5 \end{bmatrix}$, characteristic polynomial

$$p_A(x) = |xI_3 - A| =$$

$$\begin{vmatrix} x-1 & 0 & -1 \\ 0 & x-2 & 3 \\ 0 & 0 & x+5 \end{vmatrix} = (x-1)(x-2)(x+5)$$

Now, eigenvalues of A are the roots of $p_A(x)$. Hence, eigenvalues are $\lambda_1 = 1$, $\lambda_2 = 2$ and $\lambda_3 = -5$.



Algebraic Multiplicity of an Eigenvalue



Algebraic Multiplicity of an Eigenvalue

Let A be an $n \times n$ matrix and λ be an eigenvalue for A .



Algebraic Multiplicity of an Eigenvalue

Let A be an $n \times n$ matrix and λ be an eigenvalue for A . Suppose that $(x - \lambda)^k$ is the highest power of $(x - \lambda)$ that divides $p_A(x)$.



Algebraic Multiplicity of an Eigenvalue

Let A be an $n \times n$ matrix and λ be an eigenvalue for A . Suppose that $(x - \lambda)^k$ is the highest power of $(x - \lambda)$ that divides $p_A(x)$. Then k is called the **algebraic multiplicity** of λ .



Algebraic Multiplicity of an Eigenvalue

Let A be an $n \times n$ matrix and λ be an eigenvalue for A . Suppose that $(x - \lambda)^k$ is the highest power of $(x - \lambda)$ that divides $p_A(x)$. Then k is called the **algebraic multiplicity of λ** .

Corresponding to Example 2, the algebraic multiplicities of $\lambda_1 = 1$, $\lambda_2 = 2$ and $\lambda_3 = -5$ are 1, 1, 1 respectively.



Example 3



Example 3

Q:. Find all eigenvalues of the given matrix and their corresponding algebraic multiplicities.



Example 3

Q:. Find all eigenvalues of the given matrix and their corresponding algebraic multiplicities.

$$\begin{bmatrix} 4 & 0 & -2 \\ 6 & 2 & -6 \\ 4 & 0 & -2 \end{bmatrix}$$



Sol. For $A = \begin{bmatrix} 4 & 0 & -2 \\ 6 & 2 & -6 \\ 4 & 0 & -2 \end{bmatrix}$, characteristic polynomial



Sol. For $A = \begin{bmatrix} 4 & 0 & -2 \\ 6 & 2 & -6 \\ 4 & 0 & -2 \end{bmatrix}$, characteristic polynomial

$$p_A(x) = |xI_3 - A| =$$


Sol. For $A = \begin{bmatrix} 4 & 0 & -2 \\ 6 & 2 & -6 \\ 4 & 0 & -2 \end{bmatrix}$, characteristic polynomial

$$p_A(x) = |xI_3 - A| =$$

$$\begin{vmatrix} x-4 & 0 & 2 \\ -6 & x-2 & 6 \\ -4 & 0 & x+2 \end{vmatrix} = x(x-2)^2$$



Sol. For $A = \begin{bmatrix} 4 & 0 & -2 \\ 6 & 2 & -6 \\ 4 & 0 & -2 \end{bmatrix}$, characteristic polynomial
 $p_A(x) = |xI_3 - A| =$

$$\begin{vmatrix} x-4 & 0 & 2 \\ -6 & x-2 & 6 \\ -4 & 0 & x+2 \end{vmatrix} = x(x-2)^2$$

Now, the eigenvalues of A are the roots of $p_A(x)$, i.e., eigenvalues are $\lambda_1 = 0$ and $\lambda_2 = 2$.



Sol. For $A = \begin{bmatrix} 4 & 0 & -2 \\ 6 & 2 & -6 \\ 4 & 0 & -2 \end{bmatrix}$, characteristic polynomial
 $p_A(x) = |xI_3 - A| =$

$$\begin{vmatrix} x-4 & 0 & 2 \\ -6 & x-2 & 6 \\ -4 & 0 & x+2 \end{vmatrix} = x(x-2)^2$$

Now, the eigenvalues of A are the roots of $p_A(x)$, i.e., eigenvalues are $\lambda_1 = 0$ and $\lambda_2 = 2$. Hence, the algebraic multiplicity of λ_1 is one



Sol. For $A = \begin{bmatrix} 4 & 0 & -2 \\ 6 & 2 & -6 \\ 4 & 0 & -2 \end{bmatrix}$, characteristic polynomial

$$p_A(x) = |xI_3 - A| =$$

$$\begin{vmatrix} x-4 & 0 & 2 \\ -6 & x-2 & 6 \\ -4 & 0 & x+2 \end{vmatrix} = x(x-2)^2$$

Now, the eigenvalues of A are the roots of $p_A(x)$, i.e., eigenvalues are $\lambda_1 = 0$ and $\lambda_2 = 2$. Hence, the algebraic multiplicity of λ_1 is one and λ_2 is two.



Example 4



Example 4

Q:. Find the characteristic polynomial and eigenvalues of the matrix.



Example 4

Q:. Find the characteristic polynomial and eigenvalues of the matrix.

$$A = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$$



Example 4

Q:. Find the characteristic polynomial and eigenvalues of the matrix.

$$A = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$$

Sol. There are no eigenvalues of A .



Example 5



Example 5

Q:.. Let

$$A = \begin{bmatrix} 1 & 2 & -1 \\ 1 & 0 & 1 \\ 4 & -4 & 5 \end{bmatrix}$$



Example 5

Q.: Let

$$A = \begin{bmatrix} 1 & 2 & -1 \\ 1 & 0 & 1 \\ 4 & -4 & 5 \end{bmatrix}$$

Verify that

- $v = \begin{pmatrix} -1 \\ 1 \\ 2 \end{pmatrix}$ is an eigenvector of A .



Example 5

Q.: Let

$$A = \begin{bmatrix} 1 & 2 & -1 \\ 1 & 0 & 1 \\ 4 & -4 & 5 \end{bmatrix}$$

Verify that

- $v = \begin{pmatrix} -1 \\ 1 \\ 2 \end{pmatrix}$ is an eigenvector of A . Find the corresponding eigenvalue.



Example 5

Q: Let

$$A = \begin{bmatrix} 1 & 2 & -1 \\ 1 & 0 & 1 \\ 4 & -4 & 5 \end{bmatrix}$$

Verify that

- $v = \begin{pmatrix} -1 \\ 1 \\ 2 \end{pmatrix}$ is an eigenvector of A . Find the corresponding eigenvalue.
- 2 is an eigenvalue of A .



Sol. v is an eigenvector if $Av = \lambda v$ for some scalar λ .



Sol. v is an eigenvector if $Av = \lambda v$ for some scalar λ .
Now it is easy to verify that $Av = v = 1v \implies$



Sol. v is an eigenvector if $Av = \lambda v$ for some scalar λ .
Now it is easy to verify that $Av = v = 1v \implies v$ is an eigenvector of A and the corresponding eigenvalue is 1.



Sol. v is an eigenvector if $Av = \lambda v$ for some scalar λ .

Now it is easy to verify that $Av = v = 1v \implies v$ is an eigenvector of A and the corresponding eigenvalue is 1.

2 is an eigenvalue of A if there exists a non-zero vector v such that $Av = 2v \implies$



Sol. v is an eigenvector if $Av = \lambda v$ for some scalar λ .
Now it is easy to verify that $Av = v = 1v \implies v$ is an eigenvector of A and the corresponding eigenvalue is 1.

2 is an eigenvalue of A if there exists a non-zero vector v such that $Av = 2v \implies (A - 2I)v = 0$ has a non-zero solution.



Sol. v is an eigenvector if $Av = \lambda v$ for some scalar λ .
Now it is easy to verify that $Av = v = 1v \implies v$ is an eigenvector of A and the corresponding eigenvalue is 1.

2 is an eigenvalue of A if there exists a non-zero vector v such that $Av = 2v \implies (A - 2I)v = 0$ has a non-zero solution.

It can be verified that the homogeneous system of equation $(A - 2I)v = 0$ has infinitely many solutions.



Sol. v is an eigenvector if $Av = \lambda v$ for some scalar λ .
Now it is easy to verify that $Av = v = 1v \implies v$ is an eigenvector of A and the corresponding eigenvalue is 1.

2 is an eigenvalue of A if there exists a non-zero vector v such that $Av = 2v \implies (A - 2I)v = 0$ has a non-zero solution.

It can be verified that the homogeneous system of equation $(A - 2I)v = 0$ has infinitely many solutions.
Hence, 2 is an eigenvalue of A .



Eigenspace



Eigenspace

Let A be $n \times n$ matrix and λ be an eigenvalue for A .



Eigenspace

Let A be $n \times n$ matrix and λ be an eigenvalue for A .

Then the set $E_\lambda = \{X | AX = \lambda X\}$ is called the eigenspace of λ ,



Eigenspace

Let A be $n \times n$ matrix and λ be an eigenvalue for A .

Then the set $E_\lambda = \{X | AX = \lambda X\}$ is called the **eigenspace** of λ , i.e., E_λ consists of all set of all eigenvectors for A associated with λ ,



Eigenspace

Let A be $n \times n$ matrix and λ be an eigenvalue for A .

Then the set $E_\lambda = \{X | AX = \lambda X\}$ is called the **eigenspace** of λ , i.e., E_λ consists of all set of all eigenvectors for A associated with λ , together with zero vector $\mathbf{0}$.



Eigenspace

Let A be $n \times n$ matrix and λ be an eigenvalue for A . Then the set $E_\lambda = \{X | AX = \lambda X\}$ is called the **eigenspace** of λ , i.e., E_λ consists of all set of all eigenvectors for A associated with λ , together with zero vector $\mathbf{0}$.

Theorem: Let A be an $n \times n$ matrix and λ be an eigenvalue for A having eigenspace E_λ .



Eigenspace

Let A be $n \times n$ matrix and λ be an eigenvalue for A . Then the set $E_\lambda = \{X | AX = \lambda X\}$ is called the **eigenspace** of λ , i.e., E_λ consists of all set of all eigenvectors for A associated with λ , together with zero vector $\mathbf{0}$.

Theorem: Let A be an $n \times n$ matrix and λ be an eigenvalue for A having eigenspace E_λ . Then E_λ is a subspace of \mathbb{R}^n .



Example 6



Example 6

Q:. Find all eigenvalues and corresponding eigenspace for the given matrix.



Example 6

Q:. Find all eigenvalues and corresponding eigenspace for the given matrix.

$$\begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix}$$



Sol. For $A = \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix}$, characteristic polynomial

$$p_A(x) = |xI_2 - A| = \begin{vmatrix} x-1 & -3 \\ 0 & x-1 \end{vmatrix} = (x-1)^2.$$



Sol. For $A = \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix}$, characteristic polynomial

$$p_A(x) = |xI_2 - A| = \begin{vmatrix} x-1 & -3 \\ 0 & x-1 \end{vmatrix} = (x-1)^2. \text{ Hence,} \\ \text{eigenvalues are } \lambda = 1, 1.$$



Sol. For $A = \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix}$, characteristic polynomial

$$p_A(x) = |xI_2 - A| = \begin{vmatrix} x-1 & -3 \\ 0 & x-1 \end{vmatrix} = (x-1)^2. \text{ Hence,}$$

eigenvalues are $\lambda = 1, 1$.

To compute eigenspace E_1 for $\lambda = 1$, we need to solve the homogeneous system



Sol. For $A = \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix}$, characteristic polynomial

$$p_A(x) = |xI_2 - A| = \begin{vmatrix} x-1 & -3 \\ 0 & x-1 \end{vmatrix} = (x-1)^2. \text{ Hence,}$$

eigenvalues are $\lambda = 1, 1$.

To compute eigenspace E_1 for $\lambda = 1$, we need to solve the homogeneous system $\lambda I_2 - AX = 0$, i.e., $I_2 - AX = 0$.



The augmented matrix is $[I_2 - A|0]$, i.e., $\left[\begin{array}{cc|c} 0 & -3 & 0 \\ 0 & 0 & 0 \end{array} \right]$,



The augmented matrix is $[I_2 - A|0]$, i.e., $\left[\begin{array}{cc|c} 0 & -3 & 0 \\ 0 & 0 & 0 \end{array} \right]$,
which reduces to $\left[\begin{array}{cc|c} 0 & 1 & 0 \\ 0 & 0 & 0 \end{array} \right]$.



The augmented matrix is $[I_2 - A|0]$, i.e., $\left[\begin{array}{cc|c} 0 & -3 & 0 \\ 0 & 0 & 0 \end{array} \right]$,

which reduces to $\left[\begin{array}{cc|c} 0 & 1 & 0 \\ 0 & 0 & 0 \end{array} \right]$.

The associated system is



The augmented matrix is $[I_2 - A|0]$, i.e., $\left[\begin{array}{cc|c} 0 & -3 & 0 \\ 0 & 0 & 0 \end{array} \right]$,

which reduces to $\left[\begin{array}{cc|c} 0 & 1 & 0 \\ 0 & 0 & 0 \end{array} \right]$.

The associated system is $\begin{cases} x_2 = 0 \\ 0 = 0 \end{cases}$



The augmented matrix is $[I_2 - A|0]$, i.e., $\left[\begin{array}{cc|c} 0 & -3 & 0 \\ 0 & 0 & 0 \end{array} \right]$,

which reduces to $\left[\begin{array}{cc|c} 0 & 1 & 0 \\ 0 & 0 & 0 \end{array} \right]$.

The associated system is $\begin{cases} x_2 = 0 \\ 0 = 0 \end{cases}$

Since column 1 is not a pivot column, x_1 is an independent variable.



The augmented matrix is $[I_2 - A|0]$, i.e., $\left[\begin{array}{cc|c} 0 & -3 & 0 \\ 0 & 0 & 0 \end{array} \right]$,

which reduces to $\left[\begin{array}{cc|c} 0 & 1 & 0 \\ 0 & 0 & 0 \end{array} \right]$.

The associated system is $\begin{cases} x_2 = 0 \\ 0 = 0 \end{cases}$

Since column 1 is not a pivot column, x_1 is an independent variable. Let $x_1 = a$.



The augmented matrix is $[I_2 - A|0]$, i.e., $\left[\begin{array}{cc|c} 0 & -3 & 0 \\ 0 & 0 & 0 \end{array} \right]$,

which reduces to $\left[\begin{array}{cc|c} 0 & 1 & 0 \\ 0 & 0 & 0 \end{array} \right]$.

The associated system is $\begin{cases} x_2 = 0 \\ 0 = 0 \end{cases}$

Since column 1 is not a pivot column, x_1 is an independent variable. Let $x_1 = a$.

Then $E_1 = \{[a, 0] | a \in \mathbb{R}\} = \{a[1, 0] | a \in \mathbb{R}\}$.



Geometric Multiplicity (G.M.) of an Eigenvalue



Geometric Multiplicity (G.M.) of an Eigenvalue

The **geometric multiplicity** of an eigenvalue λ is the dimension of its corresponding eigenspace E_λ i.e.



Geometric Multiplicity (G.M.) of an Eigenvalue

The **geometric multiplicity** of an eigenvalue λ is the dimension of its corresponding eigenspace E_λ i.e.

$$\text{G.M. of } \lambda = \dim(E_\lambda).$$



Example 7



Example 7

Q: Consider

$$A = \begin{bmatrix} 4 & 0 & -2 \\ 6 & 2 & -6 \\ 4 & 0 & -2 \end{bmatrix}$$



Example 7

Q.: Consider

$$A = \begin{bmatrix} 4 & 0 & -2 \\ 6 & 2 & -6 \\ 4 & 0 & -2 \end{bmatrix}$$

- Find all the eigenvalue of A and compute their algebraic multiplicity.
- Find eigenspaces corresponding to each of the eigenvalues of A and compute their geometric multiplicity.



For $A = \begin{bmatrix} 4 & 0 & -2 \\ 6 & 2 & -6 \\ 4 & 0 & -2 \end{bmatrix}$, characteristic polynomial

$$p_A(x) = |xI_3 - A| = \begin{vmatrix} x-4 & 0 & 2 \\ -6 & x-2 & 6 \\ -4 & 0 & x+2 \end{vmatrix} = x(x-2)^2.$$



For $A = \begin{bmatrix} 4 & 0 & -2 \\ 6 & 2 & -6 \\ 4 & 0 & -2 \end{bmatrix}$, characteristic polynomial

$$p_A(x) = |xI_3 - A| = \begin{vmatrix} x-4 & 0 & 2 \\ -6 & x-2 & 6 \\ -4 & 0 & x+2 \end{vmatrix} = x(x-2)^2. \text{ Hence,}$$

eigenvalues are $\lambda_1 = 0$ and $\lambda_2 = 2$.



For $A = \begin{bmatrix} 4 & 0 & -2 \\ 6 & 2 & -6 \\ 4 & 0 & -2 \end{bmatrix}$, characteristic polynomial

$$p_A(x) = |xI_3 - A| = \begin{vmatrix} x-4 & 0 & 2 \\ -6 & x-2 & 6 \\ -4 & 0 & x+2 \end{vmatrix} = x(x-2)^2. \text{ Hence,}$$

eigenvalues are $\lambda_1 = 0$ and $\lambda_2 = 2$.

To compute eigenspace E_0 for $\lambda = 0$, we need to solve the homogeneous system



For $A = \begin{bmatrix} 4 & 0 & -2 \\ 6 & 2 & -6 \\ 4 & 0 & -2 \end{bmatrix}$, characteristic polynomial

$$p_A(x) = |xI_3 - A| = \begin{vmatrix} x-4 & 0 & 2 \\ -6 & x-2 & 6 \\ -4 & 0 & x+2 \end{vmatrix} = x(x-2)^2. \text{ Hence,}$$

eigenvalues are $\lambda_1 = 0$ and $\lambda_2 = 2$.

To compute eigenspace E_0 for $\lambda = 0$, we need to solve the homogeneous system $\lambda I_3 - AX = 0$, i.e., $-AX = 0$.



The augmented matrix is $[-A|0]$, i.e., $\left[\begin{array}{ccc|c} -4 & 0 & 2 & 0 \\ -6 & -2 & 6 & 0 \\ -4 & 0 & 2 & 0 \end{array} \right]$,

which reduces to



The augmented matrix is $[-A|0]$, i.e., $\left[\begin{array}{ccc|c} -4 & 0 & 2 & 0 \\ -6 & -2 & 6 & 0 \\ -4 & 0 & 2 & 0 \end{array} \right],$

which reduces to $\left[\begin{array}{ccc|c} 1 & 0 & -1/2 & 0 \\ 0 & 1 & -3/2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right].$



The associated system is



The associated system is
$$\begin{cases} x_1 - \frac{1}{2}x_3 = 0 \\ x_2 - \frac{3}{2}x_3 = 0 \\ 0 = 0 \end{cases}$$



The associated system is
$$\begin{cases} x_1 - \frac{1}{2}x_3 = 0 \\ x_2 - \frac{3}{2}x_3 = 0 \\ 0 = 0 \end{cases}$$

Since column 3 is not a pivot column, x_3 is an independent variable.



The associated system is
$$\begin{cases} x_1 - \frac{1}{2}x_3 = 0 \\ x_2 - \frac{3}{2}x_3 = 0 \\ 0 = 0 \end{cases}$$

Since column 3 is not a pivot column, x_3 is an independent variable. Let $x_3 = 2c \implies$



The associated system is
$$\begin{cases} x_1 - \frac{1}{2}x_3 = 0 \\ x_2 - \frac{3}{2}x_3 = 0 \\ 0 = 0 \end{cases}$$

Since column 3 is not a pivot column, x_3 is an independent variable. Let $x_3 = 2c \implies x_1 = c, x_2 = 3c$.



The associated system is
$$\begin{cases} x_1 - \frac{1}{2}x_3 = 0 \\ x_2 - \frac{3}{2}x_3 = 0 \\ 0 = 0 \end{cases}$$

Since column 3 is not a pivot column, x_3 is an independent variable. Let $x_3 = 2c \implies x_1 = c, x_2 = 3c$.

Then $E_0 = \{[c, 3c, 2c] | c \in \mathbb{R}\} = \{c[1, 3, 2] | c \in \mathbb{R}\}$.



The associated system is
$$\begin{cases} x_1 - \frac{1}{2}x_3 = 0 \\ x_2 - \frac{3}{2}x_3 = 0 \\ 0 = 0 \end{cases}$$

Since column 3 is not a pivot column, x_3 is an independent variable. Let $x_3 = 2c \implies x_1 = c, x_2 = 3c$.

Then $E_0 = \{[c, 3c, 2c] | c \in \mathbb{R}\} = \{c[1, 3, 2] | c \in \mathbb{R}\}$.

Now $E_0 = \text{span}\{[1, 3, 2]\} = \text{span}(B)$.



The associated system is
$$\begin{cases} x_1 - \frac{1}{2}x_3 = 0 \\ x_2 - \frac{3}{2}x_3 = 0 \\ 0 = 0 \end{cases}$$

Since column 3 is not a pivot column, x_3 is an independent variable. Let $x_3 = 2c \implies x_1 = c, x_2 = 3c$.

Then $E_0 = \{[c, 3c, 2c] | c \in \mathbb{R}\} = \{c[1, 3, 2] | c \in \mathbb{R}\}$.

Now $E_0 = \text{span}\{[1, 3, 2]\} = \text{span}(B)$. Since, B is LI, it is a basis for E_0 . Note that

G.M. of eigenvalue $0 = \dim(E_0) = 1$.



To compute eigenspace E_2 for $\lambda = 2$, we need to solve the homogeneous system



To compute eigenspace E_2 for $\lambda = 2$, we need to solve the homogeneous system $\lambda I_3 - AX = 0$, i.e., $2I_3 - AX = 0$.



To compute eigenspace E_2 for $\lambda = 2$, we need to solve the homogeneous system $\lambda I_3 - AX = 0$, i.e., $2I_3 - AX = 0$.

The augmented matrix is $[-A|0]$, i.e., $\left[\begin{array}{ccc|c} -2 & 0 & 2 & 0 \\ -6 & 0 & 6 & 0 \\ -4 & 0 & 4 & 0 \end{array} \right]$,

which reduces to



To compute eigenspace E_2 for $\lambda = 2$, we need to solve the homogeneous system $\lambda I_3 - AX = 0$, i.e., $2I_3 - AX = 0$.

The augmented matrix is $[-A|0]$, i.e., $\left[\begin{array}{ccc|c} -2 & 0 & 2 & 0 \\ -6 & 0 & 6 & 0 \\ -4 & 0 & 4 & 0 \end{array} \right]$,

which reduces to $\left[\begin{array}{ccc|c} 1 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$.



The associated system is



The associated system is
$$\begin{cases} x_1 - x_3 = 0 \\ 0 = 0 \\ 0 = 0 \end{cases}$$



The associated system is $\begin{cases} x_1 - x_3 = 0 \\ 0 = 0 \\ 0 = 0 \end{cases}$

Since columns 2 and 3 are not pivot columns, x_2 and x_3 are independent variables.



The associated system is $\begin{cases} x_1 - x_3 = 0 \\ 0 = 0 \\ 0 = 0 \end{cases}$

Since columns 2 and 3 are not pivot columns, x_2 and x_3 are independent variables. Let $x_2 = b$ and $x_3 = c \implies$



The associated system is
$$\begin{cases} x_1 - x_3 = 0 \\ 0 = 0 \\ 0 = 0 \end{cases}$$

Since columns 2 and 3 are not pivot columns, x_2 and x_3 are independent variables. Let $x_2 = b$ and $x_3 = c \implies x_1 = c$.



The associated system is
$$\begin{cases} x_1 - x_3 = 0 \\ 0 = 0 \\ 0 = 0 \end{cases}$$

Since columns 2 and 3 are not pivot columns, x_2 and x_3 are independent variables. Let $x_2 = b$ and $x_3 = c \implies x_1 = c$.

Then

$$E_2 = \{[c, b, c] | b, c \in \mathbb{R}\} = \{b[0, 1, 0] + c[1, 0, 1] | b, c \in \mathbb{R}\}.$$



The associated system is
$$\begin{cases} x_1 - x_3 = 0 \\ 0 = 0 \\ 0 = 0 \end{cases}$$

Since columns 2 and 3 are not pivot columns, x_2 and x_3 are independent variables. Let $x_2 = b$ and $x_3 = c \implies x_1 = c$.

Then

$$E_2 = \{[c, b, c] | b, c \in \mathbb{R}\} = \{b[0, 1, 0] + c[1, 0, 1] | b, c \in \mathbb{R}\}.$$

$$\text{Now } E_2 = \text{span}\{[0, 1, 0], [1, 0, 1]\} = \text{span}(B).$$



The associated system is
$$\begin{cases} x_1 - x_3 = 0 \\ 0 = 0 \\ 0 = 0 \end{cases}$$

Since columns 2 and 3 are not pivot columns, x_2 and x_3 are independent variables. Let $x_2 = b$ and $x_3 = c \implies x_1 = c$.

Then

$$E_2 = \{[c, b, c] | b, c \in \mathbb{R}\} = \{b[0, 1, 0] + c[1, 0, 1] | b, c \in \mathbb{R}\}.$$

Now $E_2 = \text{span}\{[0, 1, 0], [1, 0, 1]\} = \text{span}(B)$. Since, B is LI, it is a basis for E_2 .



The associated system is
$$\begin{cases} x_1 - x_3 = 0 \\ 0 = 0 \\ 0 = 0 \end{cases}$$

Since columns 2 and 3 are not pivot columns, x_2 and x_3 are independent variables. Let $x_2 = b$ and $x_3 = c \implies x_1 = c$.

Then

$$E_2 = \{[c, b, c] | b, c \in \mathbb{R}\} = \{b[0, 1, 0] + c[1, 0, 1] | b, c \in \mathbb{R}\}.$$

Now $E_2 = \text{span}\{[0, 1, 0], [1, 0, 1]\} = \text{span}(B)$. Since, B is LI, it is a basis for E_2 . Note that

$$\text{G.M. of eigenvalue } 2 = \dim(E_2) = 2.$$



Exercise: Consider

$$A = \begin{bmatrix} 1 & 1 & 3 \\ 1 & 5 & 1 \\ 3 & 1 & 1 \end{bmatrix}$$



Exercise: Consider

$$A = \begin{bmatrix} 1 & 1 & 3 \\ 1 & 5 & 1 \\ 3 & 1 & 1 \end{bmatrix}$$

- Find all the eigenvalue of A and compute their algebraic multiplicity.
- Find eigenspaces corresponding to each of the eigenvalues of A and compute their geometric multiplicity.



Exercise: Consider

$$A = \begin{bmatrix} 1 & 1 & 3 \\ 1 & 5 & 1 \\ 3 & 1 & 1 \end{bmatrix}$$

- Find all the eigenvalue of A and compute their algebraic multiplicity.
- Find eigenspaces corresponding to each of the eigenvalues of A and compute their geometric multiplicity.

Sol. $\lambda = -2, 3, 6$ where $E_{-2} = \{[-z, 0, z] | z \in \mathbb{R}\}$,
 $E_3 = \{[z, -z, z] | z \in \mathbb{R}\}$ and $E_6 = \{[z, 2z, z] | z \in \mathbb{R}\}$.



Exercise: Consider

$$A = \begin{bmatrix} -9 & 4 & 4 \\ -8 & 3 & 4 \\ -16 & 8 & 7 \end{bmatrix}$$



Exercise: Consider

$$A = \begin{bmatrix} -9 & 4 & 4 \\ -8 & 3 & 4 \\ -16 & 8 & 7 \end{bmatrix}$$

- Find all the eigenvalue of A and compute their algebraic multiplicity.
- Find eigenspaces corresponding to each of the eigenvalues of A and compute their geometric multiplicity.



Exercise: Consider

$$A = \begin{bmatrix} -9 & 4 & 4 \\ -8 & 3 & 4 \\ -16 & 8 & 7 \end{bmatrix}$$

- Find all the eigenvalue of A and compute their algebraic multiplicity.
- Find eigenspaces corresponding to each of the eigenvalues of A and compute their geometric multiplicity.

Sol. $\lambda = -1, -1, 3$ where $E_{-1} = \{[x, 2x - z, z] | x, z \in \mathbb{R}\}$ and $E_3 = \{[z/2, z/2, z] | z \in \mathbb{R}\}$.



Result: Let $\lambda_1, \lambda_2, \dots, \lambda_n$ be n eigenvalues of a $n \times n$ matrix A .



Result: Let $\lambda_1, \lambda_2, \dots, \lambda_n$ be n eigenvalues of a $n \times n$ matrix A . Then

$$\det A = \lambda_1 \lambda_2 \cdots \lambda_n$$



Result: Let $\lambda_1, \lambda_2, \dots, \lambda_n$ be n eigenvalues of a $n \times n$ matrix A . Then

$$\det A = \lambda_1 \lambda_2 \cdots \lambda_n$$

Theorem: Let A be a square matrix with eigenvalue λ and corresponding eigenvector X .

- If λ is an eigenvalue of a matrix A , then for any positive integer n , λ^n is an eigenvalue of A^n with corresponding eigenvector X .



Result: Let $\lambda_1, \lambda_2, \dots, \lambda_n$ be n eigenvalues of a $n \times n$ matrix A . Then

$$\det A = \lambda_1 \lambda_2 \cdots \lambda_n$$

Theorem: Let A be a square matrix with eigenvalue λ and corresponding eigenvector X .

- If λ is an eigenvalue of a matrix A , then for any positive integer n , λ^n is an eigenvalue of A^n with corresponding eigenvector X .
- If A is nonsingular, then for any integer n , λ^n is an eigenvalue of A^n with corresponding eigenvector X .



Example 8

Q:. Let A be a 2×2 matrix with eigenvectors $v_1 = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$ and $v_2 = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$ corresponding to eigenvalues $\lambda_1 = 1, \lambda_2 = 4$.



Example 8

Q:. Let A be a 2×2 matrix with eigenvectors $v_1 = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$ and $v_2 = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$ corresponding to eigenvalues $\lambda_1 = 1, \lambda_2 = 4$.
Let $X = \begin{pmatrix} -1 \\ -5 \end{pmatrix}$.



Example 8

Q:. Let A be a 2×2 matrix with eigenvectors $v_1 = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$ and $v_2 = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$ corresponding to eigenvalues $\lambda_1 = 1, \lambda_2 = 4$. Let $X = \begin{pmatrix} -1 \\ -5 \end{pmatrix}$. Find $A^4 X$ without computing the matrix A .



Example 8

Q:. Let A be a 2×2 matrix with eigenvectors $v_1 = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$ and $v_2 = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$ corresponding to eigenvalues $\lambda_1 = 1, \lambda_2 = 4$. Let $X = \begin{pmatrix} -1 \\ -5 \end{pmatrix}$. Find $A^4 X$ without computing the matrix A .

Sol. $\begin{pmatrix} -1021 \\ -515 \end{pmatrix}$



Example 9

Q:. Let A be an $n \times n$ matrix and $\lambda \in \mathbb{R}$ is not an eigenvalue for A .



Example 9

Q:. Let A be an $n \times n$ matrix and $\lambda \in \mathbb{R}$ is not an eigenvalue for A . Determine exactly which vectors are in $S = \{X \in \mathbb{R}^n | AX = \lambda X\}$.



Example 9

Q:. Let A be an $n \times n$ matrix and $\lambda \in \mathbb{R}$ is not an eigenvalue for A . Determine exactly which vectors are in $S = \{X \in \mathbb{R}^n | AX = \lambda X\}$. Is this set a subspace of \mathbb{R}^n .



Example 9

Q.: Let A be an $n \times n$ matrix and $\lambda \in \mathbb{R}$ is not an eigenvalue for A . Determine exactly which vectors are in $S = \{X \in \mathbb{R}^n | AX = \lambda X\}$. Is this set a subspace of \mathbb{R}^n .

Sol. If λ is not an eigenvalue for A , then, by the definition of an eigenvalue, there are no nonzero vectors X such that $AX = \lambda X$.



Example 9

Q.: Let A be an $n \times n$ matrix and $\lambda \in \mathbb{R}$ is not an eigenvalue for A . Determine exactly which vectors are in $S = \{X \in \mathbb{R}^n | AX = \lambda X\}$. Is this set a subspace of \mathbb{R}^n .

Sol. If λ is not an eigenvalue for A , then, by the definition of an eigenvalue, there are no nonzero vectors X such that $AX = \lambda X$. Thus, no nonzero vector can be in S .



Example 9

Q.: Let A be an $n \times n$ matrix and $\lambda \in \mathbb{R}$ is not an eigenvalue for A . Determine exactly which vectors are in $S = \{X \in \mathbb{R}^n | AX = \lambda X\}$. Is this set a subspace of \mathbb{R}^n .

Sol. If λ is not an eigenvalue for A , then, by the definition of an eigenvalue, there are no nonzero vectors X such that $AX = \lambda X$. Thus, no nonzero vector can be in S . However, $A0 = 0 = \lambda 0$, hence $0 \in S$.



Example 9

Q.: Let A be an $n \times n$ matrix and $\lambda \in \mathbb{R}$ is not an eigenvalue for A . Determine exactly which vectors are in $S = \{X \in \mathbb{R}^n | AX = \lambda X\}$. Is this set a subspace of \mathbb{R}^n .

Sol. If λ is not an eigenvalue for A , then, by the definition of an eigenvalue, there are no nonzero vectors X such that $AX = \lambda X$. Thus, no nonzero vector can be in S . However, $A0 = 0 = \lambda 0$, hence $0 \in S$. $S = \{0\}$ is a trivial subspace of \mathbb{R}^n .

