

MATHEMATICS-II (MATH F112)

Dr. Krishnendra Shekhawat

BITS PILANI
Department of Mathematics



Section 5.3

The Dimension Theorem



Kernel of a LT



Kernel of a LT

Let $L : V \rightarrow W$ be a LT.



Kernel of a LT

Let $L : V \rightarrow W$ be a LT. The **kernel** of L , denoted by $\ker(L)$,



Kernel of a LT

Let $L : V \rightarrow W$ be a LT. The **kernel** of L , denoted by $\ker(L)$, is the subset of all vectors in V that map to $\mathbf{0}_W$, i.e.,



Kernel of a LT

Let $L : V \rightarrow W$ be a LT. The **kernel** of L , denoted by $\ker(L)$, is the subset of all vectors in V that map to $\mathbf{0}_W$, i.e.,

$$\ker(L) = \{v \in V | L(v) = \mathbf{0}_W\}$$



Example 1



Example 1

Q:. Let $L : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a LT given by $L([x, y, z]) = [0, y]$.



Example 1

Q.: Let $L : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a LT given by $L([x, y, z]) = [0, y]$. Find $\ker(L)$.



Example 1

Q.: Let $L : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a LT given by $L([x, y, z]) = [0, y]$. Find $\ker(L)$.

Sol.

$$\left\{ \ker(L) = \{[x, y, z] \in \mathbb{R}^3 \mid L([x, y, z]) = \mathbf{0}_{\mathbb{R}^2}\} \right.$$



Example 1

Q.: Let $L : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a LT given by $L([x, y, z]) = [0, y]$. Find $\ker(L)$.

Sol.

$$\left\{ \begin{aligned} \ker(L) &= \{[x, y, z] \in \mathbb{R}^3 \mid L([x, y, z]) = \mathbf{0}_{\mathbb{R}^2}\} \\ &= \{[x, y, z] \in \mathbb{R}^3 \mid [0, y] = [0, 0]\} \end{aligned} \right.$$



Example 1

Q.: Let $L : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a LT given by $L([x, y, z]) = [0, y]$. Find $\ker(L)$.

Sol.

$$\left\{ \begin{aligned} \ker(L) &= \{[x, y, z] \in \mathbb{R}^3 \mid L([x, y, z]) = \mathbf{0}_{\mathbb{R}^2}\} \\ &= \{[x, y, z] \in \mathbb{R}^3 \mid [0, y] = [0, 0]\} \\ &= \{[x, y, z] \in \mathbb{R}^3 \mid y = 0\} \end{aligned} \right.$$



Example 1

Q.: Let $L : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a LT given by $L([x, y, z]) = [0, y]$. Find $\ker(L)$.

Sol.

$$\left\{ \begin{aligned} \ker(L) &= \{[x, y, z] \in \mathbb{R}^3 \mid L([x, y, z]) = \mathbf{0}_{\mathbb{R}^2}\} \\ &= \{[x, y, z] \in \mathbb{R}^3 \mid [0, y] = [0, 0]\} \\ &= \{[x, y, z] \in \mathbb{R}^3 \mid y = 0\} \\ &= \{[x, 0, z] \mid x, z \in \mathbb{R}\} \end{aligned} \right.$$



In Example 1, **Note that**

$$\ker(L) = \{[x, 0, z] \in \mathbb{R}^3 \mid x, z \in \mathbb{R}\}$$



In Example 1, **Note that**

$$\ker(L) = \{[x, 0, z] \in \mathbb{R}^3 \mid x, z \in \mathbb{R}\}$$

is a subspace of the vector space \mathbb{R}^3 .



In Example 1, **Note that**

$$\ker(L) = \{[x, 0, z] \in \mathbb{R}^3 \mid x, z \in \mathbb{R}\}$$

is a subspace of the vector space \mathbb{R}^3 .

Result: If $L : V \rightarrow W$ is a LT, then $\ker(L)$ is a subspace of V .



Range of a LT



Range of a LT

Let $L : V \rightarrow W$ be a LT.



Range of a LT

Let $L : V \rightarrow W$ be a LT. The **range** of L , denoted by $\text{range}(L)$,



Range of a LT

Let $L : V \rightarrow W$ be a LT. The **range** of L , denoted by $\text{range}(L)$, is the subset of all vectors in W that are image of some vector in V , i.e.,



Range of a LT

Let $L : V \rightarrow W$ be a LT. The **range** of L , denoted by $\text{range}(L)$, is the subset of all vectors in W that are image of some vector in V , i.e.,

$$\text{range}(L) = \{L(v) | v \in V\}$$



Range of a LT

Let $L : V \rightarrow W$ be a LT. The **range** of L , denoted by $\text{range}(L)$, is the subset of all vectors in W that are image of some vector in V , i.e.,

$$\text{range}(L) = \{L(v) | v \in V\}$$

Thus a vector $w \in \text{range}(L)$



Range of a LT

Let $L : V \rightarrow W$ be a LT. The **range** of L , denoted by $\text{range}(L)$, is the subset of all vectors in W that are image of some vector in V , i.e.,

$$\text{range}(L) = \{L(v) | v \in V\}$$

Thus a vector $w \in \text{range}(L)$ if there exists some vector $v \in V$ such that $L(v) = w$.



Example 2



Example 2

Q:. Let $L : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a LT given by
 $L([x, y, z]) = ([0, y])$.



Example 2

Q:. Let $L : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a LT given by $L([x, y, z]) = ([0, y])$. Find $\text{range}(L)$.



Example 2

Q:. Let $L : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a LT given by $L([x, y, z]) = ([0, y])$. Find $\text{range}(L)$. Also, find basis for $\ker(L)$ and $\text{range}(L)$.



Example 2

Q:. Let $L : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a LT given by $L([x, y, z]) = ([0, y])$. Find $\text{range}(L)$. Also, find basis for $\ker(L)$ and $\text{range}(L)$.

Sol.

$$\left\{ \text{range}(L) = \{L([x, y, z]) \mid [x, y, z] \in \mathbb{R}^3\} \right.$$



Example 2

Q.: Let $L : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a LT given by $L([x, y, z]) = ([0, y])$. Find $\text{range}(L)$. Also, find basis for $\ker(L)$ and $\text{range}(L)$.

Sol.

$$\left\{ \begin{aligned} \text{range}(L) &= \{L([x, y, z]) \mid [x, y, z] \in \mathbb{R}^3\} \\ &= \{[0, y] \mid y \in \mathbb{R}\} \\ &= \{y[0, 1] \mid y \in \mathbb{R}\} \end{aligned} \right.$$



Note that $\text{range}(L) = \text{span}\{[0, 1]\}$.



Note that $\text{range}(L) = \text{span}\{[0, 1]\}$. Since $\{[0, 1]\}$ is LI subset of \mathbb{R}^2 (Why?).



Note that $\text{range}(L) = \text{span}\{[0, 1]\}$. Since $\{[0, 1]\}$ is LI subset of \mathbb{R}^2 (Why?). Thus, the set $B = \{[0, 1]\}$ is a basis of $\text{range}(L)$.



Note that $\text{range}(L) = \text{span}\{[0, 1]\}$. Since $\{[0, 1]\}$ is LI subset of \mathbb{R}^2 (Why?). Thus, the set $B = \{[0, 1]\}$ is a basis of $\text{range}(L)$.

Also, $\text{range}(L)$



Note that $\text{range}(L) = \text{span}\{[0, 1]\}$. Since $\{[0, 1]\}$ is LI subset of \mathbb{R}^2 (Why?). Thus, the set $B = \{[0, 1]\}$ is a basis of $\text{range}(L)$.

Also, $\text{range}(L)$ is a subspace of the vector space \mathbb{R}^2 .



Note that $\text{range}(L) = \text{span}\{[0, 1]\}$. Since $\{[0, 1]\}$ is LI subset of \mathbb{R}^2 (Why?). Thus, the set $B = \{[0, 1]\}$ is a basis of $\text{range}(L)$.

Also, $\text{range}(L)$ is a subspace of the vector space \mathbb{R}^2 .

Result: If $L : V \rightarrow W$ is a LT, then $\text{range}(L)$ is a subspace of W .



$$\left\{ \ker(L) = \{[x, 0, z] | x, z \in \mathbb{R}\} \right.$$



$$\left\{ \begin{aligned} \ker(L) &= \{[x, 0, z] | x, z \in \mathbb{R}\} \\ &= \{x[1, 0, 0] + z[0, 0, 1] | x, z \in \mathbb{R}\} \end{aligned} \right.$$



$$\left\{ \begin{aligned} \ker(L) &= \{[x, 0, z] | x, z \in \mathbb{R}\} \\ &= \{x[1, 0, 0] + z[0, 0, 1] | x, z \in \mathbb{R}\} \\ &= \text{span}\{[1, 0, 0], [0, 0, 1]\} \end{aligned} \right.$$



$$\left\{ \begin{aligned} \ker(L) &= \{[x, 0, z] | x, z \in \mathbb{R}\} \\ &= \{x[1, 0, 0] + z[0, 0, 1] | x, z \in \mathbb{R}\} \\ &= \text{span}\{[1, 0, 0], [0, 0, 1]\} \end{aligned} \right.$$

Now, $\{[1, 0, 0], [0, 0, 1]\}$ is a LI subset of \mathbb{R}^3 , hence, they form a basis for $\ker(L)$.



Example 3

Q:. Let $L : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a LT given by
 $L([x, y, z]) = [x - 2y, y + z]$.



Example 3

Q:. Let $L : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a LT given by
 $L([x, y, z]) = [x - 2y, y + z]$. Find $\ker(L)$ and $\text{range}(L)$.



Example 3

Q:. Let $L : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a LT given by $L([x, y, z]) = [x - 2y, y + z]$. Find $\ker(L)$ and $\text{range}(L)$. Also, find basis for $\ker(L)$ and $\text{range}(L)$.



Example 3

Q:. Let $L : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a LT given by $L([x, y, z]) = [x - 2y, y + z]$. Find $\ker(L)$ and $\text{range}(L)$. Also, find basis for $\ker(L)$ and $\text{range}(L)$.

Sol.

$$\left\{ \ker(L) = \{[x, y, z] \in \mathbb{R}^3 \mid L([x, y, z]) = \mathbf{0}_{\mathbb{R}^2}\} \right.$$



Example 3

Q:. Let $L : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a LT given by $L([x, y, z]) = [x - 2y, y + z]$. Find $\ker(L)$ and $\text{range}(L)$. Also, find basis for $\ker(L)$ and $\text{range}(L)$.

Sol.

$$\left\{ \begin{aligned} \ker(L) &= \{[x, y, z] \in \mathbb{R}^3 \mid L([x, y, z]) = \mathbf{0}_{\mathbb{R}^2}\} \\ &= \{[x, y, z] \in \mathbb{R}^3 \mid [x - 2y, y + z] = [0, 0]\} \end{aligned} \right.$$



Example 3

Q:. Let $L : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a LT given by $L([x, y, z]) = [x - 2y, y + z]$. Find $\ker(L)$ and $\text{range}(L)$. Also, find basis for $\ker(L)$ and $\text{range}(L)$.

Sol.

$$\left\{ \begin{aligned} \ker(L) &= \{[x, y, z] \in \mathbb{R}^3 \mid L([x, y, z]) = \mathbf{0}_{\mathbb{R}^2}\} \\ &= \{[x, y, z] \in \mathbb{R}^3 \mid [x - 2y, y + z] = [0, 0]\} \\ &= \{[x, y, z] \in \mathbb{R}^3 \mid x = 2y, z = -y\} \end{aligned} \right.$$



Example 3

Q:. Let $L : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a LT given by $L([x, y, z]) = [x - 2y, y + z]$. Find $\ker(L)$ and $\text{range}(L)$. Also, find basis for $\ker(L)$ and $\text{range}(L)$.

Sol.

$$\left\{ \begin{aligned} \ker(L) &= \{[x, y, z] \in \mathbb{R}^3 \mid L([x, y, z]) = \mathbf{0}_{\mathbb{R}^2}\} \\ &= \{[x, y, z] \in \mathbb{R}^3 \mid [x - 2y, y + z] = [0, 0]\} \\ &= \{[x, y, z] \in \mathbb{R}^3 \mid x = 2y, z = -y\} \\ &= \{[2y, y, -y] \mid y \in \mathbb{R}\} \end{aligned} \right.$$



Example 3

Q:. Let $L : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a LT given by $L([x, y, z]) = [x - 2y, y + z]$. Find $\ker(L)$ and $\text{range}(L)$. Also, find basis for $\ker(L)$ and $\text{range}(L)$.

Sol.

$$\left\{ \begin{aligned} \ker(L) &= \{[x, y, z] \in \mathbb{R}^3 \mid L([x, y, z]) = \mathbf{0}_{\mathbb{R}^2}\} \\ &= \{[x, y, z] \in \mathbb{R}^3 \mid [x - 2y, y + z] = [0, 0]\} \\ &= \{[x, y, z] \in \mathbb{R}^3 \mid x = 2y, z = -y\} \\ &= \{[2y, y, -y] \mid y \in \mathbb{R}\} \\ &= \text{span}\{[2, 1, -1]\} \end{aligned} \right.$$



Basis for $\ker(L)$ is $\{[2, 1, -1]\}$.



Basis for $\ker(L)$ is $\{[2, 1, -1]\}$.

$$\left\{ \begin{array}{l} \text{range}(L) = \{L([x, y, z]) \mid [x, y, z] \in \mathbb{R}^3\} \end{array} \right.$$



Basis for $\ker(L)$ is $\{[2, 1, -1]\}$.

$$\left\{ \begin{aligned} \text{range}(L) &= \{L([x, y, z]) | [x, y, z] \in \mathbb{R}^3\} \\ &= \{[x - 2y, y + z] | x, y, z \in \mathbb{R}\} \end{aligned} \right.$$



Basis for $\ker(L)$ is $\{[2, 1, -1]\}$.

$$\left\{ \begin{aligned} \text{range}(L) &= \{L([x, y, z]) | [x, y, z] \in \mathbb{R}^3\} \\ &= \{[x - 2y, y + z] | x, y, z \in \mathbb{R}\} \\ &= \{x[1, 0] + y[-2, 1] + z[0, 1] | x, y, z \in \mathbb{R}\} \end{aligned} \right.$$



Basis for $\ker(L)$ is $\{[2, 1, -1]\}$.

$$\left\{ \begin{aligned} \text{range}(L) &= \{L([x, y, z]) | [x, y, z] \in \mathbb{R}^3\} \\ &= \{[x - 2y, y + z] | x, y, z \in \mathbb{R}\} \\ &= \{x[1, 0] + y[-2, 1] + z[0, 1] | x, y, z \in \mathbb{R}\} \\ &= \text{span}\{[1, 0], [-2, 1], [0, 1]\} \end{aligned} \right.$$



Basis for $\ker(L)$ is $\{[2, 1, -1]\}$.

$$\left\{ \begin{aligned} \text{range}(L) &= \{L([x, y, z]) | [x, y, z] \in \mathbb{R}^3\} \\ &= \{[x - 2y, y + z] | x, y, z \in \mathbb{R}\} \\ &= \{x[1, 0] + y[-2, 1] + z[0, 1] | x, y, z \in \mathbb{R}\} \\ &= \text{span}\{[1, 0], [-2, 1], [0, 1]\} \\ &= \text{span}\{[1, 0], [0, 1]\} \end{aligned} \right.$$



Basis for $\ker(L)$ is $\{[2, 1, -1]\}$.

$$\left\{ \begin{aligned} \text{range}(L) &= \{L([x, y, z]) | [x, y, z] \in \mathbb{R}^3\} \\ &= \{[x - 2y, y + z] | x, y, z \in \mathbb{R}\} \\ &= \{x[1, 0] + y[-2, 1] + z[0, 1] | x, y, z \in \mathbb{R}\} \\ &= \text{span}\{[1, 0], [-2, 1], [0, 1]\} \\ &= \text{span}\{[1, 0], [0, 1]\} \end{aligned} \right.$$

Basis for $\text{range}(L)$ is $\{[1, 0], [0, 1]\}$.



Exercises

Q:. Suppose $L : P_3 \rightarrow P_2$ is a LT given by $L_1(at^3 + bt^2 + ct + d) = 3at^2 + 2bt + c$. Find $\ker(L)$ and $\text{range}(L)$.



Exercises

Q:. Suppose $L : P_3 \rightarrow P_2$ is a LT given by $L_1(at^3 + bt^2 + ct + d) = 3at^2 + 2bt + c$. Find $\ker(L)$ and $\text{range}(L)$.

Sol. $\ker(L) = \{0t^3 + 0t^2 + 0t + d \mid d \in \mathbb{R}\}$ and $\text{range}(L) = P_2$.



Exercises

Q.: Suppose $L : P_3 \rightarrow P_2$ is a LT given by $L_1(at^3 + bt^2 + ct + d) = 3at^2 + 2bt + c$. Find $\ker(L)$ and $\text{range}(L)$.

Sol. $\ker(L) = \{0t^3 + 0t^2 + 0t + d \mid d \in \mathbb{R}\}$ and $\text{range}(L) = P_2$.

Q.: Let $L : \mathbb{R}^4 \rightarrow P_2$ be a LT given by $L([a, b, c, d]) = a + (b + c)x + (b - d)x^2$.



Exercises

Q.: Suppose $L : P_3 \rightarrow P_2$ is a LT given by $L_1(at^3 + bt^2 + ct + d) = 3at^2 + 2bt + c$. Find $\ker(L)$ and $\text{range}(L)$.

Sol. $\ker(L) = \{0t^3 + 0t^2 + 0t + d | d \in \mathbb{R}\}$ and $\text{range}(L) = P_2$.

Q.: Let $L : \mathbb{R}^4 \rightarrow P_2$ be a LT given by $L([a, b, c, d]) = a + (b + c)x + (b - d)x^2$. Find $\ker(L)$ and $\text{range}(L)$.



Exercises

Q.: Suppose $L : P_3 \rightarrow P_2$ is a LT given by $L_1(at^3 + bt^2 + ct + d) = 3at^2 + 2bt + c$. Find $\ker(L)$ and $\text{range}(L)$.

Sol. $\ker(L) = \{0t^3 + 0t^2 + 0t + d \mid d \in \mathbb{R}\}$ and $\text{range}(L) = P_2$.

Q.: Let $L : \mathbb{R}^4 \rightarrow P_2$ be a LT given by $L([a, b, c, d]) = a + (b + c)x + (b - d)x^2$. Find $\ker(L)$ and $\text{range}(L)$. Also, find basis for $\ker(L)$ and $\text{range}(L)$.



Exercises

Q:. Suppose $L : P_3 \rightarrow P_2$ is a LT given by $L_1(at^3 + bt^2 + ct + d) = 3at^2 + 2bt + c$. Find $\ker(L)$ and $\text{range}(L)$.

Sol. $\ker(L) = \{0t^3 + 0t^2 + 0t + d | d \in \mathbb{R}\}$ and $\text{range}(L) = P_2$.

Q:. Let $L : \mathbb{R}^4 \rightarrow P_2$ be a LT given by $L([a, b, c, d]) = a + (b + c)x + (b - d)x^2$. Find $\ker(L)$ and $\text{range}(L)$. Also, find basis for $\ker(L)$ and $\text{range}(L)$.

Sol. $\ker(L) = \{[0, b, -b, b] | b \in \mathbb{R}\} = \text{span}\{[0, 1, -1, 1]\}$ and $\text{range}(L) = \text{span}\{1, x + x^2, x, x^2\} = \text{span}\{1, x, x^2\}$.



Example 4

Q:. Let $L : \mathbb{R}^3 \rightarrow \mathbb{R}^4$ be a LT given by $L([x, y, z]) = [x, y - z, x - y + z, x + y - z]$. Find a basis for $\ker(L)$ and $\text{range}(L)$.



Example 4

Q:. Let $L : \mathbb{R}^3 \rightarrow \mathbb{R}^4$ be a LT given by $L([x, y, z]) = [x, y - z, x - y + z, x + y - z]$. Find a basis for $\ker(L)$ and $\text{range}(L)$.

Sol. $\{[0, 1, 1]\}$ is a basis for $\ker(L)$ and $\{[1, 0, 1, 1], [0, 1, -1, 1]\}$ is a basis for $\text{range}(L)$.



Alternative approach for finding a basis for $\ker(L)$ (Kernel Method)



Alternative approach for finding a basis for $\ker(L)$ (Kernel Method)

Let $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a LT.



Alternative approach for finding a basis for $\ker(L)$ (Kernel Method)

Let $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a LT.

Step 1: Express $L(X) = AX$ for some $m \times n$ matrix A .



Alternative approach for finding a basis for $\ker(L)$ (Kernel Method)

Let $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a LT.

Step 1: Express $L(X) = AX$ for some $m \times n$ matrix A .

For Example 4, we have $L(X) = AX$ where



Alternative approach for finding a basis for $\ker(L)$ (Kernel Method)

Let $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a LT.

Step 1: Express $L(X) = AX$ for some $m \times n$ matrix A .

For Example 4, we have $L(X) = AX$ where

$$X = \begin{bmatrix} x \\ y \\ z \end{bmatrix} \text{ and } A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{bmatrix}$$



Step 2: Find matrix B , the RREF of A .



Step 2: Find matrix B, the RREF of A.

$$B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$



Step 2: Find matrix B , the RREF of A .

$$B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Step 3: Solve the system $BX = 0$ to compute $\ker(L)$ such that $\ker(L) = \text{span}(S)$ for some $S \subseteq \mathbb{R}^n$



Step 2: Find matrix B , the RREF of A .

$$B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Step 3: Solve the system $BX = 0$ to compute $\ker(L)$ such that $\ker(L) = \text{span}(S)$ for some $S \subseteq \mathbb{R}^n$ ($\ker(L) = \{X \in \mathbb{R}^n \mid L(X) = 0\}$).



Step 2: Find matrix B , the RREF of A .

$$B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Step 3: Solve the system $BX = 0$ to compute $\ker(L)$ such that $\ker(L) = \text{span}(S)$ for some $S \subseteq \mathbb{R}^n$ ($\ker(L) = \{X \in \mathbb{R}^n \mid L(X) = 0\}$).

$BX = 0$ gives

$$x = 0, y = z \implies \ker(L) = \{[0, z, z] \mid z \in \mathbb{R}\} = \text{span}\{[0, 1, 1]\}.$$



Step 4: Find a LI subset of \mathcal{S} which forms a basis for $\ker(L)$.



Step 4: Find a LI subset of \mathcal{S} which forms a basis for $\ker(L)$.

For Example 4, $\{[0, 1, 1]\}$ is a basis for $\ker(L)$.



Finding a basis for $\text{range}(L)$ (Range Method)

Step 1: Find matrix B , the RREF of A .



Finding a basis for $\text{range}(L)$ (Range Method)

Step 1: Find matrix B , the RREF of A .

$$B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$



Finding a basis for $\text{range}(L)$ (Range Method)

Step 1: Find matrix B , the RREF of A .

$$B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Step 2: Column vectors in A corresponding to **pivot columns** of $\text{RREF}(A)$ forms a basis for $\text{range}(L)$.



Finding a basis for range(L) (Range Method)

Step 1: Find matrix B , the RREF of A .

$$B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Step 2: Column vectors in A corresponding to **pivot columns** of $\text{RREF}(A)$ forms a basis for $\text{range}(L)$. Note that, Columns I and II have leading entry.



Finding a basis for range(L) (Range Method)

Step 1: Find matrix B , the RREF of A .

$$B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Step 2: Column vectors in A corresponding to **pivot columns** of $\text{RREF}(A)$ forms a basis for $\text{range}(L)$. Note that, Columns I and II have leading entry. Thus, the corresponding column vector of A , i.e., $\{[1, 0, 1, 1], [0, 1, -1, 1]\}$ is a basis of $\text{range}(L)$.



Example 5

Q:. Let $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be a LT given by

$$L\left(\begin{bmatrix} x \\ y \\ z \end{bmatrix}\right) = \begin{bmatrix} 5 & 1 & -1 \\ -3 & 0 & 1 \\ 1 & -1 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$



Example 5

Q:. Let $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be a LT given by

$$L\left(\begin{bmatrix} x \\ y \\ z \end{bmatrix}\right) = \begin{bmatrix} 5 & 1 & -1 \\ -3 & 0 & 1 \\ 1 & -1 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

- Is $[1, -2, 3]^T \in \ker(L)$.



Example 5

Q:. Let $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be a LT given by

$$L\left(\begin{bmatrix} x \\ y \\ z \end{bmatrix}\right) = \begin{bmatrix} 5 & 1 & -1 \\ -3 & 0 & 1 \\ 1 & -1 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

- Is $[1, -2, 3]^T \in \ker(L)$.
- Is $[2, -1, 4]^T \in \text{range}(L)$.



Example 5

Q:. Let $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be a LT given by

$$L\left(\begin{bmatrix} x \\ y \\ z \end{bmatrix}\right) = \begin{bmatrix} 5 & 1 & -1 \\ -3 & 0 & 1 \\ 1 & -1 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

- Is $[1, -2, 3]^T \in \ker(L)$.
- Is $[2, -1, 4]^T \in \text{range}(L)$.

Sol. Yes, $[1, -2, 3]^T \in \ker(L)$ because
 $L([1, -2, 3]^T) = [0, 0, 0]^T$.



No, $[2, -1, 4]^T \notin \text{range}(L)$.



No, $[2, -1, 4]^T \notin \text{range}(L)$.

If $v = [2, -1, 4]^T \in \text{range}(L)$, then there would exist a vector $X = [x, y, z]^T$ such that $AX = v$.



No, $[2, -1, 4]^T \notin \text{range}(L)$.

If $v = [2, -1, 4]^T \in \text{range}(L)$, then there would exist a vector $X = [x, y, z]^T$ such that $AX = v$. This is equivalent to the system



No, $[2, -1, 4]^T \notin \text{range}(L)$.

If $v = [2, -1, 4]^T \in \text{range}(L)$, then there would exist a vector $X = [x, y, z]^T$ such that $AX = v$. This is equivalent to the system

$$\begin{cases} 5x + y - z = 2 \\ -3x + z = -1 \\ x - y - z = 4 \end{cases}$$



No, $[2, -1, 4]^T \notin \text{range}(L)$.

If $v = [2, -1, 4]^T \in \text{range}(L)$, then there would exist a vector $X = [x, y, z]^T$ such that $AX = v$. This is equivalent to the system

$$\begin{cases} 5x + y - z = 2 \\ -3x + z = -1 \\ x - y - z = 4 \end{cases} \quad \text{which has no solution.}$$



The Dimension Theorem



The Dimension Theorem

If $L : V \rightarrow W$ is a LT and V is finite dimensional, then $\text{range}(L)$ is finite dimensional, and

$$\dim(\ker(L)) + \dim(\text{range}(L)) = \dim(V)$$



The Dimension Theorem

If $L : V \rightarrow W$ is a LT and V is finite dimensional, then $\text{range}(L)$ is finite dimensional, and

$$\dim(\ker(L)) + \dim(\text{range}(L)) = \dim(V)$$

Note: $\dim(\ker(L))$ is called nullity(L) and $\dim(\text{range}(L))$ is called rank(L).



Example 6



Example 6

Q:. Consider a LT $L : P_2 \rightarrow P_3$ given by $L(a + bx + cx^2) = x(a + bx + cx^2)$.



Example 6

Q:. Consider a LT $L : P_2 \rightarrow P_3$ given by $L(a + bx + cx^2) = x(a + bx + cx^2)$. Find $\dim(\ker(L))$ and $\dim(\text{range}(L))$ independently and hence verify the dimension theorem.



Example 6

Q.: Consider a LT $L : P_2 \rightarrow P_3$ given by $L(a + bx + cx^2) = x(a + bx + cx^2)$. Find $\dim(\ker(L))$ and $\dim(\text{range}(L))$ independently and hence verify the dimension theorem.

Sol.

$$\begin{aligned}\ker(L) &= \{(a + bx + cx^2) \in P_2 \mid L(a + bx + cx^2) = 0_{P_3}\} \\ &= \{(a + bx + cx^2) \in P_2 \mid ax + bx^2 + cx^3 = 0\} \\ &= \{(a + bx + cx^2) \in P_2 \mid a = b = c = 0\} \\ &= \{0_{P_2}\}\end{aligned}$$



$$\implies \dim(\ker(L)) = 0.$$



$$\implies \dim(\ker(L)) = 0.$$

Also $\dim(\text{range}(L)) = 3$ (why).



$$\implies \dim(\ker(L)) = 0.$$

Also $\dim(\text{range}(L)) = 3$ (why). Since, $\dim(P_2) = 3$,



$$\implies \dim(\ker(L)) = 0.$$

Also $\dim(\text{range}(L)) = 3$ (why). Since, $\dim(P_2) = 3$, clearly, we have



$$\implies \dim(\ker(L)) = 0.$$

Also $\dim(\text{range}(L)) = 3$ (why). Since, $\dim(P_2) = 3$, clearly, we have $\dim(\ker(L)) + \dim(\text{range}(L)) = \dim(P_2)$.



Example 7



Example 7

Q:. Consider a LT $L : M_{33} \rightarrow \mathbb{R}$ given by $L(A) = \text{trace}(A)$.



Example 7

Q:. Consider a LT $L : M_{33} \rightarrow \mathbb{R}$ given by $L(A) = \text{trace}(A)$. Find $\ker(L)$, $\dim(\ker(L))$, $\text{range}(L)$ and $\dim(\text{range}(L))$.



Example 7

Q.: Consider a LT $L : M_{33} \rightarrow \mathbb{R}$ given by $L(A) = \text{trace}(A)$. Find $\ker(L)$, $\dim(\ker(L))$, $\text{range}(L)$ and $\dim(\text{range}(L))$.

Sol. $\ker(L) =$

$$\left\{ \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & -a-e \end{bmatrix} \in M_{33} \mid a, b, c, d, e, f, g, h \in \mathbb{R} \right\} \Rightarrow$$



Example 7

Q:. Consider a LT $L : M_{33} \rightarrow \mathbb{R}$ given by $L(A) = \text{trace}(A)$. Find $\ker(L)$, $\dim(\ker(L))$, $\text{range}(L)$ and $\dim(\text{range}(L))$.

Sol. $\ker(L) =$

$$\left\{ \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & -a-e \end{bmatrix} \in M_{33} \mid a, b, c, d, e, f, g, h \in \mathbb{R} \right\} \Rightarrow$$

$$\dim(\ker(L)) = 8.$$



Example 7

Q:. Consider a LT $L : M_{33} \rightarrow \mathbb{R}$ given by $L(A) = \text{trace}(A)$. Find $\ker(L)$, $\dim(\ker(L))$, $\text{range}(L)$ and $\dim(\text{range}(L))$.

Sol. $\ker(L) =$

$$\left\{ \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & -a-e \end{bmatrix} \in M_{33} \mid a, b, c, d, e, f, g, h \in \mathbb{R} \right\} \Rightarrow$$

$\dim(\ker(L)) = 8$. Also, $\text{range}(L) = \mathbb{R}$



Example 7

Q.: Consider a LT $L : M_{33} \rightarrow \mathbb{R}$ given by $L(A) = \text{trace}(A)$. Find $\ker(L)$, $\dim(\ker(L))$, $\text{range}(L)$ and $\dim(\text{range}(L))$.

Sol. $\ker(L) =$

$$\left\{ \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & -a-e \end{bmatrix} \in M_{33} \mid a, b, c, d, e, f, g, h \in \mathbb{R} \right\} \Rightarrow$$

$\dim(\ker(L)) = 8$. Also, $\text{range}(L) = \mathbb{R}$ and $\dim(\text{range}(L)) = 1$.



Example 7

Q.: Consider a LT $L : M_{33} \rightarrow \mathbb{R}$ given by $L(A) = \text{trace}(A)$. Find $\ker(L)$, $\dim(\ker(L))$, $\text{range}(L)$ and $\dim(\text{range}(L))$.

Sol. $\ker(L) =$

$$\left\{ \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & -a-e \end{bmatrix} \in M_{33} \mid a, b, c, d, e, f, g, h \in \mathbb{R} \right\} \Rightarrow$$

$\dim(\ker(L)) = 8$. Also, $\text{range}(L) = \mathbb{R}$ and $\dim(\text{range}(L)) = 1$. Hence,



Example 7

Q.: Consider a LT $L : M_{33} \rightarrow \mathbb{R}$ given by $L(A) = \text{trace}(A)$. Find $\ker(L)$, $\dim(\ker(L))$, $\text{range}(L)$ and $\dim(\text{range}(L))$.

Sol. $\ker(L) =$

$$\left\{ \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & -a-e \end{bmatrix} \in M_{33} \mid a, b, c, d, e, f, g, h \in \mathbb{R} \right\} \Rightarrow$$

$\dim(\ker(L)) = 8$. Also, $\text{range}(L) = \mathbb{R}$ and $\dim(\text{range}(L)) = 1$. Hence,

$$\dim(\ker(L)) + \dim(\text{range}(L)) = \dim(M_{33})$$



Example 8



Example 8

Q:. Consider a LT $L : M_{33} \rightarrow M_{33}$ given by $L(A) = A + A^T$.



Example 8

Q:. Consider a LT $L : M_{33} \rightarrow M_{33}$ given by $L(A) = A + A^T$. Find $\ker(L)$, $\text{range}(L)$ and verify dimension theorem.



Example 8

Q:. Consider a LT $L : M_{33} \rightarrow M_{33}$ given by $L(A) = A + A^T$. Find $\ker(L)$, $\text{range}(L)$ and verify dimension theorem.

Sol. $\ker(L)$ is precisely the set of all skew-symmetric 3×3 matrices.



Example 8

Q:. Consider a LT $L : M_{33} \rightarrow M_{33}$ given by $L(A) = A + A^T$. Find $\ker(L)$, $\text{range}(L)$ and verify dimension theorem.

Sol. $\ker(L)$ is precisely the set of all skew-symmetric 3×3 matrices.

$\text{range}(L)$ is the set of all symmetric 3×3 matrices.



Example 8

Q.: Consider a LT $L : M_{33} \rightarrow M_{33}$ given by $L(A) = A + A^T$. Find $\ker(L)$, $\text{range}(L)$ and verify dimension theorem.

Sol. $\ker(L)$ is precisely the set of all skew-symmetric 3×3 matrices.

$\text{range}(L)$ is the set of all symmetric 3×3 matrices.

$$\dim(\ker(L)) = 3$$



Example 8

Q.: Consider a LT $L : M_{33} \rightarrow M_{33}$ given by $L(A) = A + A^T$. Find $\ker(L)$, $\text{range}(L)$ and verify dimension theorem.

Sol. $\ker(L)$ is precisely the set of all skew-symmetric 3×3 matrices.

$\text{range}(L)$ is the set of all symmetric 3×3 matrices.

$\dim(\ker(L)) = 3$ and $\dim(\text{range}(L)) = 6$.



True or False



True or False

Q:. If $L : V \rightarrow W$ is a LT and $\dim(V) = 5, \dim(W) = 3$, then the Dimension Theorem implies that $\dim(\ker(L)) = 2$.



True or False

Q:. If $L : V \rightarrow W$ is a LT and $\dim(V) = 5, \dim(W) = 3$, then the Dimension Theorem implies that $\dim(\ker(L)) = 2$.

Sol. False



True or False

Q:. If $L : V \rightarrow W$ is a LT and $\dim(V) = 5, \dim(W) = 3$, then the Dimension Theorem implies that $\dim(\ker(L)) = 2$.

Sol. False Counterexample:



True or False

Q:. If $L : V \rightarrow W$ is a LT and $\dim(V) = 5, \dim(W) = 3$, then the Dimension Theorem implies that $\dim(\ker(L)) = 2$.

Sol. False Counterexample: Let $L : \mathbb{R}^5 \rightarrow \mathbb{R}^3$ given by $L(v) = 0_{\mathbb{R}^3}$ for all $v \in \mathbb{R}^5$.



True or False

Q:. If $L : V \rightarrow W$ is a LT and $\dim(V) = 5, \dim(W) = 3$, then the Dimension Theorem implies that $\dim(\ker(L)) = 2$.

Sol. False Counterexample: Let $L : \mathbb{R}^5 \rightarrow \mathbb{R}^3$ given by $L(v) = 0_{\mathbb{R}^3}$ for all $v \in \mathbb{R}^5$. $\ker(L) = \mathbb{R}^5 \implies$



True or False

Q:. If $L : V \rightarrow W$ is a LT and $\dim(V) = 5, \dim(W) = 3$, then the Dimension Theorem implies that $\dim(\ker(L)) = 2$.

Sol. False Counterexample: Let $L : \mathbb{R}^5 \rightarrow \mathbb{R}^3$ given by $L(v) = 0_{\mathbb{R}^3}$ for all $v \in \mathbb{R}^5$. $\ker(L) = \mathbb{R}^5 \implies \dim(\ker(L)) = 5 \neq 2$.



True or False

Q:. If $L : V \rightarrow W$ is a LT and $\dim(V) = 5, \dim(W) = 3$, then the Dimension Theorem implies that $\dim(\ker(L)) = 2$.

Sol. False Counterexample: Let $L : \mathbb{R}^5 \rightarrow \mathbb{R}^3$ given by $L(v) = 0_{\mathbb{R}^3}$ for all $v \in \mathbb{R}^5$. $\ker(L) = \mathbb{R}^5 \implies \dim(\ker(L)) = 5 \neq 2$.

Q:. If $L : V \rightarrow W$ is a LT, then $\text{range}(L)$ is a subspace of V .



True or False

Q:. If $L : V \rightarrow W$ is a LT and $\dim(V) = 5, \dim(W) = 3$, then the Dimension Theorem implies that $\dim(\ker(L)) = 2$.

Sol. False Counterexample: Let $L : \mathbb{R}^5 \rightarrow \mathbb{R}^3$ given by $L(v) = 0_{\mathbb{R}^3}$ for all $v \in \mathbb{R}^5$. $\ker(L) = \mathbb{R}^5 \implies \dim(\ker(L)) = 5 \neq 2$.

Q:. If $L : V \rightarrow W$ is a LT, then $\text{range}(L)$ is a subspace of V .

Sol. False,



True or False

Q:. If $L : V \rightarrow W$ is a LT and $\dim(V) = 5, \dim(W) = 3$, then the Dimension Theorem implies that $\dim(\ker(L)) = 2$.

Sol. False Counterexample: Let $L : \mathbb{R}^5 \rightarrow \mathbb{R}^3$ given by $L(v) = 0_{\mathbb{R}^3}$ for all $v \in \mathbb{R}^5$. $\ker(L) = \mathbb{R}^5 \implies \dim(\ker(L)) = 5 \neq 2$.

Q:. If $L : V \rightarrow W$ is a LT, then $\text{range}(L)$ is a subspace of V .

Sol. False, $\text{range}(L)$ is a subspace of W .



Exercises

Q:. Let W be the vector space of all 2×2 symmetric matrices. Define a LT $L : W \rightarrow P_2$ by

$$L\left(\begin{bmatrix} a & b \\ b & c \end{bmatrix}\right) = (a - b) + (b - c)x + (c - a)x^2$$

Find $\dim(\ker(L))$ and $\dim(\text{range}(L))$.



Exercises

Q:. Let W be the vector space of all 2×2 symmetric matrices. Define a LT $L : W \rightarrow P_2$ by

$$L\left(\begin{bmatrix} a & b \\ b & c \end{bmatrix}\right) = (a - b) + (b - c)x + (c - a)x^2$$

Find $\dim(\ker(L))$ and $\dim(\text{range}(L))$.

Sol. $\dim(\ker(L)) = 1$ and $\dim(\text{range}(L)) = 2$.



Q:. Let $\{e_1, e_2, e_3, e_4\}$ be standard basis for \mathbb{R}^4 and $L : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ be a LT given by



Q:. Let $\{e_1, e_2, e_3, e_4\}$ be standard basis for \mathbb{R}^4 and $L : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ be a LT given by $T(e_1) = [1, 1, 1], T(e_2) = [1, -1, 1], T(e_3) = [1, 0, 0], T(e_4) = [1, 0, 1]$.



Q:. Let $\{e_1, e_2, e_3, e_4\}$ be standard basis for \mathbb{R}^4 and $L : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ be a LT given by $T(e_1) = [1, 1, 1], T(e_2) = [1, -1, 1], T(e_3) = [1, 0, 0], T(e_4) = [1, 0, 1]$. Find $\ker(L)$, $\dim(\ker(L))$, $\text{range}(L)$ and $\dim(\text{range}(L))$.



Q:. Let $\{e_1, e_2, e_3, e_4\}$ be standard basis for \mathbb{R}^4 and $L : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ be a LT given by $T(e_1) = [1, 1, 1], T(e_2) = [1, -1, 1], T(e_3) = [1, 0, 0], T(e_4) = [1, 0, 1]$. Find $\ker(L)$, $\dim(\ker(L))$, $\text{range}(L)$ and $\dim(\text{range}(L))$.

Sol. $\dim(\ker(L)) = 1$ and $\dim(\text{range}(L)) = 3$.



Q:. Let $\{e_1, e_2, e_3, e_4\}$ be standard basis for \mathbb{R}^4 and $L : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ be a LT given by $T(e_1) = [1, 1, 1], T(e_2) = [1, -1, 1], T(e_3) = [1, 0, 0], T(e_4) = [1, 0, 1]$. Find $\ker(L)$, $\dim(\ker(L))$, $\text{range}(L)$ and $\dim(\text{range}(L))$.

Sol. $\dim(\ker(L)) = 1$ and $\dim(\text{range}(L)) = 3$.

Q:. Consider $L : P_2 \rightarrow P_4$ given by $L(p) = x^2 p$. Find $\ker(L)$, $\dim(\ker(L))$, $\text{range}(L)$ and $\dim(\text{range}(L))$.



Q:. Let $\{e_1, e_2, e_3, e_4\}$ be standard basis for \mathbb{R}^4 and $L : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ be a LT given by $T(e_1) = [1, 1, 1], T(e_2) = [1, -1, 1], T(e_3) = [1, 0, 0], T(e_4) = [1, 0, 1]$. Find $\ker(L)$, $\dim(\ker(L))$, $\text{range}(L)$ and $\dim(\text{range}(L))$.

Sol. $\dim(\ker(L)) = 1$ and $\dim(\text{range}(L)) = 3$.

Q:. Consider $L : P_2 \rightarrow P_4$ given by $L(p) = x^2 p$. Find $\ker(L)$, $\dim(\ker(L))$, $\text{range}(L)$ and $\dim(\text{range}(L))$.

Sol. $\dim(\ker(L)) = 0$ and $\dim(\text{range}(L)) = 3$.

