

MATHEMATICS-II (MATH F112)

Dr. Krishnendra Shekhawat

BITS PILANI
Department of Mathematics



CHAPTER 5

Linear Transformations



Topics to be covered in Chapter 5



Topics to be covered in Chapter 5

- Introduction to Linear Transformations



Topics to be covered in Chapter 5

- Introduction to Linear Transformations
- The Matrix of a Linear Transformation



Topics to be covered in Chapter 5

- Introduction to Linear Transformations
- The Matrix of a Linear Transformation
- The Dimension Theorem



Topics to be covered in Chapter 5

- Introduction to Linear Transformations
- The Matrix of a Linear Transformation
- The Dimension Theorem
- One-to-One and Onto Linear Transformations



Topics to be covered in Chapter 5

- Introduction to Linear Transformations
- The Matrix of a Linear Transformation
- The Dimension Theorem
- One-to-One and Onto Linear Transformations
- Isomorphism



Section 5.1

Introduction to Linear Transformations



Linear transformation (LT)



Linear transformation (LT)

Let U and V be real vector spaces.



Linear transformation (LT)

Let U and V be real vector spaces. A map $L : U \rightarrow V$ is called a **Linear map**, or **Linear transformation (LT)**



Linear transformation (LT)

Let U and V be real vector spaces. A map $L : U \rightarrow V$ is called a **Linear map**, or **Linear transformation (LT)** if and only if both of the following are true:



Linear transformation (LT)

Let U and V be real vector spaces. A map $L : U \rightarrow V$ is called a **Linear map**, or **Linear transformation (LT)** if and only if both of the following are true:

- $L(u + v) = L(u) + L(v)$ for all $u, v \in U$



Linear transformation (LT)

Let U and V be real vector spaces. A map $L : U \rightarrow V$ is called a **Linear map**, or **Linear transformation (LT)** if and only if both of the following are true:

- $L(u + v) = L(u) + L(v)$ for all $u, v \in U$ (L preserves vector addition)



Linear transformation (LT)

Let U and V be real vector spaces. A map $L : U \rightarrow V$ is called a **Linear map**, or **Linear transformation (LT)** if and only if both of the following are true:

- $L(u + v) = L(u) + L(v)$ for all $u, v \in U$ (L preserves vector addition)
- $L(cu) = cL(u)$ for all $c \in \mathbb{R}$ and all $u \in U$



Linear transformation (LT)

Let U and V be real vector spaces. A map $L : U \rightarrow V$ is called a **Linear map**, or **Linear transformation (LT)** if and only if both of the following are true:

- $L(u + v) = L(u) + L(v)$ for all $u, v \in U$ (L preserves vector addition)
- $L(cu) = cL(u)$ for all $c \in \mathbb{R}$ and all $u \in U$ (L preserves scalar multiplication)



Example 1

Q:. Consider the mapping $L : M_{mn} \rightarrow M_{nm}$, given by $L(A) = A^T$ for any $m \times n$ matrix A .



Example 1

Q:. Consider the mapping $L : M_{mn} \rightarrow M_{nm}$, given by $L(A) = A^T$ for any $m \times n$ matrix A . Check whether L is a LT.



Example 1

Q:. Consider the mapping $L : M_{mn} \rightarrow M_{nm}$, given by $L(A) = A^T$ for any $m \times n$ matrix A . Check whether L is a LT.

Sol. Let $A, B \in M_{mn}$ and $c \in \mathbb{R}$.



Example 1

Q:. Consider the mapping $L : M_{mn} \rightarrow M_{nm}$, given by $L(A) = A^T$ for any $m \times n$ matrix A . Check whether L is a LT.

Sol. Let $A, B \in M_{mn}$ and $c \in \mathbb{R}$.

- $$L(A + B) = (A + B)^T = A^T + B^T = L(A) + L(B)$$



Example 1

Q:. Consider the mapping $L : M_{mn} \rightarrow M_{nm}$, given by $L(A) = A^T$ for any $m \times n$ matrix A . Check whether L is a LT.

Sol. Let $A, B \in M_{mn}$ and $c \in \mathbb{R}$.

- $L(A + B) = (A + B)^T = A^T + B^T = L(A) + L(B)$
- $L(cA) = (cA)^T = cA^T = cL(A)$



Example 1

Q.: Consider the mapping $L : M_{mn} \rightarrow M_{nm}$, given by $L(A) = A^T$ for any $m \times n$ matrix A . Check whether L is a LT.

Sol. Let $A, B \in M_{mn}$ and $c \in \mathbb{R}$.

- $L(A + B) = (A + B)^T = A^T + B^T = L(A) + L(B)$
- $L(cA) = (cA)^T = cA^T = cL(A)$

Hence, L is a LT.



Example 2

Q:. Consider the mapping $L : \mathbb{R}^2 \rightarrow \mathbb{R}^3$, given by $L([x, y]) = ([x, y, xy])$.



Example 2

Q:. Consider the mapping $L : \mathbb{R}^2 \rightarrow \mathbb{R}^3$, given by $L([x, y]) = ([x, y, xy])$. Check whether L is a LT.



Example 2

Q:. Consider the mapping $L : \mathbb{R}^2 \rightarrow \mathbb{R}^3$, given by $L([x,y]) = ([x,y,xy])$. Check whether L is a LT.

Sol. Note that



Example 2

Q:. Consider the mapping $L : \mathbb{R}^2 \rightarrow \mathbb{R}^3$, given by $L([x,y]) = ([x,y,xy])$. Check whether L is a LT.

Sol. Note that

$$L(c([x,y])) =$$



Example 2

Q:. Consider the mapping $L : \mathbb{R}^2 \rightarrow \mathbb{R}^3$, given by $L([x,y]) = ([x,y,xy])$. Check whether L is a LT.

Sol. Note that

$$L(c([x,y])) = L([cx,cy]) = [cx,cy,c^2xy] = c[x,y,cxy]$$



Example 2

Q:. Consider the mapping $L : \mathbb{R}^2 \rightarrow \mathbb{R}^3$, given by $L([x,y]) = ([x,y,xy])$. Check whether L is a LT.

Sol. Note that

$$L(c([x,y])) = L([cx,cy]) = [cx,cy,c^2xy] = c[x,y,cxy] \neq c([x,y,xy]),$$



Example 2

Q:. Consider the mapping $L : \mathbb{R}^2 \rightarrow \mathbb{R}^3$, given by $L([x,y]) = ([x,y,xy])$. Check whether L is a LT.

Sol. Note that

$$L(c([x,y])) = L([cx,cy]) = [cx,cy,c^2xy] = c[x,y,cxy] \\ \neq c([x,y,xy]), \text{ i.e., } L(c([x,y])) \neq cL([x,y]) \text{ for some } c \in \mathbb{R}.$$



Example 2

Q:. Consider the mapping $L : \mathbb{R}^2 \rightarrow \mathbb{R}^3$, given by $L([x, y]) = ([x, y, xy])$. Check whether L is a LT.

Sol. Note that

$$L(c([x, y])) = L([cx, cy]) = [cx, cy, c^2xy] = c[x, y, cxy] \\ \neq c([x, y, xy]), \text{ i.e., } L(c([x, y])) \neq cL([x, y]) \text{ for some } c \in \mathbb{R}.$$

Hence, L is not a LT.



Exercises

Q: Check which of the following maps are LT.



Exercises

Q.: Check which of the following maps are LT.

- $L : P_2 \rightarrow \mathbb{R}^3$, given by $L(a + bx + cx^2) = (a, b, c)$.



Exercises

Q:. Check which of the following maps are LT.

- $L : P_2 \rightarrow \mathbb{R}^3$, given by $L(a + bx + cx^2) = (a, b, c)$.
- $L : \mathbb{R}^3 \rightarrow \mathbb{R}^2$, given by $L([x, y, z]) = ([x - y, y + z])$.



Exercises

Q:. Check which of the following maps are LT.

- $L : P_2 \rightarrow \mathbb{R}^3$, given by $L(a + bx + cx^2) = (a, b, c)$.
- $L : \mathbb{R}^3 \rightarrow \mathbb{R}^2$, given by $L([x, y, z]) = ([x - y, y + z])$.
- $L : \mathbb{R}^2 \rightarrow \mathbb{R}^2$, given by $L([a, b]) = [a, -b]$.



Linear Operator



Linear Operator

Let V be a vector space.



Linear Operator

Let V be a vector space. A **linear operator** on V is a LT whose domain and codomain are both V .



Linear Operator

Let V be a vector space. A **linear operator** on V is a LT whose domain and codomain are both V .

For example, the mapping $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$, given by $L([x, y, z]) = [x, y, -z]$ is a linear operator.



Theorem: Let V and W be vector spaces, and let $L : V \rightarrow W$ be a LT.



Theorem: Let V and W be vector spaces, and let $L : V \rightarrow W$ be a LT. Let $\mathbf{0}_V$ be the zero vector in V and $\mathbf{0}_W$ be the zero vector in W . Then



Theorem: Let V and W be vector spaces, and let $L : V \rightarrow W$ be a LT. Let $\mathbf{0}_V$ be the zero vector in V and $\mathbf{0}_W$ be the zero vector in W . Then

$$\textcircled{1} \quad L(\mathbf{0}_V) = \mathbf{0}_W$$



Theorem: Let V and W be vector spaces, and let $L : V \rightarrow W$ be a LT. Let $\mathbf{0}_V$ be the zero vector in V and $\mathbf{0}_W$ be the zero vector in W . Then

- ❶ $L(\mathbf{0}_V) = \mathbf{0}_W$
- ❷ $L(-\mathbf{v}) = -L(\mathbf{v})$, for all $\mathbf{v} \in V$



Theorem: Let V and W be vector spaces, and let $L : V \rightarrow W$ be a LT. Let $\mathbf{0}_V$ be the zero vector in V and $\mathbf{0}_W$ be the zero vector in W . Then

- ❶ $L(\mathbf{0}_V) = \mathbf{0}_W$
- ❷ $L(-\mathbf{v}) = -L(\mathbf{v})$, for all $\mathbf{v} \in V$
- ❸ For $n \geq 2$,



Theorem: Let V and W be vector spaces, and let $L : V \rightarrow W$ be a LT. Let $\mathbf{0}_V$ be the zero vector in V and $\mathbf{0}_W$ be the zero vector in W . Then

① $L(\mathbf{0}_V) = \mathbf{0}_W$

② $L(-\mathbf{v}) = -L(\mathbf{v})$, for all $\mathbf{v} \in V$

③ For $n \geq 2$,

$$\begin{aligned} L(a_1\mathbf{v}_1 + a_2\mathbf{v}_2 + \cdots + a_n\mathbf{v}_n) \\ = a_1L(\mathbf{v}_1) + a_2L(\mathbf{v}_2) + \cdots + a_nL(\mathbf{v}_n), \end{aligned}$$



Theorem: Let V and W be vector spaces, and let $L : V \rightarrow W$ be a LT. Let $\mathbf{0}_V$ be the zero vector in V and $\mathbf{0}_W$ be the zero vector in W . Then

① $L(\mathbf{0}_V) = \mathbf{0}_W$

② $L(-\mathbf{v}) = -L(\mathbf{v})$, for all $\mathbf{v} \in V$

③ For $n \geq 2$,

$$\begin{aligned} L(a_1\mathbf{v}_1 + a_2\mathbf{v}_2 + \cdots + a_n\mathbf{v}_n) \\ = a_1L(\mathbf{v}_1) + a_2L(\mathbf{v}_2) + \cdots + a_nL(\mathbf{v}_n), \end{aligned}$$

for all $a_1, a_2, \dots, a_n \in \mathbb{R}$ and $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n \in V$.



Example 3

Q.: Consider the mapping $L : M_{22} \rightarrow \mathbb{R}$ given by

$$L \begin{bmatrix} a & b \\ c & d \end{bmatrix} = a + b + c + d - 1.$$


Example 3

Q:. Consider the mapping $L : M_{22} \rightarrow \mathbb{R}$ given by

$$L \begin{bmatrix} a & b \\ c & d \end{bmatrix} = a + b + c + d - 1. \text{ Check whether } L \text{ is a LT.}$$


Example 3

Q.: Consider the mapping $L : M_{22} \rightarrow \mathbb{R}$ given by
$$L \begin{bmatrix} a & b \\ c & d \end{bmatrix} = a + b + c + d - 1.$$
 Check whether L is a LT.

Sol.
$$L \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = 0 + 0 + 0 + 0 - 1 = -1 \neq 0.$$



Example 3

Q.: Consider the mapping $L : M_{22} \rightarrow \mathbb{R}$ given by
$$L \begin{bmatrix} a & b \\ c & d \end{bmatrix} = a + b + c + d - 1.$$
 Check whether L is a LT.

Sol.
$$L \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = 0 + 0 + 0 + 0 - 1 = -1 \neq 0.$$

Since $L(0_{M_{22}}) \neq 0_{\mathbb{R}}$,



Example 3

Q.: Consider the mapping $L : M_{22} \rightarrow \mathbb{R}$ given by
$$L \begin{bmatrix} a & b \\ c & d \end{bmatrix} = a + b + c + d - 1.$$
 Check whether L is a LT.

Sol.
$$L \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = 0 + 0 + 0 + 0 - 1 = -1 \neq 0.$$

Since $L(0_{M_{22}}) \neq 0_{\mathbb{R}}$, L is not a LT.



Example 4

Q:. Suppose $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is a linear operator and $L([1, 0, 0]) = [-2, 1, 0]$, $L([0, 1, 0]) = [3, -2, 1]$, and $L([0, 0, 1]) = [0, -1, 3]$. Find $L([-3, 2, 4])$.



Example 4

Q:. Suppose $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is a linear operator and $L([1, 0, 0]) = [-2, 1, 0]$, $L([0, 1, 0]) = [3, -2, 1]$, and $L([0, 0, 1]) = [0, -1, 3]$. Find $L([-3, 2, 4])$. Give a formula for $L([x, y, z])$ for $[x, y, z] \in \mathbb{R}^3$.



Example 4

Q:. Suppose $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is a linear operator and $L([1, 0, 0]) = [-2, 1, 0]$, $L([0, 1, 0]) = [3, -2, 1]$, and $L([0, 0, 1]) = [0, -1, 3]$. Find $L([-3, 2, 4])$. Give a formula for $L([x, y, z])$ for $[x, y, z] \in \mathbb{R}^3$.

Sol.

$$\left\{ \begin{aligned} L([-3, 2, 4]) &= L(-3[1, 0, 0] + 2[0, 1, 0] + 4[0, 0, 1]) \end{aligned} \right.$$



Example 4

Q:. Suppose $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is a linear operator and $L([1, 0, 0]) = [-2, 1, 0]$, $L([0, 1, 0]) = [3, -2, 1]$, and $L([0, 0, 1]) = [0, -1, 3]$. Find $L([-3, 2, 4])$. Give a formula for $L([x, y, z])$ for $[x, y, z] \in \mathbb{R}^3$.

Sol.

$$\left\{ \begin{aligned} L([-3, 2, 4]) &= L(-3[1, 0, 0] + 2[0, 1, 0] + 4[0, 0, 1]) \\ &= -3L([1, 0, 0]) + 2L([0, 1, 0]) + 4L([0, 0, 1]) \end{aligned} \right.$$



Example 4

Q:. Suppose $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is a linear operator and $L([1, 0, 0]) = [-2, 1, 0]$, $L([0, 1, 0]) = [3, -2, 1]$, and $L([0, 0, 1]) = [0, -1, 3]$. Find $L([-3, 2, 4])$. Give a formula for $L([x, y, z])$ for $[x, y, z] \in \mathbb{R}^3$.

Sol.

$$\left\{ \begin{aligned} L([-3, 2, 4]) &= L(-3[1, 0, 0] + 2[0, 1, 0] + 4[0, 0, 1]) \\ &= -3L([1, 0, 0]) + 2L([0, 1, 0]) + 4L([0, 0, 1]) \\ &= -3[-2, 1, 0] + 2[3, -2, 1] + 4[0, -1, 3] \end{aligned} \right.$$



Example 4

Q:. Suppose $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is a linear operator and $L([1, 0, 0]) = [-2, 1, 0]$, $L([0, 1, 0]) = [3, -2, 1]$, and $L([0, 0, 1]) = [0, -1, 3]$. Find $L([-3, 2, 4])$. Give a formula for $L([x, y, z])$ for $[x, y, z] \in \mathbb{R}^3$.

Sol.

$$\left\{ \begin{aligned} L([-3, 2, 4]) &= L(-3[1, 0, 0] + 2[0, 1, 0] + 4[0, 0, 1]) \\ &= -3L([1, 0, 0]) + 2L([0, 1, 0]) + 4L([0, 0, 1]) \\ &= -3[-2, 1, 0] + 2[3, -2, 1] + 4[0, -1, 3] \\ &= [12, -11, 14] \end{aligned} \right.$$



Similarly, $L([x, y, z]) = L(x[1, 0, 0] + y[0, 1, 0] + z[0, 0, 1])$



Similarly, $L([x, y, z]) = L(x[1, 0, 0] + y[0, 1, 0] + z[0, 0, 1])$

$$L([x, y, z]) = x[-2, 1, 0] + y[3, -2, 1] + z[0, -1, 3]$$



Similarly, $L([x, y, z]) = L(x[1, 0, 0] + y[0, 1, 0] + z[0, 0, 1])$

$$L([x, y, z]) = x[-2, 1, 0] + y[3, -2, 1] + z[0, -1, 3]$$

$$L([x, y, z]) = [-2x + 3y, x - 2y - z, y + 3z]$$



Similarly, $L([x, y, z]) = L(x[1, 0, 0] + y[0, 1, 0] + z[0, 0, 1])$

$$L([x, y, z]) = x[-2, 1, 0] + y[3, -2, 1] + z[0, -1, 3]$$

$$L([x, y, z]) = [-2x + 3y, x - 2y - z, y + 3z]$$

Note that

$$L \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -2 & 3 & 0 \\ 1 & -2 & -1 \\ 0 & 1 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$



Exercise

Q:. Suppose $L : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a linear operator and $L([1, 1]) = [3, 0]$ and $L([-1, 1]) = [0, 1]$. Compute $L([x, y])$.

Sol. $L([x, y]) = \left[\frac{3x+3y}{2}, \frac{-x+y}{2} \right]$



Exercise

Q:. Suppose $L : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a linear operator and $L([1, 1]) = [3, 0]$ and $L([-1, 1]) = [0, 1]$. Compute $L([x, y])$.

Sol. $L([x, y]) = \left[\frac{3x+3y}{2}, \frac{-x+y}{2} \right]$

Remark

Let V and W be vector spaces, and let $L : V \rightarrow W$ be a LT. Also, let $\{v_1, v_2, \dots, v_n\}$ be a basis for V .



Exercise

Q:. Suppose $L : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a linear operator and $L([1, 1]) = [3, 0]$ and $L([-1, 1]) = [0, 1]$. Compute $L([x, y])$.

Sol.
$$L([x, y]) = \left[\frac{3x+3y}{2}, \frac{-x+y}{2} \right]$$

Remark

Let V and W be vector spaces, and let $L : V \rightarrow W$ be a LT. Also, let $\{v_1, v_2, \dots, v_n\}$ be a basis for V . If $v \in V$, $L(v)$ is completely determined by $\{L(v_1), L(v_2), \dots, L(v_n)\}$.



Composition of Linear transformations



Composition of Linear transformations

Theorem: Let V_1 , V_2 , and V_3 be vector spaces. Let $L_1 : V_1 \rightarrow V_2$ and $L_2 : V_2 \rightarrow V_3$ be linear transformations. Then $L_2 \circ L_1 : V_1 \rightarrow V_3$ given by $(L_2 \circ L_1)(\mathbf{v}) = L_2(L_1(\mathbf{v}))$, for all $\mathbf{v} \in V_1$, is a LT.



Composition of Linear transformations

Theorem: Let V_1 , V_2 , and V_3 be vector spaces. Let $L_1 : V_1 \rightarrow V_2$ and $L_2 : V_2 \rightarrow V_3$ be linear transformations. Then $L_2 \circ L_1 : V_1 \rightarrow V_3$ given by $(L_2 \circ L_1)(\mathbf{v}) = L_2(L_1(\mathbf{v}))$, for all $\mathbf{v} \in V_1$, is a LT.

Note: $(L_2 \circ L_1)(v)$ is called composite of L_2 with L_1 .



Example 5

Q:. Let $L_1 : P_2 \rightarrow P_2$ and $L_2 : P_2 \rightarrow P_2$ be linear operators.



Example 5

Q:. Let $L_1 : P_2 \rightarrow P_2$ and $L_2 : P_2 \rightarrow P_2$ be linear operators. Also, let $L_1(at^2 + bt + c) = 2at + b$ and $L_2(at^2 + bt + c) = 2at^2 + bt$.



Example 5

Q:. Let $L_1 : P_2 \rightarrow P_2$ and $L_2 : P_2 \rightarrow P_2$ be linear operators. Also, let $L_1(at^2 + bt + c) = 2at + b$ and $L_2(at^2 + bt + c) = 2at^2 + bt$. Compute $L_2 \circ L_1$ and $L_1 \circ L_2$.



Example 5

Q:. Let $L_1 : P_2 \rightarrow P_2$ and $L_2 : P_2 \rightarrow P_2$ be linear operators. Also, let $L_1(at^2 + bt + c) = 2at + b$ and $L_2(at^2 + bt + c) = 2at^2 + bt$. Compute $L_2 \circ L_1$ and $L_1 \circ L_2$.

Sol. $L_2 \circ L_1(at^2 + bt + c) = L_2(L_1(at^2 + bt + c)) = L_2(2at + b) = 2at$.



Example 5

Q:. Let $L_1 : P_2 \rightarrow P_2$ and $L_2 : P_2 \rightarrow P_2$ be linear operators. Also, let $L_1(at^2 + bt + c) = 2at + b$ and $L_2(at^2 + bt + c) = 2at^2 + bt$. Compute $L_2 \circ L_1$ and $L_1 \circ L_2$.

Sol. $L_2 \circ L_1(at^2 + bt + c) = L_2(L_1(at^2 + bt + c)) = L_2(2at + b) = 2at$.

Similarly, $L_1 \circ L_2(at^2 + bt + c) = 4at + b$.



Example 5

Q.: Let $L_1 : P_2 \rightarrow P_2$ and $L_2 : P_2 \rightarrow P_2$ be linear operators. Also, let $L_1(at^2 + bt + c) = 2at + b$ and $L_2(at^2 + bt + c) = 2at^2 + bt$. Compute $L_2 \circ L_1$ and $L_1 \circ L_2$.

Sol. $L_2 \circ L_1(at^2 + bt + c) = L_2(L_1(at^2 + bt + c)) = L_2(2at + b) = 2at$.

Similarly, $L_1 \circ L_2(at^2 + bt + c) = 4at + b$.

Clearly, $L_2 \circ L_1 \neq L_1 \circ L_2$.



Example 6

Q:. Find two linear operators $L_1 : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ and $L_2 : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that $L_2 \circ L_1([a, b]) = [0, 0]$ and $L_1 \circ L_2([a, b]) \neq [0, 0]$.



Example 6

Q:. Find two linear operators $L_1 : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ and $L_2 : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that $L_2 \circ L_1([a, b]) = [0, 0]$ and $L_1 \circ L_2([a, b]) \neq [0, 0]$.

Sol. Let $L_2([a, b]) = [a, 0]$ and $L_1([a, b]) = [0, a]$.



Example 6

Q:. Find two linear operators $L_1 : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ and $L_2 : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that $L_2 \circ L_1([a, b]) = [0, 0]$ and $L_1 \circ L_2([a, b]) \neq [0, 0]$.

Sol. Let $L_2([a, b]) = [a, 0]$ and $L_1([a, b]) = [0, a]$.
 $L_2 \circ L_1([a, b]) = L_2(L_1([a, b])) = L_2([0, a]) = [0, 0]$.



Example 6

Q:. Find two linear operators $L_1 : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ and $L_2 : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that $L_2 \circ L_1([a, b]) = [0, 0]$ and $L_1 \circ L_2([a, b]) \neq [0, 0]$.

Sol. Let $L_2([a, b]) = [a, 0]$ and $L_1([a, b]) = [0, a]$.
 $L_2 \circ L_1([a, b]) = L_2(L_1([a, b])) = L_2([0, a]) = [0, 0]$.
Similarly,
 $L_1 \circ L_2([a, b]) = L_1(L_2([a, b])) = L_1([a, 0]) = [0, a]$.

