

Mathematics-II (MATH F112)

Linear Algebra and Complex Variables

Jitender Kumar

Department of Mathematics
Birla Institute of Technology and Science Pilani
Pilani-333031



Announcements



Announcements

Instructor-Incharge: Dr. Trilok Mathur



Announcements

Instructor-Incharge: Dr. Trilok Mathur

Quizzes: There will be **four unannounced quizzes** of 20 marks each (of time duration 15 minutes) to be conducted in tutorial classes



Announcements

Instructor-Incharge: Dr. Trilok Mathur

Quizzes: There will be **four unannounced quizzes** of 20 marks each (of time duration 15 minutes) to be conducted in tutorial classes . Marks of **Best three** quizzes out of **Four** will be considered for grading.



Students have to write all the Quizzes in their
Registered Tutorial Section Only.



Students have to write all the Quizzes in their
Registered Tutorial Section Only.

Assignments: Two assignments will be given for your practice and does not require submission. However, some questions from assignments may be asked in Mid-sem/Comprehensive examination.



Students have to write all the Quizzes in their
Registered Tutorial Section Only.

Assignments: Two assignments will be given for your practice and does not require submission. However, some questions from assignments may be asked in Mid-sem/Comprehensive examination.

Notices: All course notices will be posted on **NALANDA** and **Department Notice Board.**



Students have to write all the Quizzes in their **Registered Tutorial Section Only.**

Assignments: Two assignments will be given for your practice and does not require submission. However, some questions from assignments may be asked in Mid-sem/Comprehensive examination.

Notices: All course notices will be posted on **NALANDA** and **Department Notice Board.**


For more details about the course structure, please go through the course (MATH F112) handout available on ID website.



Text Book: For Linear Algebra



Text Book: For Linear Algebra



Chapter: 2

- System of Linear equations
- Row Echelon Form
- Elementary Row Operations
- Gaussian Elimination Method
- Reduced Row Echelon Form
- Gauss-Jordan Row Reduction Method
- Rank
- Inverse of a Matrix



An Example for Motivation: Solve the system of linear equations

$$x - y - z = 2$$

$$3x - 3y + 2z = 16$$

$$2x - y + z = 9.$$



An Example for Motivation: Solve the system of linear equations

$$x - y - z = 2$$

$$3x - 3y + 2z = 16$$

$$2x - y + z = 9.$$

Step 1: Represent the given system of equations as follows:

$$\begin{aligned}x - y - z &= 2 \\ 3x - 3y + 2z &= 16 \\ 2x - y + z &= 9\end{aligned}$$

$$\left[\begin{array}{ccc|c} 1 & -1 & -1 & 2 \\ 3 & -3 & 2 & 16 \\ 2 & -1 & 1 & 9 \end{array} \right]$$



Step 2: Multiply the first equation by 3 and subtract it from the 2nd equation; **Multiply the first row by 3 and subtract it from the 2nd row**



Step 2: Multiply the first equation by 3 and subtract it from the 2nd equation; **Multiply the first row by 3 and subtract it from the 2nd row**

$$\begin{aligned}x - y - z &= 2 \\5z &= 10 \\2x - y + z &= 9\end{aligned}$$

$$\left[\begin{array}{ccc|c} 1 & -1 & -1 & 2 \\ 0 & 0 & 5 & 10 \\ 2 & -1 & 1 & 9 \end{array} \right]$$



Step 2: Multiply the first equation by 3 and subtract it from the 2nd equation; **Multiply the first row by 3 and subtract it from the 2nd row**

$$\begin{array}{rcl} x - y - z & = & 2 \\ 5z & = & 10 \\ 2x - y + z & = & 9 \end{array} \quad \left[\begin{array}{ccc|c} 1 & -1 & -1 & 2 \\ 0 & 0 & 5 & 10 \\ 2 & -1 & 1 & 9 \end{array} \right]$$

Step 3: Multiply the first equation by 2 and subtract it from the 3rd equation; **Multiply the first row by 2 and subtract it from the 3rd row**



Step 2: Multiply the first equation by 3 and subtract it from the 2nd equation; **Multiply the first row by 3 and subtract it from the 2nd row**

$$\begin{array}{rcl} x - y - z & = & 2 \\ 5z & = & 10 \\ 2x - y + z & = & 9 \end{array} \quad \left[\begin{array}{ccc|c} 1 & -1 & -1 & 2 \\ 0 & 0 & 5 & 10 \\ 2 & -1 & 1 & 9 \end{array} \right]$$

Step 3: Multiply the first equation by 2 and subtract it from the 3rd equation; **Multiply the first row by 2 and subtract it from the 3rd row**

$$\begin{array}{rcl} x - y - z & = & 2 \\ 5z & = & 10 \\ y + 3z & = & 5 \end{array} \quad \left[\begin{array}{ccc|c} 1 & -1 & -1 & 2 \\ 0 & 0 & 5 & 10 \\ 0 & 1 & 3 & 5 \end{array} \right]$$



Step 4: Interchange 2nd and 3rd equation;
Interchange 2nd and 3rd row



Step 4: Interchange 2nd and 3rd equation;
Interchange 2nd and 3rd row

$$x - y - z = 2$$

$$y + 3z = 5$$

$$5z = 10$$

$$\left[\begin{array}{ccc|c} 1 & -1 & -1 & 2 \\ 0 & 1 & 3 & 5 \\ 0 & 0 & 5 & 10 \end{array} \right]$$



Step 4: Interchange 2nd and 3rd equation;
Interchange 2nd and 3rd row

$$\begin{aligned}x - y - z &= 2 \\y + 3z &= 5 \\5z &= 10\end{aligned}$$

$$\left[\begin{array}{ccc|c} 1 & -1 & -1 & 2 \\ 0 & 1 & 3 & 5 \\ 0 & 0 & 5 & 10 \end{array} \right]$$

Step 5: Multiply the 3rd equation by $\frac{1}{5}$; **Multiply the 3rd row by $\frac{1}{5}$**



Step 4: Interchange 2nd and 3rd equation;
Interchange 2nd and 3rd row

$$\begin{array}{rcl} x - y - z & = & 2 \\ y + 3z & = & 5 \\ 5z & = & 10 \end{array} \quad \left[\begin{array}{ccc|c} 1 & -1 & -1 & 2 \\ 0 & 1 & 3 & 5 \\ 0 & 0 & 5 & 10 \end{array} \right]$$

Step 5: Multiply the 3rd equation by $\frac{1}{5}$; **Multiply the 3rd row by $\frac{1}{5}$**

$$\begin{array}{rcl} x - y - z & = & 2 \\ y + 3z & = & 5 \\ z & = & 2 \end{array} \quad \left[\begin{array}{ccc|c} 1 & -1 & -1 & 2 \\ 0 & 1 & 3 & 5 \\ 0 & 0 & 1 & 2 \end{array} \right]$$



By Backward substitution we find

$$z = 2, \quad y = -1, \quad x = 3$$

is a solution of the given system of equations.



Row Echelon Form (REF): A matrix A is said to be in row echelon form if it satisfies the following properties:



Row Echelon Form (REF): A matrix A is said to be in row echelon form if it satisfies the following properties:

- 1 Any rows consisting entirely of 0's are at the bottom.



Row Echelon Form (REF): A matrix A is said to be in **row echelon form** if it satisfies the following properties:

- 1 Any rows consisting entirely of 0's are at the bottom.
- 2 In each nonzero row, the first nonzero entry (called the **leading entry** or **pivot**) is 1 and it is in a column to the left (strictly) of any leading entry below it.



Row Echelon Form (REF): A matrix A is said to be in **row echelon form** if it satisfies the following properties:

- 1 Any rows consisting entirely of 0's are at the bottom.
- 2 In each nonzero row, the first nonzero entry (called the **leading entry** or **pivot**) is 1 and it is in a column to the left (strictly) of any leading entry below it. (The column containing a pivot element is called a **pivot column**).



- The following matrices are in row echelon form:

$$\begin{bmatrix} \boxed{1} & 2 \\ 0 & \boxed{1} \end{bmatrix}, \begin{bmatrix} \boxed{1} & -1 & -1 \\ 0 & \boxed{1} & 3 \\ 0 & 0 & \boxed{1} \end{bmatrix}, \begin{bmatrix} 0 & \boxed{1} & -1 & 2 & 1 \\ 0 & 0 & \boxed{1} & 4 & 8 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$



- The following matrices are in row echelon form:

$$\begin{bmatrix} \boxed{1} & 2 \\ 0 & \boxed{1} \end{bmatrix}, \begin{bmatrix} \boxed{1} & -1 & -1 \\ 0 & \boxed{1} & 3 \\ 0 & 0 & \boxed{1} \end{bmatrix}, \begin{bmatrix} 0 & \boxed{1} & -1 & 2 & 1 \\ 0 & 0 & \boxed{1} & 4 & 8 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

- If a matrix A is in row echelon form, then in each column of A containing a leading entry, the entries below that leading entry are zero.



- The following matrices are **not** in row echelon form:

$$\begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}, \begin{bmatrix} 1 & -1 & -1 \\ 0 & 0 & 1 \\ 0 & 1 & 3 \end{bmatrix}, \begin{bmatrix} 0 & -1 & -1 & 2 & 1 \\ 1 & 0 & 5 & 10 & 8 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$



Elementary Row Operations: The following row operations are called **elementary row operations** of a matrix:

- Interchange of two rows ($R_i \leftrightarrow R_j$)
- Multiply a row R_i by a nonzero constant c
($R_i \rightarrow cR_i$)
- Add a multiple of a row R_j to another row R_i
($R_i \rightarrow R_i + cR_j$)



Example: Transform the following matrix into row echelon form

$$\begin{bmatrix} 1 & 1 & 1 & 3 \\ 1 & 2 & 2 & 5 \\ 3 & 4 & 4 & 12 \end{bmatrix}$$



Example: Transform the following matrix into row echelon form

$$\begin{bmatrix} 1 & 1 & 1 & 3 \\ 1 & 2 & 2 & 5 \\ 3 & 4 & 4 & 12 \end{bmatrix}$$

Answer:

$$\begin{bmatrix} \boxed{1} & 1 & 1 & 3 \\ 0 & \boxed{1} & 1 & 2 \\ 0 & 0 & 0 & \boxed{1} \end{bmatrix}$$



Example: Transform the following matrix into row echelon form

$$\begin{bmatrix} 1 & 1 & 1 & 3 \\ 1 & 2 & 2 & 5 \\ 3 & 4 & 4 & 12 \end{bmatrix}$$

Answer:

$$\begin{bmatrix} \boxed{1} & 1 & 1 & 3 \\ 0 & \boxed{1} & 1 & 2 \\ 0 & 0 & 0 & \boxed{1} \end{bmatrix}$$

- Row echelon form of a matrix is not **unique**.



Row Equivalent Matrices: Matrices A and B are said to be **row equivalent** if there is a finite sequence of elementary row operations that converts A into B or B into A .



Row Equivalent Matrices: Matrices A and B are said to be **row equivalent** if there is a finite sequence of elementary row operations that converts A into B or B into A .

Example: Matrices

$$A = \begin{bmatrix} 3 & 2 & 7 \\ -4 & 1 & 6 \\ 2 & 5 & 4 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 3 & 2 & 7 \\ -2 & 6 & 10 \\ 2 & 5 & 4 \end{bmatrix}$$

are row equivalent



Row Equivalent Matrices: Matrices A and B are said to be **row equivalent** if there is a finite sequence of elementary row operations that converts A into B or B into A .

Example: Matrices

$$A = \begin{bmatrix} 3 & 2 & 7 \\ -4 & 1 & 6 \\ 2 & 5 & 4 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 3 & 2 & 7 \\ -2 & 6 & 10 \\ 2 & 5 & 4 \end{bmatrix}$$

are row equivalent (**Why?**).



- Every matrix is row equivalent to its row echelon form.



- Every matrix is row equivalent to its row echelon form.
- If a matrix A is row equivalent to a matrix B , then B is row equivalent to A



- Every matrix is row equivalent to its row echelon form.
- If a matrix A is row equivalent to a matrix B , then B is row equivalent to A (Why?).



- Every matrix is row equivalent to its row echelon form.
- If a matrix A is row equivalent to a matrix B , then B is row equivalent to A (Why?).

Result: Matrices A and B are row equivalent if and only if they can be reduced to same row echelon form.



Recall:

- A **vector** is a directed line segment that corresponds to a displacement from one point A to another point B .



Recall:

- A **vector** is a directed line segment that corresponds to a displacement from one point A to another point B . The vector from A to B is denoted by \overrightarrow{AB} .



Recall:

- A **vector** is a directed line segment that corresponds to a displacement from one point A to another point B . The vector from A to B is denoted by \overrightarrow{AB} .
- The point A is called its **initial point** or **tail**, and the point B is called its **terminal point** or **head**.



Recall:

- A **vector** is a directed line segment that corresponds to a displacement from one point A to another point B . The vector from A to B is denoted by \overrightarrow{AB} .
- The point A is called its **initial point** or **tail**, and the point B is called its **terminal point** or **head**.
- The set of all ordered pair of real numbers is denoted by \mathbb{R}^2 i.e. $\mathbb{R}^2 = \{(a, b) \mid a, b \in \mathbb{R}\}$.



Recall:

- A **vector** is a directed line segment that corresponds to a displacement from one point A to another point B . The vector from A to B is denoted by \overrightarrow{AB} .
- The point A is called its **initial point** or **tail**, and the point B is called its **terminal point** or **head**.
- The set of all ordered pair of real numbers is denoted by \mathbb{R}^2 i.e. $\mathbb{R}^2 = \{(a, b) \mid a, b \in \mathbb{R}\}$.
- The set \mathbb{R}^2 corresponds to the set of vectors whose tails are at the origin O .



- For example, the ordered pair $A = (3, 2) \in \mathbb{R}^2$ corresponds to the vector \overrightarrow{OA} and we denote it as square bracket $[3, 2]$.



For $n \in \mathbb{N}$, \mathbb{R}^n is the set of all ordered n -tuples (x_1, x_2, \dots, x_n) , where $x_i \in \mathbb{R}$.

- We can think the point $(x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ as vector and write it as $[x_1, x_2, \dots, x_n]$ (row vector). Thus,

$$\mathbb{R}^n = \{[x_1, x_2, \dots, x_n] \mid x_i \in \mathbb{R}\}.$$



- Sometime we will write a vector of \mathbb{R}^n as a **column vector**

$$\begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = [x_1, x_2, \dots, x_n]^T,$$



- Sometime we will write a vector of \mathbb{R}^n as a **column vector**

$$\begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = [x_1, x_2, \dots, x_n]^T,$$

depend on the situation.



- Sometime we will write a vector of \mathbb{R}^n as a **column vector**

$$\begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = [x_1, x_2, \dots, x_n]^T,$$

depend on the situation.

- The vector $[0, 0, \dots, 0]$ of \mathbb{R}^n , called the **zero vector** of \mathbb{R}^n and it is denoted by the symbol **0**.



Let $\mathbf{u} = [u_1, u_2, \dots, u_n]$ and $\mathbf{v} = [v_1, v_2, \dots, v_n] \in \mathbb{R}^n$
and $k \in \mathbb{R}$.



Let $\mathbf{u} = [u_1, u_2, \dots, u_n]$ and $\mathbf{v} = [v_1, v_2, \dots, v_n] \in \mathbb{R}^n$
and $k \in \mathbb{R}$. Define

- $\mathbf{u} + \mathbf{v} = [u_1 + v_1, u_2 + v_2, \dots, u_n + v_n]$ (Vector addition)



Let $\mathbf{u} = [u_1, u_2, \dots, u_n]$ and $\mathbf{v} = [v_1, v_2, \dots, v_n] \in \mathbb{R}^n$ and $k \in \mathbb{R}$. Define

- $\mathbf{u} + \mathbf{v} = [u_1 + v_1, u_2 + v_2, \dots, u_n + v_n]$ (Vector addition)
- $k\mathbf{u} = [ku_1, ku_2, \dots, ku_n]$ (Scalar Multiplication)



Some Basic Properties: Let $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathbb{R}^n$ and $c, d \in \mathbb{R}$. Then



Some Basic Properties: Let $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathbb{R}^n$ and $c, d \in \mathbb{R}$. Then

- $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$ (commutativity).
- $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}$ (associativity).
- $\mathbf{u} + \mathbf{0} = \mathbf{u}$.
- $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$, where $-\mathbf{u} = [-u_1, -u_2, \dots, -u_n]$.
- $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$ (distributivity over vector addition).
- $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$ (distributivity over scalar addition).
- $c(d\mathbf{u}) = (cd)\mathbf{u}$.
- $1\mathbf{u} = \mathbf{u}$.
- $0\mathbf{u} = \mathbf{0}$.



System of Linear Equations



System of Linear Equations

A system of **m** linear equations in **n** unknown variables x_1, x_2, \dots, x_n is given by

$$\begin{array}{cccc} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n & = & b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n & = & b_2 \\ \vdots & & \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n & = & b_m \end{array}$$

where $a_{ij}, b_i \in \mathbb{R}$ and $1 \leq i \leq m, 1 \leq j \leq n$.



System of Linear Equations

A system of m linear equations in n unknown variables x_1, x_2, \dots, x_n is given by

$$\begin{aligned}a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n &= b_1 \\a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n &= b_2 \\ \vdots & \quad \quad \quad \vdots \quad \quad \quad \vdots \\a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n &= b_m\end{aligned}$$

where $a_{ij}, b_i \in \mathbb{R}$ and $1 \leq i \leq m, 1 \leq j \leq n$.

- A solution of the linear system is an n -tuple (s_1, s_2, \dots, s_n) such that each equation of the system is satisfied by substituting s_i in place of x_i .



Above linear system of equations can be written in the form $AX = B$, where



Above linear system of equations can be written in the form $AX = B$, where

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \vdots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}, X = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \text{ and } B = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$$



Above linear system of equations can be written in the form $AX = B$, where

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \vdots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}, X = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \text{ and } B = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$$

- The matrix A is called the **coefficient matrix**.



- The matrix $[A|B]$ which is formed by inserting the column of matrix B next to the column of A , is called the **augmented matrix** of the linear system $AX = B$ i.e.



- The matrix $[A|B]$ which is formed by inserting the column of matrix B next to the column of A , is called the **augmented matrix** of the linear system $AX = B$ i.e.

$$[A|B] = \left[\begin{array}{cccc|c} a_{11} & a_{12} & \dots & a_{1n} & b_1 \\ a_{21} & a_{22} & \dots & a_{2n} & b_2 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} & b_m \end{array} \right]$$



- The matrix $[A|B]$ which is formed by inserting the column of matrix B next to the column of A , is called the **augmented matrix** of the linear system $AX = B$ i.e.

$$[A|B] = \left[\begin{array}{cccc|c} a_{11} & a_{12} & \dots & a_{1n} & b_1 \\ a_{21} & a_{22} & \dots & a_{2n} & b_2 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} & b_m \end{array} \right]$$

- The vertical bar is used in the augmented matrix $[A|B]$ only to distinguish the column vector B from the coefficient matrix A .



- If $B = 0 = [0, 0, \dots, 0]^T$ i.e. $b_1 = 0 = b_2 = \dots = b_m$, the system $AX = 0$ is called **homogenous** system of equations.



- If $B = 0 = [0, 0, \dots, 0]^T$ i.e. $b_1 = 0 = b_2 = \dots = b_m$, the system $AX = 0$ is called **homogenous** system of equations.
- If $B \neq 0$, then the system $AX = B$ is called **non-homogenous** system of equations.



- If $B = 0 = [0, 0, \dots, 0]^T$ i.e. $b_1 = 0 = b_2 = \dots = b_m$, the system $AX = 0$ is called **homogenous** system of equations.
- If $B \neq 0$, then the system $AX = B$ is called **non-homogenous** system of equations.
- The solution $X = 0$ of the system $AX = 0$ is called the **trivial** solution



- If $B = 0 = [0, 0, \dots, 0]^T$ i.e. $b_1 = 0 = b_2 = \dots = b_m$, the system $AX = 0$ is called **homogenous** system of equations.
- If $B \neq 0$, then the system $AX = B$ is called **non-homogenous** system of equations.
- The solution $X = 0$ of the system $AX = 0$ is called the **trivial** solution and a solution other than $X = 0$ is called a **non-trivial** solution of the homogenous system $AX = 0$.



Equivalent Systems: Two system of m linear equations in n variables are **equivalent** if and only if they have exactly the same solution set.



Equivalent Systems: Two system of m linear equations in n variables are **equivalent** if and only if they have exactly the same solution set.

Example: The systems

$$\begin{array}{l} 2x - y = 1 \\ 3x + y = 9 \end{array} \quad \text{and} \quad \begin{array}{l} x + 4y = 14 \\ 5x - 2y = 4 \end{array}$$

are equivalent



Equivalent Systems: Two system of m linear equations in n variables are **equivalent** if and only if they have exactly the same solution set.

Example: The systems

$$\begin{array}{l} 2x - y = 1 \\ 3x + y = 9 \end{array} \quad \text{and} \quad \begin{array}{l} x + 4y = 14 \\ 5x - 2y = 4 \end{array}$$

are equivalent (**Why?**).



Theorem: Let $AX = B$ be a system of linear equations. If $[C|D]$ is row equivalent to $[A|B]$, then the system $CX = D$ is equivalent to $AX = B$.



Gaussian Elimination Method: Use the following steps to solve a system of equations $AX = B$



Gaussian Elimination Method: Use the following steps to solve a system of equations $AX = B$

- Write the augmented matrix $[A \mid B]$.



Gaussian Elimination Method: Use the following steps to solve a system of equations $AX = B$

- Write the augmented matrix $[A \mid B]$.
- Find a row echelon form of the matrix $[A \mid B]$.



Gaussian Elimination Method: Use the following steps to solve a system of equations $AX = B$

- Write the augmented matrix $[A \mid B]$.
- Find a row echelon form of the matrix $[A \mid B]$.
- Use back substitution to solve the equivalent system that corresponds to row echelon form.



Exercise: Solve the linear system of equations by Gaussian elimination method

$$x + y + z = 3$$

$$2x + 3z = 5$$

$$y + z = 2$$



Exercise: Solve the linear system of equations

$$x + y + z = 3, \quad x + 2y + 2z = 5, \quad 3x + 4y + 4z = 12$$

by Gaussian elimination method.



Independent and dependent variables:



Independent and dependent variables:

- Consider the linear system $AX = B$ in n variables and m equations.



Independent and dependent variables:

- Consider the linear system $AX = B$ in n variables and m equations.
- Let $[C \mid D]$ be a row echelon form of the augmented matrix $[A \mid B]$.



Independent and dependent variables:

- Consider the linear system $AX = B$ in n variables and m equations.
- Let $[C \mid D]$ be a row echelon form of the augmented matrix $[A \mid B]$.
- The variables corresponding to the **pivot** columns in the first n columns of $[C \mid D]$ are called the **dependent** (or basic) variables.



Independent and dependent variables:

- Consider the linear system $AX = B$ in n variables and m equations.
- Let $[C \mid D]$ be a row echelon form of the augmented matrix $[A \mid B]$.
- The variables corresponding to the **pivot** columns in the first n columns of $[C \mid D]$ are called the **dependent** (or basic) variables.
- The variables which are not dependent are called **independent** (free) variables.



Exercise: Solve the system of linear equations

$$x + y + z = 3, \quad x + 2y + 2z = 5, \quad 3x + 4y + 4z = 11$$

by Gaussian elimination method.



Reduced Row Echelon Form (RREF): A matrix A is said to be in **reduced row echelon form** if it satisfies the following properties:



Reduced Row Echelon Form (RREF): A matrix A is said to be in **reduced row echelon form** if it satisfies the following properties:

- A is in row echelon form.



Reduced Row Echelon Form (RREF): A matrix A is said to be in **reduced row echelon form** if it satisfies the following properties:

- A is in row echelon form.
- If a column contains a **leading entry (or pivot)** then all other entries in that column must be zero.



Reduced Row Echelon Form (RREF): A matrix A is said to be in **reduced row echelon form** if it satisfies the following properties:

- A is in row echelon form.
- If a column contains a **leading entry (or pivot)** then all other entries in that column must be zero.

Example: The following matrices are in reduced row echelon form

$$\begin{bmatrix} \boxed{1} & 0 \\ 0 & \boxed{1} \end{bmatrix}, \begin{bmatrix} \boxed{1} & 0 & 0 \\ 0 & 0 & \boxed{1} \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & \boxed{1} & 0 & 2 & 1 \\ 0 & 0 & \boxed{1} & 4 & 8 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$



Result: Every matrix has a **unique** reduced row echelon form.



Gauss-Jordan Row Reduction Method: Use the following steps to solve a system of equations

$$AX = B$$



Gauss-Jordan Row Reduction Method: Use the following steps to solve a system of equations

$$AX = B$$

- Write the augmented matrix $[A \mid B]$.



Gauss-Jordan Row Reduction Method: Use the following steps to solve a system of equations

$$AX = B$$

- Write the augmented matrix $[A \mid B]$.
- Find the reduced row echelon form of the matrix $[A \mid B]$.



Gauss-Jordan Row Reduction Method: Use the following steps to solve a system of equations

$$AX = B$$

- Write the augmented matrix $[A \mid B]$.
- Find the reduced row echelon form of the matrix $[A \mid B]$.
- Use back substitution to solve the equivalent system that corresponds to the reduced row echelon form.



Exercise: Solve the system of linear equations equations

$$x + y + z = 5, \quad 2x + 3y + 5z = 8, \quad 4x + 5z = 2$$

by Gauss-Jordan method.



Exercise: Solve the system of linear equations

$$x + y + z = 5, \quad 2x + 3y + 5z = 8, \quad 4x + 5z = 2$$

by Gauss-Jordan method.

Answer: $x = 3, y = 4$ and $z = -2$.



Exercise: Solve the system of linear equations

$$4y + z = 2, \quad 2x + 6y - 2z = 3, \quad 4x + 8y - 5z = 4$$

by Gauss-Jordan method.



Exercise: Solve the system of linear equations

$$4y + z = 2, \quad 2x + 6y - 2z = 3, \quad 4x + 8y - 5z = 4$$

by Gauss-Jordan method.

Answer: Infinitely many solutions and the solution set is

$$\left\{ \left(\frac{7}{4}d, \frac{1}{2} - \frac{1}{4}d, d \right) \mid d \in \mathbb{R} \right\}.$$



Exercise: Solve the system of linear equations

$$x + 2y - 3z = 2, \quad 6x + 3y - 9z = 6, \quad 7x + 14y - 21z = 13$$

by Gauss-Jordan method.



Exercise: Solve the system of linear equations

$$x + 2y - 3z = 2, \quad 6x + 3y - 9z = 6, \quad 7x + 14y - 21z = 13$$

by Gauss-Jordan method.

Answer: No solution.



So far we solved linear systems using Gauss elimination method or the Gauss-Jordan method. In the examples considered, we have encountered three possibilities, namely

- existence of a unique solution



So far we solved linear systems using Gauss elimination method or the Gauss-Jordan method. In the examples considered, we have encountered three possibilities, namely

- existence of a unique solution
- existence of an infinite number of solutions, and



So far we solved linear systems using Gauss elimination method or the Gauss-Jordan method. In the examples considered, we have encountered three possibilities, namely

- existence of a unique solution
- existence of an infinite number of solutions, and
- no solution.



So far we solved linear systems using Gauss elimination method or the Gauss-Jordan method. In the examples considered, we have encountered three possibilities, namely

- existence of a unique solution
 - existence of an infinite number of solutions, and
 - no solution.
-
- If the system $AX = B$ has some solution then it is called a **consistent** system. Otherwise it is called an **inconsistent** system.



Question: Whether there are conditions under which the linear system $AX = B$ is consistent?



Question: Whether there are conditions under which the linear system $AX = B$ is consistent?

Rank: The **rank** of a matrix A is the number of nonzero rows in its row echelon form.



Question: Whether there are conditions under which the linear system $AX = B$ is consistent?

Rank: The **rank** of a matrix A is the number of nonzero rows in its row echelon form. It is denoted by $\text{rank}(A)$.



Question: Whether there are conditions under which the linear system $AX = B$ is consistent?

Rank: The **rank** of a matrix A is the number of nonzero rows in its row echelon form. It is denoted by $\text{rank}(A)$.

Remark: The number of nonzero rows in either the row echelon form or the reduced row echelon form of a matrix are same.



Exercise: Determine the rank of $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 3 & 1 \\ 1 & 1 & 0 \end{bmatrix}$.



Theorem: Let $AX = B$ be a system of equations with n variables.

- 1 If $\text{rank}(A) = \text{rank}([A \mid B]) = n$ then the system $AX = B$ has a unique solution.



Theorem: Let $AX = B$ be a system of equations with n variables.

- 1 If $\text{rank}(A) = \text{rank}([A \mid B]) = n$ then the system $AX = B$ has a unique solution.
- 2 if $\text{rank}(A) = \text{rank}([A \mid B]) < n$ then the system $AX = B$ has a infinitely many solutions.



Theorem: Let $AX = B$ be a system of equations with n variables.

- 1 If $\text{rank}(A) = \text{rank}([A \mid B]) = n$ then the system $AX = B$ has a unique solution.
- 2 if $\text{rank}(A) = \text{rank}([A \mid B]) < n$ then the system $AX = B$ has a infinitely many solutions.
- 3 If $\text{rank}(A) \neq \text{rank}([A \mid B])$ then the system $AX = B$ is inconsistent.



Theorem: Let $AX = 0$ be a homogenous system of equations with n variables.

- 1 If $\text{rank}(A) = n$ then the system has a unique solution (**trivial solution**).



Theorem: Let $AX = 0$ be a homogenous system of equations with n variables.

- 1 If $\text{rank}(A) = n$ then the system has a unique solution (**trivial solution**).
- 2 If $\text{rank}(A) < n$ then the system $AX = B$ has infinitely many solutions.



Exercise: Test the consistency of the given system of equations

$$3x + y + w = -9$$

$$-2y + 12z - 8w = -6$$

$$2x - 3y + 22z - 14w = -17.$$

Find all the solutions, if it is consistent.



Exercise: For what value of $\lambda \in \mathbb{R}$, the following system of equations has (i) a unique solution (ii) infinitely many solutions and (iii) no solution

$$(5 - \lambda)x + 4y + 2z = 4$$

$$4x + (5 - \lambda)y + 2z = 4$$

$$2x + 2y + (2 - \lambda)z = 2.$$



Exercise: For what value of $\lambda \in \mathbb{R}$, the following system of equations has (i) a unique solution (ii) infinitely many solutions and (iii) no solution

$$(5 - \lambda)x + 4y + 2z = 4$$

$$4x + (5 - \lambda)y + 2z = 4$$

$$2x + 2y + (2 - \lambda)z = 2.$$

Also find the solutions, whenever they exist.



Definition: Let A be an $n \times n$ matrix. Then an $n \times n$ matrix B is a (multiplicative) **inverse** of A if and only if

$$AB = BA = I_n,$$

where I_n is the $n \times n$ identity matrix.



Definition: Let A be an $n \times n$ matrix. Then an $n \times n$ matrix B is a (multiplicative) **inverse** of A if and only if

$$AB = BA = I_n,$$

where I_n is the $n \times n$ identity matrix.

- If such a matrix B exists then A is called **nonsingular**. Otherwise it is called **singular**.



Example: Show that the matrix $A = \begin{bmatrix} 2 & 5 \\ 1 & 3 \end{bmatrix}$ is nonsingular.



Example: Show that the matrix $A = \begin{bmatrix} 2 & 5 \\ 1 & 3 \end{bmatrix}$ is nonsingular.

Theorem: Let A and B be $n \times n$ matrices.

- If $AB = I_n$ then $BA = I_n$.



Example: Show that the matrix $A = \begin{bmatrix} 2 & 5 \\ 1 & 3 \end{bmatrix}$ is nonsingular.

Theorem: Let A and B be $n \times n$ matrices.

- If $AB = I_n$ then $BA = I_n$.
- If $BA = I_n$ then $AB = I_n$.



Example: Show that the matrix $A = \begin{bmatrix} 2 & 5 \\ 1 & 3 \end{bmatrix}$ is nonsingular.

Theorem: Let A and B be $n \times n$ matrices.

- If $AB = I_n$ then $BA = I_n$.
- If $BA = I_n$ then $AB = I_n$.

Theorem: Inverse of a matrix is **unique** if it exists.



As the inverse of a matrix A is unique, we denote it by A^{-1} . That is, $AA^{-1} = A^{-1}A = I$.



As the inverse of a matrix A is unique, we denote it by A^{-1} . That is, $AA^{-1} = A^{-1}A = I$.

Theorem: Let A and B be an $n \times n$ nonsingular matrices. Then

- $(A^{-1})^{-1} = A$.



As the inverse of a matrix A is unique, we denote it by A^{-1} . That is, $AA^{-1} = A^{-1}A = I$.

Theorem: Let A and B be an $n \times n$ nonsingular matrices. Then

- $(A^{-1})^{-1} = A$.
- $(AB)^{-1} = B^{-1}A^{-1}$.



As the inverse of a matrix A is unique, we denote it by A^{-1} . That is, $AA^{-1} = A^{-1}A = I$.

Theorem: Let A and B be an $n \times n$ nonsingular matrices. Then

- $(A^{-1})^{-1} = A$.
- $(AB)^{-1} = B^{-1}A^{-1}$.
- $(A^T)^{-1} = (A^{-1})^T$.



Question:

- How can we know when a matrix has an inverse?



Question:

- How can we know when a matrix has an inverse?
- If a matrix does have an inverse, how can we find it?



Method for finding the Inverse of a matrix (if it exists): Let A be a given $n \times n$ matrix.



Method for finding the Inverse of a matrix (if it exists): Let A be a given $n \times n$ matrix.

Step 1: Write the augmented matrix $[A \mid I_n]$.



Method for finding the Inverse of a matrix (if it exists): Let A be a given $n \times n$ matrix.

Step 1: Write the augmented matrix $[A \mid I_n]$.

Step 2: Transform the augmented matrix $[A \mid I_n]$ to the matrix $[C \mid D]$ in reduced row echelon form via elementary row operations.



Method for finding the Inverse of a matrix (if it exists): Let A be a given $n \times n$ matrix.

Step 1: Write the augmented matrix $[A \mid I_n]$.

Step 2: Transform the augmented matrix $[A \mid I_n]$ to the matrix $[C \mid D]$ in reduced row echelon form via elementary row operations.

Step 3: If

- $C = I_n$ then $D = A^{-1}$.



Method for finding the Inverse of a matrix (if it exists): Let A be a given $n \times n$ matrix.

Step 1: Write the augmented matrix $[A \mid I_n]$.

Step 2: Transform the augmented matrix $[A \mid I_n]$ to the matrix $[C \mid D]$ in reduced row echelon form via elementary row operations.

Step 3: If

- $C = I_n$ then $D = A^{-1}$.
- $C \neq I_n$ then A is singular and A^{-1} does not exist.



Exercise: Using row reduction method, find the inverse of $A = \begin{bmatrix} 1 & -1 & -2 \\ 2 & -3 & -5 \\ -1 & 3 & 5 \end{bmatrix}$, if it exists.



Exercise: Using row reduction method, find the inverse of $A = \begin{bmatrix} 1 & -1 & -2 \\ 2 & -3 & -5 \\ -1 & 3 & 5 \end{bmatrix}$, if it exists.

Hint: Note that reduced row echelon form of the matrix $[A|I_3]$ is

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 5 & -3 & -1 \\ 0 & 0 & 1 & -3 & 2 & 1 \end{array} \right]$$



Exercise: Using row reduction method, find the inverse of $A = \begin{bmatrix} 1 & -1 & -2 \\ 2 & -3 & -5 \\ -1 & 3 & 5 \end{bmatrix}$, if it exists.

Hint: Note that reduced row echelon form of the matrix $[A|I_3]$ is

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 5 & -3 & -1 \\ 0 & 0 & 1 & -3 & 2 & 1 \end{array} \right]$$

Thus, $A^{-1} = \begin{bmatrix} 0 & 1 & 1 \\ 5 & -3 & -1 \\ -3 & 2 & 1 \end{bmatrix}$



Theorem: Let A be an $n \times n$ matrix. The following statements are equivalent:

- A is nonsingular.
- The homogenous system $AX = 0$ has only the trivial solution.
- $\text{rank}(A) = n$.
- The reduced row echelon form of A is I_n .



Thank You

