

MATHEMATICS-I (MATH F111)

Dr. Krishnendra Shekhawat

BITS PILANI
Department of Mathematics



CHAPTER 11

Conic Sections and Polar Coordinates



Topics to be covered in this Chapter:



Topics to be covered in this Chapter:

❶ Polar Coordinates



Topics to be covered in this Chapter:

- ① Polar Coordinates
- ② Curve Tracing



Topics to be covered in this Chapter:

- ① Polar Coordinates
- ② Curve Tracing
- ③ Faster Graphing



Topics to be covered in this Chapter:

- ① Polar Coordinates
- ② Curve Tracing
- ③ Faster Graphing
- ④ Areas and Lengths of the Curves



Topics to be covered in this Chapter:

- ① Polar Coordinates
- ② Curve Tracing
- ③ Faster Graphing
- ④ Areas and Lengths of the Curves
- ⑤ Conic Sections



Section 11.3

Polar Coordinates



- How can be describe or represent a curve?



- How can be describe or represent a curve?
- Cartesian coordinates




- How can be describe or represent a curve?
- Cartesian coordinates
- Polar coordinates



Cartesian Coordinates

A point in cartesian coordinate or $x-y$ co-ordinate or rectangular co-ordinate is described in terms of horizontal and vertical distances from the origin $(0,0)$.



Polar Coordinates

To define polar coordinates in a plane, we start with an origin (called the pole) and an initial ray.




Figure: θ is an angle made by the ray OP with initial ray and r is the distance of P from O along ray OP .




Plotting of Polar co-ordinates

Plot the following points:

- $(2, 0)$
- $(3, \frac{\pi}{2})$
- $(-3, \frac{\pi}{2})$
- $(2, \pi)$





Polar Coordinates with Negative r Values

The points $(-r, \theta)$ and (r, θ) lie on the same line through the pole O and at the distance $|r|$ from O , but on opposite sides of O .



Polar Coordinates with Negative r Values

The points $(-r, \theta)$ and (r, θ) lie on the same line through the pole O and at the distance $|r|$ from O , but on opposite sides of O .

- If $r > 0$, the point (r, θ) is in the same quadrant as θ .



Polar Coordinates with Negative r Values


The points $(-r, \theta)$ and (r, θ) lie on the same line through the pole O and at the distance $|r|$ from O , but on opposite sides of O .

- If $r > 0$, the point (r, θ) is in the same quadrant as θ .
- If $r < 0$, the point (r, θ) is in the quadrant opposite of the angle θ , *i.e.*, opposite of pole.



Different representation of Polar co-ordinates

Plot $(\sqrt{2}, \pi/4)$



The number of Polar Coordinate Pairs

Q:. How many Polar Coordinate Pairs the point (r, θ) can have?



The number of Polar Coordinate Pairs

Q:. How many Polar Coordinate Pairs the point (r, θ) can have?

Sol. Infinite



The different polar coordinates of a point (r, θ) are:



The different polar coordinates of a point (r, θ) are:

- $(r, \theta + 2n\pi), \quad n = 0, \pm 1, \pm 2, \dots$



The different polar coordinates of a point (r, θ) are:

- $(r, \theta + 2n\pi), \quad n = 0, \pm 1, \pm 2, \dots$
- $(-r, \theta + (2n + 1)\pi), \quad n = 0, \pm 1, \pm 2, \dots$

Remark. If $P \equiv O$, then $r = 0$ and $(0, \theta)$ represents pole for any value of θ .






Figure: The point $(2, \frac{\pi}{6})$ has infinitely many polar coordinate pairs



Q: $0 \leq \theta \leq \frac{\pi}{6}, r \geq 0.$



Q:. $0 \leq \theta \leq \frac{\pi}{6}$, $r \geq 0$.

Sol.




Figure: An infinite region



Q:. $0 \leq \theta \leq \frac{\pi}{6}$, $r \geq 0$.

Sol.




Figure: An infinite region



Q:.. $\theta = \frac{\pi}{3}, -1 \leq r \leq 3.$



Q:. $\theta = \frac{\pi}{3}$, $-1 \leq r \leq 3$.

Sol.





Figure: A line segment



Q: $r = a$



Q:. $r = a$ (The Polar Equation of a Circle with radius $|a|$ centered at O)




Q:. Graph the set of points whose polar coordinates satisfy the inequality $1 \leq r \leq 2$.



Q:. Graph the set of points whose polar coordinates satisfy the inequality $1 \leq r \leq 2$.

Sol.




Q:.. $1 \leq |r| \leq 2, 0 \leq \theta \leq \pi/2$




Q:.. $1 \leq |r| \leq 2, 0 \leq \theta \leq \pi/2$

Q:.. $r = \theta, \theta \geq 0$









The Graph of Some Inequalities


(a)




(c)



(b)



(d)



Homework. Graph the set of points whose polar coordinates satisfy the inequality:

- $-2 \leq r \leq -1$.



Homework. Graph the set of points whose polar coordinates satisfy the inequality:

- $-2 \leq r \leq -1$. (An annulus).



Homework. Graph the set of points whose polar coordinates satisfy the inequality:

- $-2 \leq r \leq -1$. (An annulus).
- $-1 \leq r \leq 2$.



Homework. Graph the set of points whose polar coordinates satisfy the inequality:

- $-2 \leq r \leq -1$. (An annulus).
- $-1 \leq r \leq 2$. (Interior of circle $r = 2$ including boundary).



Homework. Graph the set of points whose polar coordinates satisfy the inequality:

- $-2 \leq r \leq -1$. (An annulus).
- $-1 \leq r \leq 2$. (Interior of circle $r = 2$ including boundary).
- $r \leq 1$.



Homework. Graph the set of points whose polar coordinates satisfy the inequality:

- $-2 \leq r \leq -1$. (An annulus).
- $-1 \leq r \leq 2$. (Interior of circle $r = 2$ including boundary).
- $r \leq 1$. (Whole plane).



Homework. Graph the set of points whose polar coordinates satisfy the inequality:

- $-2 \leq r \leq -1$. (An annulus).
- $-1 \leq r \leq 2$. (Interior of circle $r = 2$ including boundary).
- $r \leq 1$. (Whole plane).
- $r \geq -1$. (Whole plane).



Relation between Polar and Cartesian Coordinates




Figure: The usual way to relate polar and cartesian coordinates



From figure, we have

$$x = r \cos \theta, \quad y = r \sin \theta,$$



From figure, we have

$$x = r \cos \theta, \quad y = r \sin \theta,$$

on squaring and adding:

$$x^2 + y^2 = r^2,$$



From figure, we have

$$x = r \cos \theta, \quad y = r \sin \theta,$$

on squaring and adding:

$$x^2 + y^2 = r^2,$$

on dividing:

$$\theta = \tan^{-1} \frac{y}{x}.$$



Q:.. Replace $r = \sin \theta$ by an equivalent cartesian equation.



Q:. Replace $r = \sin \theta$ by an equivalent cartesian equation.
Sol.

$$r = \sin \theta$$

$$\Rightarrow r^2 = r \sin \theta$$

$$\Rightarrow x^2 + y^2 = y$$

$$\Rightarrow x^2 + y^2 - y = 0$$

$$\Rightarrow x^2 + \left(y - \frac{1}{2}\right)^2 = \left(\frac{1}{2}\right)^2.$$



Q:.. Replace $r^2(1 + \sin 2\theta) = 1$ by an equivalent cartesian equation.



Q:. Replace $r^2(1 + \sin 2\theta) = 1$ by an equivalent cartesian equation.

Sol.

$$\begin{aligned}r^2 + 2r^2 \sin \theta \cos \theta &= 1 \\ \Rightarrow r^2 + 2(r \sin \theta)(r \cos \theta) &= 1 \\ \Rightarrow x^2 + y^2 + 2xy &= 1 \\ \Rightarrow x + y &= \pm 1.\end{aligned}$$



Q:.. Replace $x^2 - y^2 = 9$ by an equivalent polar equation.



Q:. Replace $x^2 - y^2 = 9$ by an equivalent polar equation.
Sol.

$$\begin{aligned}x^2 - y^2 &= 9 \\ \Rightarrow r^2 \cos^2 \theta - r^2 \sin^2 \theta &= 9 \\ \Rightarrow r^2 (\cos^2 \theta - \sin^2 \theta) &= 9 \\ \Rightarrow r^2 \cos 2\theta &= 9 \\ \Rightarrow r &= \pm 3 \sqrt{\sec 2\theta}.\end{aligned}$$



Q:. Replace $(x - 5)^2 + y^2 = 25$ by an equivalent polar equation.



Q:. Replace $(x - 5)^2 + y^2 = 25$ by an equivalent polar equation.

Sol.

$$\begin{aligned}(x - 5)^2 + y^2 &= 25 \\ \Rightarrow x^2 - 10x + 25 + y^2 &= 25 \\ \Rightarrow r^2 - 10r \cos \theta &= 0 \\ \Rightarrow r &= 10 \cos \theta.\end{aligned}$$



Section 11.4

Graphing in Polar Coordinates



How to Trace a Curve in Polar Coordinate



How to Trace a Curve in Polar Coordinate

Trace the Curve $r = 1 - \cos\theta$



How to Trace a Curve in Polar Coordinate

Trace the Curve $r = 1 - \cos\theta$

θ	0	$\frac{\pi}{4}$	$\frac{\pi}{2}$	$\frac{3\pi}{4}$	π	$\frac{5\pi}{4}$	$\frac{3\pi}{2}$	$\frac{7\pi}{4}$	2π
r	0	0.29	1	1.71	2	1.71	1	0.29	0






Figure: $r = 1 - \cos\theta$



Q:.. Do you see symmetry in above Figure ?



Q:.. Do you see symmetry in above Figure ? What is its role?



Q:.. Do you see symmetry in above Figure ? What is its role?

Q:.. Where can be draw tangent in the above Figure



Q:. Do you see symmetry in above Figure ? What is its role?

Q:. Where can be draw tangent in the above Figure (slope of a curve) ?



Curves in Polar Coordinates

- For polar coordinates (r, θ) , the equation $f(r, \theta) = 0$ (implicit form) or $r = f(\theta)$ (explicit form) defines a curve C in the plane.



Curves in Polar Coordinates

- For polar coordinates (r, θ) , the equation $f(r, \theta) = 0$ (implicit form) or $r = f(\theta)$ (explicit form) defines a curve C in the plane.
- A point P lies on C if and only if for at least one polar coordinate (r_0, θ_0) of P , $f(r_0, \theta_0) = 0$.



Curves in Polar Coordinates

- For polar coordinates (r, θ) , the equation $f(r, \theta) = 0$ (implicit form) or $r = f(\theta)$ (explicit form) defines a curve C in the plane.
- A point P lies on C if and only if for at least one polar coordinate (r_0, θ_0) of P , $f(r_0, \theta_0) = 0$.

In this section, our aim is to trace the curve $r = f(\theta)$ or $f(r, \theta) = 0$.




Symmetry Test 1

- **Symmetry about x -axis:** If the point (r, θ) lies on the graph, the point $(r, -\theta)$ or $(-r, \pi - \theta)$ also lies on the graph.



Symmetry Test 1

- **Symmetry about x -axis:** If the point (r, θ) lies on the graph, the point $(r, -\theta)$ or $(-r, \pi - \theta)$ also lies on the graph.




Symmetry Test 2

- **Symmetry about the y-axis:** If the point (r, θ) lies on the graph, the point $(-r, -\theta)$ or $(r, \pi - \theta)$ also lies on the graph.



Symmetry Test 2

- **Symmetry about the y-axis:** If the point (r, θ) lies on the graph, the point $(-r, -\theta)$ or $(r, \pi - \theta)$ also lies on the graph.



Symmetry Test 3 and 4

- **Symmetry about the origin:** If the point (r, θ) lies on the graph, the point $(-r, \theta)$ or $(r, \pi + \theta)$ also lies on the graph.



Symmetry Test 3 and 4

- **Symmetry about the origin:** If the point (r, θ) lies on the graph, the point $(-r, \theta)$ or $(r, \pi + \theta)$ also lies on the graph.
- **Symmetry about the line $y = x$:** If the point (r, θ) lies on the graph, the point $(r, \frac{\pi}{2} - \theta)$ or $(-r, -\frac{\pi}{2} - \theta)$ also lies on the graph.






Figure: Symmetry about the pole and line $y = x$ respectively



Q: Identify the symmetries of the following curves.



Q:. Identify the symmetries of the following curves.

- $r = 2 + \sin \theta$



Q:. Identify the symmetries of the following curves.

- $r = 2 + \sin \theta$ (symmetry about y -axis)



Q:. Identify the symmetries of the following curves.

- $r = 2 + \sin \theta$ (symmetry about y -axis)

- $r^2 = 4 \sin 2\theta$



Q:. Identify the symmetries of the following curves.

- $r = 2 + \sin \theta$ (symmetry about y-axis)
- $r^2 = 4 \sin 2\theta$ (symmetry about the origin)



Q:. Identify the symmetries of the following curves.

- $r = 2 + \sin \theta$ (symmetry about y-axis)
- $r^2 = 4 \sin 2\theta$ (symmetry about the origin)
- $r^2 = 4 \cos 2\theta$



Q:. Identify the symmetries of the following curves.

- $r = 2 + \sin \theta$ (symmetry about y -axis)
- $r^2 = 4 \sin 2\theta$ (symmetry about the origin)
- $r^2 = 4 \cos 2\theta$ (symmetry about x -axis, y -axis, the origin)



Points to remember about Symmetry

- If a curve is symmetric about x -axis and y -axis, then it is symmetric about the pole.



Points to remember about Symmetry

- If a curve is symmetric about x -axis and y -axis, then it is symmetric about the pole.
- If a curve is symmetric about x -axis & pole, then it is symmetric about y -axis.



Points to remember about Symmetry

- If a curve is symmetric about x -axis and y -axis, then it is symmetric about the pole.
- If a curve is symmetric about x -axis & pole, then it is symmetric about y -axis.
- If a curve is symmetric about y -axis & pole, then it is symmetric about x -axis.



Points to remember about Symmetry

- If a curve is symmetric about x -axis and y -axis, then it is symmetric about the pole.
- If a curve is symmetric about x -axis & pole, then it is symmetric about y -axis.
- If a curve is symmetric about y -axis & pole, then it is symmetric about x -axis.
- Thus if a curve is symmetric about x -axis but not symmetric about y -axis (or if a curve is symmetric about y -axis but not symmetric about x -axis) then it can not be symmetric about the pole.



Slope of a Polar Curve



Slope of a Polar Curve

- The parametric equations of $r = f(\theta)$ are $x = r \cos \theta$, $y = r \sin \theta$.



Slope of a Polar Curve

- The parametric equations of $r = f(\theta)$ are $x = r \cos \theta$, $y = r \sin \theta$.
- The slope of the curve $r = f(\theta)$ at any point (r, θ) is given by

$$\left. \frac{dy}{dx} \right|_{(r, \theta)} = \frac{\left. \frac{dy}{d\theta} \right|_{(r, \theta)}}{\left. \frac{dx}{d\theta} \right|_{(r, \theta)}} = \frac{f'(\theta) \sin \theta + f(\theta) \cos \theta}{f'(\theta) \cos \theta - f(\theta) \sin \theta}$$

provided $\frac{dx}{d\theta} \neq 0$ at any point (r, θ) .



Example

Find the slope of the curve $r = 4 \sin 3\theta$ at $\theta = \frac{\pi}{6}$.



Example

Find the slope of the curve $r = 4 \sin 3\theta$ at $\theta = \frac{\pi}{6}$.

Sol.

$$\begin{aligned}\left. \frac{dy}{dx} \right|_{(r,\theta)} &= \frac{f'(\theta) \sin \theta + f(\theta) \cos \theta}{f'(\theta) \cos \theta - f(\theta) \sin \theta} \\ &= \frac{12 \cos 3\theta \sin \theta + 4 \sin 3\theta \cos \theta}{12 \cos 3\theta \cos \theta - 4 \sin 3\theta \sin \theta}.\end{aligned}$$



Example

Find the slope of the curve $r = 4 \sin 3\theta$ at $\theta = \frac{\pi}{6}$.

Sol.

$$\begin{aligned}\left. \frac{dy}{dx} \right|_{(r,\theta)} &= \frac{f'(\theta) \sin \theta + f(\theta) \cos \theta}{f'(\theta) \cos \theta - f(\theta) \sin \theta} \\ &= \frac{12 \cos 3\theta \sin \theta + 4 \sin 3\theta \cos \theta}{12 \cos 3\theta \cos \theta - 4 \sin 3\theta \sin \theta}.\end{aligned}$$

Thus

$$\left. \frac{dy}{dx} \right|_{\theta=\pi/6} = -\sqrt{3}.$$



Q:.. What does the slope of a curve represent.



Q: What does the slope of a curve represent.

$$\left. \frac{dy}{dx} \right|_{\theta=\pi/6} = -\sqrt{3}.$$



Q:. What does the slope of a curve represent.

$$\left. \frac{dy}{dx} \right|_{\theta=\pi/6} = -\sqrt{3}.$$

At $\theta = \frac{\pi}{6}$, $r = 4$ and $\tan \theta_1 = -\sqrt{3} \implies \theta_1 = \frac{2\pi}{3}$.




Q:. What does the slope of a curve represent.

$$\left. \frac{dy}{dx} \right|_{\theta=\pi/6} = -\sqrt{3}.$$

At $\theta = \frac{\pi}{6}$, $r = 4$ and $\tan \theta_1 = -\sqrt{3} \implies \theta_1 = \frac{2\pi}{3}$. Here θ_1 is the angle the tangent at $(r, \theta) = (4, \frac{\pi}{6})$ makes with the x -axis.





Slope at Pole

How to compute Slope at Pole?



Slope at Pole

How to compute Slope at Pole?

If the curve $r = f(\theta)$ passes through the pole at $\theta = \theta_0$, then $r = f(\theta_0) = 0$.



Slope at Pole

How to compute Slope at Pole?

If the curve $r = f(\theta)$ passes through the pole at $\theta = \theta_0$, then $r = f(\theta_0) = 0$. Hence the slope of the curve $r = f(\theta)$ at pole is given by

$$\left. \frac{dy}{dx} \right|_{(0, \theta_0)} = \tan \theta_0$$

.



Slope at Pole

How to compute Slope at Pole?

If the curve $r = f(\theta)$ passes through the pole at $\theta = \theta_0$, then $r = f(\theta_0) = 0$. Hence the slope of the curve $r = f(\theta)$ at pole is given by

$$\left. \frac{dy}{dx} \right|_{(0, \theta_0)} = \tan \theta_0$$

.

Hence the line $\theta = \theta_0$ is the tangent to the curve at the pole.



Q: Find the slope of the curve $r = 1 + \cos \theta$ at $\theta = \frac{\pi}{2}$.



Q:. Find the slope of the curve $r = 1 + \cos \theta$ at $\theta = \frac{\pi}{2}$.

Sol.

$$\left. \frac{dy}{dx} \right|_{\theta=\pi/2} = 1.$$



Tracing the Curve in Polar Coordinate

Q.: Trace the curve $r = 1 + \cos \theta$.



Tracing the Curve in Polar Coordinate

Q:. Trace the curve $r = 1 + \cos \theta$.

Step 1. Check for symmetries (it will reduce the work for tracing).



Tracing the Curve in Polar Coordinate

Q:. Trace the curve $r = 1 + \cos \theta$.

Step 1. Check for symmetries (it will reduce the work for tracing).

- Since $(r, -\theta)$ lies on the curve, it is symmetric about x -axis.



Tracing the Curve in Polar Coordinate

Q:. Trace the curve $r = 1 + \cos \theta$.

Step 1. Check for symmetries (it will reduce the work for tracing).

- Since $(r, -\theta)$ lies on the curve, it is symmetric about x -axis. Hence, it is enough to consider the steps for $0 \leq \theta \leq \pi$.



Step 2. Solve the equation $r = 0$ for θ .



Step 2. Solve the equation $r = 0$ for θ . If $\theta = \theta_0$ satisfies $r = 0$, then the line $\theta = \theta_0$ will be a tangent to the curve at pole.



Step 2. Solve the equation $r = 0$ for θ . If $\theta = \theta_0$ satisfies $r = 0$, then the line $\theta = \theta_0$ will be a tangent to the curve at pole.

□ $r = 0$ gives $\cos \theta = -1 \implies \theta = \pi$. Thus $\theta = \pi$ is a tangent to the curve at pole.



Step 3 Find $\frac{dr}{d\theta}$.



Step 3 Find $\frac{dr}{d\theta}$.

$$\square \frac{dr}{d\theta} = -\sin \theta.$$



Step 3 Find $\frac{dr}{d\theta}$.

$$\square \frac{dr}{d\theta} = -\sin \theta.$$

Step 3.1 Find the values of θ for which $\frac{dr}{d\theta} > 0$.



Step 3 Find $\frac{dr}{d\theta}$.

$$\square \frac{dr}{d\theta} = -\sin \theta.$$

Step 3.1 Find the values of θ for which $\frac{dr}{d\theta} > 0$. This would be an interval (or no value of θ). In this interval r is an increasing function.



Step 3 Find $\frac{dr}{d\theta}$.

$$\square \frac{dr}{d\theta} = -\sin \theta.$$

Step 3.1 Find the values of θ for which $\frac{dr}{d\theta} > 0$. This would be an interval (or no value of θ). In this interval r is an increasing function. Find max/min values of r and the associated θ values in this interval.



Step 3 Find $\frac{dr}{d\theta}$.

$$\square \frac{dr}{d\theta} = -\sin \theta.$$

Step 3.1 Find the values of θ for which $\frac{dr}{d\theta} > 0$. This would be an interval (or no value of θ). In this interval r is an increasing function. Find max/min values of r and the associated θ values in this interval.

$$\square \frac{dr}{d\theta} > 0 \Rightarrow \sin \theta < 0,$$



Step 3 Find $\frac{dr}{d\theta}$.

$$\square \frac{dr}{d\theta} = -\sin \theta.$$

Step 3.1 Find the values of θ for which $\frac{dr}{d\theta} > 0$. This would be an interval (or no value of θ). In this interval r is an increasing function. Find max/min values of r and the associated θ values in this interval.

$$\square \frac{dr}{d\theta} > 0 \Rightarrow \sin \theta < 0, \text{ thus no value of } \theta \text{ in between } 0 \text{ and } \pi.$$



Step 3.2 Similarly find the values of θ for which $\frac{dr}{d\theta} < 0$. This would be an interval (or no value of θ). In this interval r is a decreasing function. Find max/min values of r and the associated θ values in this interval.



Step 3.2 Similarly find the values of θ for which $\frac{dr}{d\theta} < 0$. This would be an interval (or no value of θ). In this interval r is a decreasing function. Find max/min values of r and the associated θ values in this interval.

- $\frac{dr}{d\theta} < 0 \Rightarrow \sin \theta > 0 \Rightarrow 0 < \theta < \pi$, thus r decreases in the interval $[0, \pi]$.



Step 3.2 Similarly find the values of θ for which $\frac{dr}{d\theta} < 0$. This would be an interval (or no value of θ). In this interval r is a decreasing function. Find max/min values of r and the associated θ values in this interval.

- $\frac{dr}{d\theta} < 0 \Rightarrow \sin \theta > 0 \Rightarrow 0 < \theta < \pi$, thus r decreases in the interval $[0, \pi]$.

Clearly $\max r = 2$ at $\theta = 0$ and $\min r = 0$ at $\theta = \pi$.



Step 4. Find slopes at some special points (like at end points of the intervals obtained in step 3).



Step 4. Find slopes at some special points (like at end points of the intervals obtained in step 3).

$$\left. \frac{dy}{dx} \right|_{\theta=0} = \infty, \quad \left. \frac{dy}{dx} \right|_{\theta=\frac{\pi}{2}} = 1.$$



Step 4. Find slopes at some special points (like at end points of the intervals obtained in step 3).

$$\left. \frac{dy}{dx} \right|_{\theta=0} = \infty, \quad \left. \frac{dy}{dx} \right|_{\theta=\frac{\pi}{2}} = 1.$$

At $\theta = 0$, $r = 2$ and $\tan \theta_1 = \infty \implies \theta_1 = \frac{\pi}{2}$.



Step 4. Find slopes at some special points (like at end points of the intervals obtained in step 3).

$$\left. \frac{dy}{dx} \right|_{\theta=0} = \infty, \quad \left. \frac{dy}{dx} \right|_{\theta=\frac{\pi}{2}} = 1.$$

At $\theta = 0$, $r = 2$ and $\tan \theta_1 = \infty \implies \theta_1 = \frac{\pi}{2}$.

At $\theta = \frac{\pi}{2}$, $r = 1$ and $\tan \theta_1 = 1 \implies \theta_1 = \frac{\pi}{4}$.



Step 5. Make a table θ vs r for different values of θ .



Step 5. Make a table θ vs r for different values of θ .

θ	0	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	π
r	2	$1 + \frac{1}{\sqrt{2}}$	$\frac{3}{2}$	1	$\frac{1}{2}$	$1 - \frac{1}{\sqrt{2}}$	0



Step 5. Make a table θ vs r for different values of θ .

θ	0	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	π
r	2	$1 + \frac{1}{\sqrt{2}}$	$\frac{3}{2}$	1	$\frac{1}{2}$	$1 - \frac{1}{\sqrt{2}}$	0

Step 6. Plot the curve while considering the above steps.

