

MATHEMATICS-I (MATH F111)

Dr. Krishnendra Shekhawat

BITS PILANI
Department of Mathematics



CHAPTER 10

Infinite Sequences and Series



Topics to be covered in Chapter 10



Topics to be covered in Chapter 10

- Sequences (Certain Theorems on Sequences)



Topics to be covered in Chapter 10

- Sequences (Certain Theorems on Sequences)
- Infinite Series



Topics to be covered in Chapter 10

- Sequences (Certain Theorems on Sequences)
- Infinite Series
- Integral Test



Topics to be covered in Chapter 10

- Sequences (Certain Theorems on Sequences)
- Infinite Series
- Integral Test
- Comparison Tests



Topics to be covered in Chapter 10

- Sequences (Certain Theorems on Sequences)
- Infinite Series
- Integral Test
- Comparison Tests
- Ratio and Root Tests



Topics to be covered in Chapter 10

- Sequences (Certain Theorems on Sequences)
- Infinite Series
- Integral Test
- Comparison Tests
- Ratio and Root Tests
- Alternating Series



Topics to be covered in Chapter 10

- Sequences (Certain Theorems on Sequences)
- Infinite Series
- Integral Test
- Comparison Tests
- Ratio and Root Tests
- Alternating Series
- Power Series



Topics to be covered in Chapter 10

- Sequences (Certain Theorems on Sequences)
- Infinite Series
- Integral Test
- Comparison Tests
- Ratio and Root Tests
- Alternating Series
- Power Series
- Taylor & Maclaurin Series



Section 10.1

Sequences [Self Study]



Sequence

A function whose domain is the set of natural numbers ($f : \mathbb{N} \rightarrow \mathbb{R}$) is called a **sequence of real numbers**.

We write



Sequence

A function whose domain is the set of natural numbers ($f : \mathbb{N} \rightarrow \mathbb{R}$) is called a **sequence of real numbers**.

We write $f(n) = a_n$, and the sequence is denoted by



Sequence

A function whose domain is the set of natural numbers ($f : \mathbb{N} \rightarrow \mathbb{R}$) is called a **sequence of real numbers**.

We write $f(n) = a_n$, and the sequence is denoted by $\{a_1, a_2, \dots\}$ or by $\{a_n\}_{n=1}^{\infty}$ or simply by $\{a_n\}$.



Sequence

A function whose domain is the set of natural numbers ($f : \mathbb{N} \rightarrow \mathbb{R}$) is called a **sequence of real numbers**.

We write $f(n) = a_n$, and the sequence is denoted by $\{a_1, a_2, \dots\}$ or by $\{a_n\}_{n=1}^{\infty}$ or simply by $\{a_n\}$.

We call a_n the n^{th} term of the sequence or the value of the sequence at n .



Sequence

A function whose domain is the set of natural numbers ($f : \mathbb{N} \rightarrow \mathbb{R}$) is called a **sequence of real numbers**.

We write $f(n) = a_n$, and the sequence is denoted by $\{a_1, a_2, \dots\}$ or by $\{a_n\}_{n=1}^{\infty}$ or simply by $\{a_n\}$.

We call a_n the n^{th} term of the sequence or the value of the sequence at n .

Note. Sequence is a special case for Infinite series.



Examples: Sequences

- $\{n\} = \{1, 2, 3, \dots, n, \dots\}$



Examples: Sequences

- $\{n\} = \{1, 2, 3, \dots, n, \dots\}$
- $\left\{\frac{1}{n}\right\} = \left\{1, \frac{1}{2}, \frac{1}{3}, \dots, \frac{1}{n}, \dots\right\}$



Examples: Sequences

- $\{n\} = \{1, 2, 3, \dots, n, \dots\}$
- $\left\{\frac{1}{n}\right\} = \left\{1, \frac{1}{2}, \frac{1}{3}, \dots, \frac{1}{n}, \dots\right\}$
- $\{(-1)^{n+1}\} = \{1, -1, 1, -1, \dots, (-1)^{n+1}, \dots\}$



Examples: Sequences

- $\{n\} = \{1, 2, 3, \dots, n, \dots\}$
- $\left\{\frac{1}{n}\right\} = \left\{1, \frac{1}{2}, \frac{1}{3}, \dots, \frac{1}{n}, \dots\right\}$
- $\{(-1)^{n+1}\} = \{1, -1, 1, -1, \dots, (-1)^{n+1}, \dots\}$
- $\left\{1 - \frac{1}{n}\right\} = \left\{0, \frac{1}{2}, \frac{2}{3}, \dots, 1 - \frac{1}{n}, \dots\right\}$



Convergence

A sequence $\{a_n\}$ is said to converge to a number L ,



Convergence

A sequence $\{a_n\}$ is said to **converge to a number L** , if for every positive number ϵ , however small, we can find a positive integer N (depending on ϵ) such that

$$|a_n - L| < \epsilon, \quad \forall n > N.$$



- If no such number L exists, the sequence is said to



- If no such number L exists, the sequence is said to **diverge**.



- If no such number L exists, the sequence is said to **diverge**.
- If $\{a_n\}$ converges to L , then we write



- If no such number L exists, the sequence is said to **diverge**.
- If $\{a_n\}$ converges to L , then we write $\lim_{n \rightarrow \infty} a_n = L$ or simply $a_n \rightarrow L$.



- If no such number L exists, the sequence is said to **diverge**.
- If $\{a_n\}$ converges to L , then we write $\lim_{n \rightarrow \infty} a_n = L$ or simply $a_n \rightarrow L$.
- The number L is called **limit** of the sequence.



- If no such number L exists, the sequence is said to **diverge**.
- If $\{a_n\}$ converges to L , then we write $\lim_{n \rightarrow \infty} a_n = L$ or simply $a_n \rightarrow L$.
- The number L is called **limit** of the sequence.

Note. A sequence **can not converges to more than one limit** *i.e.*, the limit of a sequence is unique.



Examples: Converge or Diverge ?

- The sequence $\left\{\frac{1}{n}\right\}$ converges to



Examples: Converge or Diverge ?

- The sequence $\left\{\frac{1}{n}\right\}$ converges to the number 0



Examples: Converge or Diverge ?

- The sequence $\left\{\frac{1}{n}\right\}$ converges to the number 0
- The sequence $\{n\}$



Examples: Converge or Diverge ?

- The sequence $\left\{\frac{1}{n}\right\}$ converges to the number 0
- The sequence $\{n\}$ diverges.



Examples: Converge or Diverge ?

- The sequence $\left\{\frac{1}{n}\right\}$ converges to the number 0
- The sequence $\{n\}$ diverges.
- The sequence $\left\{\frac{(-1)^n}{n}\right\}$ converges to



Examples: Converge or Diverge ?

- The sequence $\left\{\frac{1}{n}\right\}$ converges to the number 0
- The sequence $\{n\}$ diverges.
- The sequence $\left\{\frac{(-1)^n}{n}\right\}$ converges to the number 0



Examples: Converge or Diverge ?

- The sequence $\left\{\frac{1}{n}\right\}$ converges to the number 0
- The sequence $\{n\}$ diverges.
- The sequence $\left\{\frac{(-1)^n}{n}\right\}$ converges to the number 0
- The sequence $\{(-1)^n\}$



Examples: Converge or Diverge ?

- The sequence $\left\{\frac{1}{n}\right\}$ converges to the number 0
- The sequence $\{n\}$ diverges.
- The sequence $\left\{\frac{(-1)^n}{n}\right\}$ converges to the number 0
- The sequence $\{(-1)^n\}$ diverges.



Evaluate $\lim \frac{5n+7}{3n-5}$



Evaluate $\lim_{n \rightarrow \infty} \frac{5n+7}{3n-5}$ (divide numerator and denominator by n)



Evaluate $\lim_{n \rightarrow \infty} \frac{5n+7}{3n-5}$ (divide numerator and denominator by n)

Theorem (Theorem 1.)

Let $\{a_n\}$ and $\{b_n\}$ be two convergent sequences. Then

- ① $\lim ka_n = k \lim a_n$
- ② $\lim(a_n \pm b_n) = \lim a_n \pm \lim b_n$
- ③ $\lim(a_n \cdot b_n) = \lim a_n \cdot \lim b_n$
- ④ $\lim\left(\frac{a_n}{b_n}\right) = \frac{\lim a_n}{\lim b_n}$, if $b_n \neq 0$ for all n and $\lim b_n \neq 0$.



Find $\lim_{n \rightarrow \infty} \frac{\cos n}{n}$ (If exists).



Find $\lim \frac{\cos n}{n}$ (If exists).

Theorem (Theorem 2. Sandwich Theorem or Squeeze Theorem)

Let $\{a_n\}$, $\{b_n\}$, and $\{c_n\}$ be sequences of real numbers.



Find $\lim \frac{\cos n}{n}$ (If exists).

Theorem (Theorem 2. Sandwich Theorem or Squeeze Theorem)

Let $\{a_n\}$, $\{b_n\}$, and $\{c_n\}$ be sequences of real numbers. If $a_n \leq b_n \leq c_n$ for all $n \geq N$ and if $\lim a_n = \lim c_n = L$, then $\lim b_n = L$.



Find $\lim \frac{\cos n}{n}$ (If exists).

Theorem (Theorem 2. Sandwich Theorem or Squeeze Theorem)

Let $\{a_n\}$, $\{b_n\}$, and $\{c_n\}$ be sequences of real numbers. If $a_n \leq b_n \leq c_n$ for all $n \geq N$ and if $\lim a_n = \lim c_n = L$, then $\lim b_n = L$.

Example. Find $\lim \frac{\cos n}{n}$.



Find $\lim \frac{\cos n}{n}$ (If exists).

Theorem (Theorem 2. Sandwich Theorem or Squeeze Theorem)

Let $\{a_n\}$, $\{b_n\}$, and $\{c_n\}$ be sequences of real numbers. If $a_n \leq b_n \leq c_n$ for all $n \geq N$ and if $\lim a_n = \lim c_n = L$, then $\lim b_n = L$.

Example. Find $\lim \frac{\cos n}{n}$.

Sol. $\frac{-1}{n} \leq \frac{\cos n}{n} \leq \frac{1}{n}$.



Find $\lim \frac{\cos n}{n}$ (If exists).

Theorem (Theorem 2. Sandwich Theorem or Squeeze Theorem)

Let $\{a_n\}$, $\{b_n\}$, and $\{c_n\}$ be sequences of real numbers. If $a_n \leq b_n \leq c_n$ for all $n \geq N$ and if $\lim a_n = \lim c_n = L$, then $\lim b_n = L$.

Example. Find $\lim \frac{\cos n}{n}$.

Sol. $\frac{-1}{n} \leq \frac{\cos n}{n} \leq \frac{1}{n}$. Now, $\lim \frac{1}{n} = \lim \frac{-1}{n} = 0$, hence, using Sandwich theorem we have $\lim \frac{\cos n}{n} = 0$.



Find $\lim \cos \frac{1}{n}$.



Find $\lim \cos \frac{1}{n}$.

Theorem (Theorem 3. Continuous Function Theorem)

Let $\{a_n\}$ be a sequence of real numbers.



Find $\lim \cos \frac{1}{n}$.

Theorem (Theorem 3. Continuous Function Theorem)

Let $\{a_n\}$ be a sequence of real numbers. If $a_n \rightarrow L$ and $f(x)$ is a function that is continuous at $x = L$



Find $\lim \cos \frac{1}{n}$.

Theorem (Theorem 3. Continuous Function Theorem)

Let $\{a_n\}$ be a sequence of real numbers. If $a_n \rightarrow L$ and $f(x)$ is a function that is continuous at $x = L$ and defined at all a_n ,



Find $\lim \cos \frac{1}{n}$.

Theorem (Theorem 3. Continuous Function Theorem)

Let $\{a_n\}$ be a sequence of real numbers. If $a_n \rightarrow L$ and $f(x)$ is a function that is continuous at $x = L$ and defined at all a_n , then $f(a_n) \rightarrow f(L)$.



Example

Find $\lim \cos \frac{1}{n}$ (If exists).



Example

Find $\lim \cos \frac{1}{n}$ (If exists).

Sol. Let $a_n = \frac{1}{n}$. Then $\lim a_n = \lim \frac{1}{n} = 0$. Since $\cos x$ is continuous at $x = 0$ and defined at $a_n = \frac{1}{n}$ for all n , therefore

$$\lim \cos \frac{1}{n} = \cos(0) = 1.$$



Q: Find $\lim \frac{n+1}{e^n}$.



Q: Find $\lim \frac{n+1}{e^n}$.

Q: Can we use L'Hôpital's rule ?



Q.: Find $\lim \frac{n+1}{e^n}$.

Q.: Can we use L'Hôpital's rule ?

Theorem (Theorem 4.)

Suppose $f(x)$ is a function defined for all $x \geq n_0$ and that $\{a_n\}$ is a sequence of real numbers s.t. $a_n = f(n)$ for $n \geq n_0$, then

$$\lim_{x \rightarrow \infty} f(x) = L \Rightarrow \lim_{n \rightarrow \infty} f(n) = L.$$



Q: Find $\lim \frac{n+1}{e^n}$.



Q:. Find $\lim \frac{n+1}{e^n}$.

Sol.

$$\begin{aligned}\lim \frac{x+1}{e^x} &= \lim \frac{1}{e^x}, \quad \text{by L'Hôpital's rule} \\ &= 0.\end{aligned}$$



Q:. Find $\lim \frac{n+1}{e^n}$.

Sol.

$$\begin{aligned}\lim \frac{x+1}{e^x} &= \lim \frac{1}{e^x}, \quad \text{by L'Hôpital's rule} \\ &= 0.\end{aligned}$$

Hence $\lim \frac{n+1}{e^n} = 0$.



Q:. Find $\lim \frac{n+1}{e^n}$.

Sol.

$$\begin{aligned}\lim \frac{x+1}{e^x} &= \lim \frac{1}{e^x}, \quad \text{by L'Hôpital's rule} \\ &= 0.\end{aligned}$$

Hence $\lim \frac{n+1}{e^n} = 0$.

Q:. Find $\lim \cos(2\pi x)$?



Q:. Find $\lim \frac{n+1}{e^n}$.

Sol.

$$\lim \frac{x+1}{e^x} = \lim \frac{1}{e^x}, \quad \text{by L'Hôpital's rule} \\ = 0.$$

Hence $\lim \frac{n+1}{e^n} = 0$.

Q:. Find $\lim \cos(2\pi x)$? (does not exist)



Q:. Find $\lim \frac{n+1}{e^n}$.

Sol.

$$\lim \frac{x+1}{e^x} = \lim \frac{1}{e^x}, \quad \text{by L'Hôpital's rule} \\ = 0.$$

Hence $\lim \frac{n+1}{e^n} = 0$.

Q:. Find $\lim \cos(2\pi x)$? (does not exist)

Q:. Find $\lim \cos(2\pi n)$?



Q:. Find $\lim \frac{n+1}{e^n}$.

Sol.

$$\lim \frac{x+1}{e^x} = \lim \frac{1}{e^x}, \quad \text{by L'Hôpital's rule} \\ = 0.$$

Hence $\lim \frac{n+1}{e^n} = 0$.

Q:. Find $\lim \cos(2\pi x)$? (does not exist)

Q:. Find $\lim \cos(2\pi n)$? ($=1$)



Q:. Find $\lim \frac{n+1}{e^n}$.

Sol.

$$\lim \frac{x+1}{e^x} = \lim \frac{1}{e^x}, \quad \text{by L'Hôpital's rule} \\ = 0.$$

Hence $\lim \frac{n+1}{e^n} = 0$.

Q:. Find $\lim \cos(2\pi x)$? (does not exist)

Q:. Find $\lim \cos(2\pi n)$? (=1)

Conclusion?



Q:. Find $\lim \frac{n+1}{e^n}$.

Sol.

$$\lim \frac{x+1}{e^x} = \lim \frac{1}{e^x}, \quad \text{by L'Hôpital's rule} \\ = 0.$$

Hence $\lim \frac{n+1}{e^n} = 0$.

Q:. Find $\lim \cos(2\pi x)$? (does not exist)

Q:. Find $\lim \cos(2\pi n)$? (=1)

Conclusion? Converse of Theorem 4 is not true.



Bounded Sequence

A sequence $\{a_n\}$ is said to be **bounded from above** if there exists a number M_1 such that

$$a_n \leq M_1, \forall n.$$

The number M_1 is called an upper bound for $\{a_n\}$.



Bounded Sequence

A sequence $\{a_n\}$ is said to be **bounded from above** if there exists a number M_1 such that

$$a_n \leq M_1, \forall n.$$

The number M_1 is called an upper bound for $\{a_n\}$.

Similarly, a sequence $\{a_n\}$ is said to be **bounded from below** if there exists a number M_2 such that

$$a_n \geq M_2, \forall n.$$

The number M_2 is called a lower bound for $\{a_n\}$.



Examples: Bounded or not ?

- The sequence $\left\{\frac{1}{n}\right\}$ is bounded below by



Examples: Bounded or not ?

- The sequence $\left\{\frac{1}{n}\right\}$ is bounded below by 0 and bounded above by



Examples: Bounded or not ?

- The sequence $\left\{\frac{1}{n}\right\}$ is bounded below by 0 and bounded above by 1.



Examples: Bounded or not ?

- The sequence $\left\{\frac{1}{n}\right\}$ is bounded below by 0 and bounded above by 1.
- The sequence $\left\{\frac{1}{2^n}\right\}$ is bounded below by



Examples: Bounded or not ?

- The sequence $\left\{\frac{1}{n}\right\}$ is bounded below by 0 and bounded above by 1.
- The sequence $\left\{\frac{1}{2^n}\right\}$ is bounded below by 0



Examples: Bounded or not ?

- The sequence $\left\{\frac{1}{n}\right\}$ is bounded below by 0 and bounded above by 1.
- The sequence $\left\{\frac{1}{2^n}\right\}$ is bounded below by 0 and bounded above by



Examples: Bounded or not ?

- The sequence $\left\{\frac{1}{n}\right\}$ is bounded below by 0 and bounded above by 1.
- The sequence $\left\{\frac{1}{2^n}\right\}$ is bounded below by 0 and bounded above by $\frac{1}{2}$.



Examples: Bounded or not ?

- The sequence $\left\{\frac{1}{n}\right\}$ is bounded below by 0 and bounded above by 1.
- The sequence $\left\{\frac{1}{2^n}\right\}$ is bounded below by 0 and bounded above by $\frac{1}{2}$.

Bounded Sequence

A sequence $\{a_n\}$ is said to be bounded if there exists a number M such that

$$|a_n| \leq M, \forall n.$$

The number M is called a bound for $\{a_n\}$.



Nondecreasing/Nonincreasing/Monotonic Sequence

A sequence $\{a_n\}$ is called a **nondecreasing sequence**, if $a_n \leq a_{n+1}$ for all n . The sequence is **nonincreasing**, if $a_n \geq a_{n+1}$ for all n . The sequence is **monotonic** if it is either nondecreasing or nonincreasing.



Nondecreasing/Nonincreasing/Monotonic Sequence

A sequence $\{a_n\}$ is called a **nondecreasing sequence**, if $a_n \leq a_{n+1}$ for all n . The sequence is **nonincreasing**, if $a_n \geq a_{n+1}$ for all n . The sequence is **monotonic** if it is either nondecreasing or nonincreasing.

Examples: Nonincreasing or Nondecreasing ?

- The sequence $\left\{\frac{n}{n+1}\right\}$ is



Nondecreasing/Nonincreasing/Monotonic Sequence

A sequence $\{a_n\}$ is called a **nondecreasing sequence**, if $a_n \leq a_{n+1}$ for all n . The sequence is **nonincreasing**, if $a_n \geq a_{n+1}$ for all n . The sequence is **monotonic** if it is either nondecreasing or nonincreasing.

Examples: Nonincreasing or Nondecreasing ?

- The sequence $\left\{\frac{n}{n+1}\right\}$ is nondecreasing.



Nondecreasing/Nonincreasing/Monotonic Sequence

A sequence $\{a_n\}$ is called a **nondecreasing sequence**, if $a_n \leq a_{n+1}$ for all n . The sequence is **nonincreasing**, if $a_n \geq a_{n+1}$ for all n . The sequence is **monotonic** if it is either nondecreasing or nonincreasing.

Examples: Nonincreasing or Nondecreasing ?

- The sequence $\left\{\frac{n}{n+1}\right\}$ is nondecreasing.
- The sequence $\left\{\frac{1}{n}\right\}$ is



Nondecreasing/Nonincreasing/Monotonic Sequence

A sequence $\{a_n\}$ is called a **nondecreasing sequence**, if $a_n \leq a_{n+1}$ for all n . The sequence is **nonincreasing**, if $a_n \geq a_{n+1}$ for all n . The sequence is **monotonic** if it is either nondecreasing or nonincreasing.

Examples: Nonincreasing or Nondecreasing ?

- The sequence $\left\{\frac{n}{n+1}\right\}$ is nondecreasing.
- The sequence $\left\{\frac{1}{n}\right\}$ is nonincreasing.



Theorem (Theorem 6. Monotonic Sequence Theorem)

A bounded monotonic sequence is convergent.



Theorem (Theorem 6. Monotonic Sequence Theorem)

A bounded monotonic sequence is convergent.

Example

The sequence $\left\{\frac{1}{n}\right\}$ is monotonic (nonincreasing) and bounded (as $|a_n| \leq 1$), therefore it is convergent.



Section 10.2

Infinite Series



Infinite Series

An **infinite series** is the sum of the infinite terms of a sequence, i.e., it consists of infinite number of real numbers separated by $+$ sign.



Infinite Series

An **infinite series** is the sum of the infinite terms of a sequence, i.e., it consists of infinite number of real numbers separated by $+$ sign. For a sequence $\{a_n\}$ the series is written as



Infinite Series

An **infinite series** is the sum of the infinite terms of a sequence, i.e., it consists of infinite number of real numbers separated by $+$ sign. For a sequence $\{a_n\}$ the series is written as $a_1 + a_2 + a_3 + \cdots + a_n + \cdots$ or as $\sum_{n=1}^{\infty} a_n$ or simply as $\sum a_n$.



Infinite Series

An **infinite series** is the sum of the infinite terms of a sequence, i.e., it consists of infinite number of real numbers separated by $+$ sign. For a sequence $\{a_n\}$ the series is written as $a_1 + a_2 + a_3 + \cdots + a_n + \cdots$ or as $\sum_{n=1}^{\infty} a_n$ or simply as $\sum a_n$. The number a_n is called the n^{th} term of the series.



Infinite Series

An **infinite series** is the sum of the infinite terms of a sequence, i.e., it consists of infinite number of real numbers separated by $+$ sign. For a sequence $\{a_n\}$ the series is written as $a_1 + a_2 + a_3 + \cdots + a_n + \cdots$ or as $\sum_{n=1}^{\infty} a_n$ or simply as $\sum a_n$. The number a_n is called the n^{th} term of the series.

Example: $1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n} + \cdots$ (Harmonic Series)





First method: Sequence of Partial Sums

Consider the series $\sum a_n$.



First method: Sequence of Partial Sums

Consider the series $\sum a_n$. The sequence $\{S_n\}$, defined by

$$S_n = a_1 + a_2 + a_3 + \cdots + a_n, \quad \forall n$$

is called the **sequence of partial sums** of the series.



First method: Sequence of Partial Sums

Consider the series $\sum a_n$. The sequence $\{S_n\}$, defined by

$$S_n = a_1 + a_2 + a_3 + \cdots + a_n, \quad \forall n$$

is called the **sequence of partial sums** of the series.

The n^{th} term S_n of the sequence is called **n^{th} partial sum** of the series.



Convergent Series

A series $\sum a_n$ is said to be **convergent** iff the sequence $\{S_n\}$ is convergent.



Convergent Series

A series $\sum a_n$ is said to be **convergent** iff the sequence $\{S_n\}$ is convergent. If $\{S_n\}$ converges to a real number L , then the series $\sum a_n$ also converges to L



Convergent Series

A series $\sum a_n$ is said to be **convergent** iff the sequence $\{S_n\}$ is convergent. If $\{S_n\}$ converges to a real number L , then the series $\sum a_n$ also converges to L and the sum of the series is L .



Convergent Series

A series $\sum a_n$ is said to be **convergent** iff the sequence $\{S_n\}$ is convergent. If $\{S_n\}$ converges to a real number L , then the series $\sum a_n$ also converges to L and the sum of the series is L .

A series which is not convergent is called **divergent**.



Convergent Series

Q:. What is the effect of addition or deletion of a finite number of terms in a series on the convergence of a series.



Convergent Series

Q:. What is the effect of addition or deletion of a finite number of terms in a series on the convergence of a series.

Sol. It does not alter the behavior (convergence or divergence) of the series. However, the sum of the series will change in the case of convergent series.



Q.: $\sum \frac{1}{(n+4)(n+5)}$



Q.: $\sum \frac{1}{(n+4)(n+5)}$

Example: Telescoping series

Q.: Find a formula for the n^{th} partial sum of $\sum \frac{1}{(n+4)(n+5)}$ and use it to determine the convergence of the series. If the series converges, find its sum.



Q.: $\sum \frac{1}{(n+4)(n+5)}$

Example: Telescoping series

Q.: Find a formula for the n^{th} partial sum of $\sum \frac{1}{(n+4)(n+5)}$ and use it to determine the convergence of the series. If the series converges, find its sum.

Sol. Here $a_n = \frac{1}{(n+4)(n+5)} = \frac{1}{n+4} - \frac{1}{n+5}$. Now,



$$\text{Q.: } \sum \frac{1}{(n+4)(n+5)}$$

Example: Telescoping series

Q.: Find a formula for the n^{th} partial sum of $\sum \frac{1}{(n+4)(n+5)}$ and use it to determine the convergence of the series. If the series converges, find its sum.

Sol. Here $a_n = \frac{1}{(n+4)(n+5)} = \frac{1}{n+4} - \frac{1}{n+5}$. Now,

$$S_n = \left(\frac{1}{5} - \frac{1}{6}\right) + \left(\frac{1}{6} - \frac{1}{7}\right) + \cdots + \left(\frac{1}{n+4} - \frac{1}{n+5}\right) = \frac{1}{5} - \frac{1}{n+5}.$$

Thus $\lim S_n = \frac{1}{5}$ and hence series converges and $\sum \frac{1}{(n+4)(n+5)} = \frac{1}{5}$.



Telescoping series

Because of the manner in which the general term of the sequence of partial sums **collapses** to two terms, the series in above Example is said to be a **telescoping series**.



Telescoping series

Because of the manner in which the general term of the sequence of partial sums **collapses** to two terms, the series in above Example is said to be a **telescoping series**.

This leads us to many more well-known series.



Second method: Comparison with well-known series

Two most commonly used series are as follows:



Second method: Comparison with well-known series

Two most commonly used series are as follows:

- ① The geometric series: $\sum ar^{n-1}$, $a \neq 0$



Second method: Comparison with well-known series

Two most commonly used series are as follows:

- ① The geometric series: $\sum ar^{n-1}$, $a \neq 0$ (e.g. $\sum 5\left(\frac{3}{2}\right)^{n-1}$)



Second method: Comparison with well-known series

Two most commonly used series are as follows:

- ① The geometric series: $\sum ar^{n-1}$, $a \neq 0$ (e.g. $\sum 5\left(\frac{3}{2}\right)^{n-1}$)
- ② The p -series: $\sum \frac{1}{n^p}$, $p \in \mathbb{R}$



Second method: Comparison with well-known series

Two most commonly used series are as follows:

- ① The geometric series: $\sum ar^{n-1}$, $a \neq 0$ (e.g. $\sum 5\left(\frac{3}{2}\right)^{n-1}$)
- ② The p -series: $\sum \frac{1}{n^p}$, $p \in \mathbb{R}$ (e.g. harmonic series).



Convergence of Geometric series

Theorem

*The geometric series $\sum ar^{n-1}$ with $a \neq 0$, is
(i) convergent if $|r| < 1$ and $\sum ar^{n-1} = \frac{a}{1-r}$.*



Convergence of Geometric series

Theorem

The geometric series $\sum ar^{n-1}$ with $a \neq 0$, is

(i) convergent if $|r| < 1$ and $\sum ar^{n-1} = \frac{a}{1-r}$.

(ii) divergent if $|r| \geq 1$.



Convergence of Geometric series

Theorem

The geometric series $\sum ar^{n-1}$ with $a \neq 0$, is

(i) convergent if $|r| < 1$ and $\sum ar^{n-1} = \frac{a}{1-r}$.

(ii) divergent if $|r| \geq 1$.

Proof. We know that

$$S_n = a + ar + ar^2 + \cdots + ar^{n-1} = \frac{a(1-r^n)}{1-r}, \quad r \neq 1.$$



Convergence of Geometric series

Theorem

The geometric series $\sum ar^{n-1}$ with $a \neq 0$, is

(i) convergent if $|r| < 1$ and $\sum ar^{n-1} = \frac{a}{1-r}$.

(ii) divergent if $|r| \geq 1$.

Proof. We know that

$$S_n = a + ar + ar^2 + \cdots + ar^{n-1} = \frac{a(1-r^n)}{1-r}, \quad r \neq 1.$$

(i) Case I: $|r| < 1$



Convergence of Geometric series

Theorem

The geometric series $\sum ar^{n-1}$ with $a \neq 0$, is

(i) convergent if $|r| < 1$ and $\sum ar^{n-1} = \frac{a}{1-r}$.

(ii) divergent if $|r| \geq 1$.

Proof. We know that

$$S_n = a + ar + ar^2 + \cdots + ar^{n-1} = \frac{a(1-r^n)}{1-r}, \quad r \neq 1.$$

(i) Case I: $|r| < 1$

In this case $r^n \rightarrow 0$ as $n \rightarrow \infty$



Convergence of Geometric series

Theorem

The geometric series $\sum ar^{n-1}$ with $a \neq 0$, is

(i) convergent if $|r| < 1$ and $\sum ar^{n-1} = \frac{a}{1-r}$.

(ii) divergent if $|r| \geq 1$.

Proof. We know that

$$S_n = a + ar + ar^2 + \cdots + ar^{n-1} = \frac{a(1-r^n)}{1-r}, \quad r \neq 1.$$

(i) Case I: $|r| < 1$

In this case $r^n \rightarrow 0$ as $n \rightarrow \infty \implies S_n \rightarrow \frac{a}{1-r}$ as $n \rightarrow \infty$.



Convergence of Geometric series

Theorem

The geometric series $\sum ar^{n-1}$ with $a \neq 0$, is

(i) convergent if $|r| < 1$ and $\sum ar^{n-1} = \frac{a}{1-r}$.

(ii) divergent if $|r| \geq 1$.

Proof. We know that

$$S_n = a + ar + ar^2 + \cdots + ar^{n-1} = \frac{a(1-r^n)}{1-r}, \quad r \neq 1.$$

(i) Case I: $|r| < 1$

In this case $r^n \rightarrow 0$ as $n \rightarrow \infty \implies S_n \rightarrow \frac{a}{1-r}$ as $n \rightarrow \infty$.

Hence the GS is convergent and its sum is $\frac{a}{1-r}$.



(ii) Case II: $|r| > 1$



(ii) Case II: $|r| > 1$

$$|r| > 1 \implies r > 1 \text{ or } r < -1.$$



(ii) Case II: $|r| > 1$

$|r| > 1 \implies r > 1$ or $r < -1$.

● For $r > 1$: We have $r^n \rightarrow \infty$ as $n \rightarrow \infty$



(ii) Case II: $|r| > 1$

$|r| > 1 \implies r > 1$ or $r < -1$.

- For $r > 1$: We have $r^n \rightarrow \infty$ as $n \rightarrow \infty$ and hence S_n also tends to ∞ or $-\infty$ (depending on the sign of a).



(ii) Case II: $|r| > 1$

$|r| > 1 \implies r > 1$ or $r < -1$.

- For $r > 1$: We have $r^n \rightarrow \infty$ as $n \rightarrow \infty$ and hence S_n also tends to ∞ or $-\infty$ (depending on the sign of a).
- For $r < -1$: r^n oscillates between ∞ and $-\infty$ and



(ii) Case II: $|r| > 1$

$|r| > 1 \implies r > 1$ or $r < -1$.

- For $r > 1$: We have $r^n \rightarrow \infty$ as $n \rightarrow \infty$ and hence S_n also tends to ∞ or $-\infty$ (depending on the sign of a).
- For $r < -1$: r^n oscillates between ∞ and $-\infty$ and hence $\lim S_n$ does not exist.



(ii) Case II: $|r| > 1$

$|r| > 1 \implies r > 1$ or $r < -1$.

- For $r > 1$: We have $r^n \rightarrow \infty$ as $n \rightarrow \infty$ and hence S_n also tends to ∞ or $-\infty$ (depending on the sign of a).
- For $r < -1$: r^n oscillates between ∞ and $-\infty$ and hence $\lim S_n$ does not exist.

Therefore, GS is divergent in this case.



(ii) Case III: $|r| = 1$, i.e., $r = 1$ or $r = -1$



(ii) Case III: $|r| = 1$, i.e., $r = 1$ or $r = -1$

- If $r = 1$, then GS becomes $a + a + \cdots$,



(ii) Case III: $|r| = 1$, i.e., $r = 1$ or $r = -1$

- If $r = 1$, then GS becomes $a + a + \cdots$, $\implies S_n = na$;



(ii) Case III: $|r| = 1$, i.e., $r = 1$ or $r = -1$

- If $r = 1$, then GS becomes $a + a + \dots$, $\implies S_n = na$; this tends to ∞ or $-\infty$ as $n \rightarrow \infty$ (depending on the sign of a); and hence GS is divergent.



(ii) Case III: $|r| = 1$, i.e., $r = 1$ or $r = -1$

- If $r = 1$, then GS becomes $a + a + \cdots$, $\implies S_n = na$; this tends to ∞ or $-\infty$ as $n \rightarrow \infty$ (depending on the sign of a); and hence GS is divergent.
- If $r = -1$, then GS becomes $a - a + a - a + \cdots \implies$,



(ii) Case III: $|r| = 1$, i.e., $r = 1$ or $r = -1$

- If $r = 1$, then GS becomes $a + a + \cdots$, $\implies S_n = na$; this tends to ∞ or $-\infty$ as $n \rightarrow \infty$ (depending on the sign of a); and hence GS is divergent.
- If $r = -1$, then GS becomes $a - a + a - a + \cdots \implies$,

$$S_n = \begin{cases} 0, & \text{if } n \text{ is even} \\ a, & \text{if } n \text{ is odd.} \end{cases}$$



(ii) Case III: $|r| = 1$, i.e., $r = 1$ or $r = -1$

- If $r = 1$, then GS becomes $a + a + \dots$, $\implies S_n = na$; this tends to ∞ or $-\infty$ as $n \rightarrow \infty$ (depending on the sign of a); and hence GS is divergent.
- If $r = -1$, then GS becomes $a - a + a - a + \dots \implies$,

$$S_n = \begin{cases} 0, & \text{if } n \text{ is even} \\ a, & \text{if } n \text{ is odd.} \end{cases}$$

Since a is not zero, S_n does not tend to a unique limiting value and hence GS is divergent.



Example: Geometric series

Q.: $\sum_{n=0}^{\infty} \frac{\cos n\pi}{5^n}.$



Example: Geometric series

Q.: $\sum_{n=0}^{\infty} \frac{\cos n\pi}{5^n}.$

Sol. Here $r = \frac{-1}{5} < 1 \implies \sum_{n=0}^{\infty} \left(\frac{\cos n\pi}{5^n} \right)$ is convergent



Example: Geometric series

Q.: $\sum_{n=0}^{\infty} \frac{\cos n\pi}{5^n}.$

Sol. Here $r = \frac{-1}{5} < 1 \implies \sum_{n=0}^{\infty} \left(\frac{\cos n\pi}{5^n} \right)$ is convergent

$$\implies \sum_{n=0}^{\infty} \frac{\cos n\pi}{5^n} = \frac{1}{1 - \frac{-1}{5}} = \frac{5}{6}.$$



Example: Geometric series

Q.: $\sum_{n=0}^{\infty} \frac{e^{n\pi}}{\pi^{ne}}.$



Example: Geometric series

Q:. $\sum_{n=0}^{\infty} \frac{e^{n\pi}}{\pi^{ne}}.$

Sol. Is $r > 1$ or $r < 1$?



Example: Geometric series

Q.: $\sum_{n=0}^{\infty} \frac{e^{n\pi}}{\pi^{ne}}.$

Sol. Is $r > 1$ or $r < 1$? $r = \frac{e^{\pi}}{\pi^e} > 1$ (**How ?**)



Example: Geometric series

Q.: $\sum_{n=0}^{\infty} \frac{e^{n\pi}}{\pi^{ne}}.$

Sol. Is $r > 1$ or $r < 1$? $r = \frac{e^{\pi}}{\pi^e} > 1$ (**How ?**)

Let $f(x) = \frac{e^x}{x^e}$



Example: Geometric series

Q.: $\sum_{n=0}^{\infty} \frac{e^{n\pi}}{\pi^{ne}}.$

Sol. Is $r > 1$ or $r < 1$? $r = \frac{e^{\pi}}{\pi^e} > 1$ (**How ?**)

Let $f(x) = \frac{e^x}{x^e} \implies f'(x) = x^{-e-1}(x-e)e^x,$



Example: Geometric series

Q.: $\sum_{n=0}^{\infty} \frac{e^{n\pi}}{\pi^{ne}}.$

Sol. Is $r > 1$ or $r < 1$? $r = \frac{e^\pi}{\pi^e} > 1$ (**How ?**)

Let $f(x) = \frac{e^x}{x^e} \implies f'(x) = x^{-e-1}(x-e)e^x$, i.e., $f'(x) > 0$ for $x > e \implies f$ is an increasing function,



Example: Geometric series

Q.: $\sum_{n=0}^{\infty} \frac{e^{n\pi}}{\pi^{ne}}.$

Sol. Is $r > 1$ or $r < 1$? $r = \frac{e^\pi}{\pi^e} > 1$ (**How ?**)

Let $f(x) = \frac{e^x}{x^e} \implies f'(x) = x^{-e-1}(x-e)e^x$, i.e., $f'(x) > 0$ for $x > e \implies f$ is an increasing function, i.e., if $x_1 > x_2$, then $f(x_1) > f(x_2)$.



Example: Geometric series

Q.: $\sum_{n=0}^{\infty} \frac{e^{n\pi}}{\pi^{ne}}.$

Sol. Is $r > 1$ or $r < 1$? $r = \frac{e^\pi}{\pi^e} > 1$ (**How ?**)

Let $f(x) = \frac{e^x}{x^e} \implies f'(x) = x^{-e-1}(x-e)e^x$, i.e., $f'(x) > 0$ for $x > e \implies f$ is an increasing function, i.e., if $x_1 > x_2$, then $f(x_1) > f(x_2)$.

Let $x_1 = \pi, x_2 = e$. Now,

$$\pi > e \implies f(\pi) > f(e) \implies \frac{e^\pi}{\pi^e} > 1$$



Example: Geometric series

Q.: $\sum_{n=0}^{\infty} \frac{e^{n\pi}}{\pi^{ne}}.$

Sol. Is $r > 1$ or $r < 1$? $r = \frac{e^\pi}{\pi^e} > 1$ (**How ?**)

Let $f(x) = \frac{e^x}{x^e} \implies f'(x) = x^{-e-1}(x-e)e^x$, i.e., $f'(x) > 0$ for $x > e \implies f$ is an increasing function, i.e., if $x_1 > x_2$, then $f(x_1) > f(x_2)$.

Let $x_1 = \pi, x_2 = e$. Now,

$$\pi > e \implies f(\pi) > f(e) \implies \frac{e^\pi}{\pi^e} > 1$$

$$\implies \sum_{n=0}^{\infty} \frac{e^{n\pi}}{\pi^{ne}} \text{ is divergent.}$$



Method 3: The n th term test for divergence



Method 3: The n th term test for divergence

If $\lim a_n$ fails to exist or is different from zero, then $\sum a_n$ is divergent.



Method 3: The n th term test for divergence

If $\lim a_n$ fails to exist or is different from zero, then $\sum a_n$ is divergent.

Q: $\sum \ln \frac{1}{3^n}$.



Method 3: The n th term test for divergence

If $\lim a_n$ fails to exist or is different from zero, then $\sum a_n$ is divergent.

Q.: $\sum \ln \frac{1}{3^n}$.

Sol. Here

$$\lim a_n = \lim \ln \frac{1}{3^n} = \lim (\ln 1 - \ln 3^n) = -\lim n \ln 3 = -\infty \neq 0,$$



Method 3: The n th term test for divergence

If $\lim a_n$ fails to exist or is different from zero, then $\sum a_n$ is divergent.

Q:. $\sum \ln \frac{1}{3^n}$.

Sol. Here

$\lim a_n = \lim \ln \frac{1}{3^n} = \lim (\ln 1 - \ln 3^n) = -\lim n \ln 3 = -\infty \neq 0$,
therefore $\sum \ln \frac{1}{3^n}$ is divergent.



Theorem (Theorem 7. The necessary condition for a series to be convergent)

If $\sum a_n$ converges, then $\lim a_n = 0$.



Theorem (Theorem 7. The necessary condition for a series to be convergent)

If $\sum a_n$ converges, then $\lim a_n = 0$.

Proof. We have

$$a_1 = S_1$$

$$a_n = S_n - S_{n-1} \text{ for } n = 2, 3, 4, \dots$$



Theorem (Theorem 7. The necessary condition for a series to be convergent)

If $\sum a_n$ converges, then $\lim a_n = 0$.

Proof. We have

$$a_1 = S_1$$

$$a_n = S_n - S_{n-1} \text{ for } n = 2, 3, 4, \dots$$

If the given series converges then $S_n \rightarrow L$. Therefore,
 $\lim a_n = \lim(S_n - S_{n-1}) \rightarrow L - L = 0$.



Q.: $\sum_{n=0}^{\infty} \frac{e^n}{e^n + n}.$



Q.: $\sum_{n=0}^{\infty} \frac{e^n}{e^{n+n}}.$

Sol. Here $\lim a_n = \lim \frac{e^n}{e^{n+n}} = \lim \frac{1}{1+\frac{n}{e^n}} = 1 \neq 0$, therefore

$\sum_{n=0}^{\infty} \frac{e^n}{e^{n+n}}$ is divergent.



Q.: $\sum_{n=0}^{\infty} \frac{e^n}{e^{n+n}}.$

Sol. Here $\lim a_n = \lim \frac{e^n}{e^{n+n}} = \lim \frac{1}{1+\frac{n}{e^n}} = 1 \neq 0$, therefore

$\sum_{n=0}^{\infty} \frac{e^n}{e^{n+n}}$ is divergent.

Q.: $\sum \frac{2^n-1}{2^{n+1}}.$



Q.: $\sum_{n=0}^{\infty} \frac{e^n}{e^{n+n}}.$

Sol. Here $\lim a_n = \lim \frac{e^n}{e^{n+n}} = \lim \frac{1}{1+\frac{n}{e^n}} = 1 \neq 0$, therefore

$\sum_{n=0}^{\infty} \frac{e^n}{e^{n+n}}$ is divergent.

Q.: $\sum \frac{2^n-1}{2^{n+1}}.$

Sol. Divergent.



Q.: $\sum_{n=0}^{\infty} \frac{e^n}{e^{n+n}}.$

Sol. Here $\lim a_n = \lim \frac{e^n}{e^{n+n}} = \lim \frac{1}{1+\frac{n}{e^n}} = 1 \neq 0$, therefore

$\sum_{n=0}^{\infty} \frac{e^n}{e^{n+n}}$ is divergent.

Q.: $\sum \frac{2^n-1}{2^{n+1}}.$

Sol. Divergent.

Q.: $\sum \left(1 - \frac{1}{n}\right).$



Q.: $\sum_{n=0}^{\infty} \frac{e^n}{e^{n+n}}.$

Sol. Here $\lim a_n = \lim \frac{e^n}{e^{n+n}} = \lim \frac{1}{1+\frac{n}{e^n}} = 1 \neq 0$, therefore

$\sum_{n=0}^{\infty} \frac{e^n}{e^{n+n}}$ is divergent.

Q.: $\sum \frac{2^n-1}{2^{n+1}}.$

Sol. Divergent.

Q.: $\sum \left(1 - \frac{1}{n}\right).$

Sol. Divergent.



Theorem (Theorem 8.)

If $\sum a_n$ and $\sum b_n$ are two convergent series having sums A and B respectively, then



Theorem (Theorem 8.)

If $\sum a_n$ and $\sum b_n$ are two convergent series having sums A and B respectively, then

- $\sum(a_n \pm b_n) = \sum a_n \pm \sum b_n = A \pm B.$
- $\sum ka_n = k \sum a_n = kA$ (for any real number k).



For next few classes, series will be assumed to be series of nonnegative terms *i.e.*, $a_n \geq 0$ for all n . This means that $S_n \leq S_{n+1}$ for all n . That is the sequence $\{S_n\}$ is nondecreasing.



For next few classes, series will be assumed to be series of nonnegative terms *i.e.*, $a_n \geq 0$ for all n . This means that $S_n \leq S_{n+1}$ for all n . That is the sequence $\{S_n\}$ is nondecreasing.

Corollary (Corollary of Theorem 6)

A series $\sum a_n$ of nonnegative terms converges iff S_n is bounded from above.



Section 10.3

The Integral Test



Method 4: The Integral Test

Theorem (Theorem 9.)

Let $\sum a_n$ be a series of positive terms.



Method 4: The Integral Test

Theorem (Theorem 9.)

Let $\sum a_n$ be a series of positive terms. Let $f(x)$ be a positive, continuous, and decreasing function for all $x \geq N$ for some \underline{N} ; and let $\underline{f(n) = a_n}$ for all n .



Method 4: The Integral Test

Theorem (Theorem 9.)

Let $\sum a_n$ be a series of positive terms. Let $f(x)$ be a positive, continuous, and decreasing function for all $x \geq N$ for some \underline{N} ; and let $\underline{f(n) = a_n}$ for all n . Then the series $\sum_{n=N}^{\infty} a_n$ and the integral $\int_N^{\infty} f(x)dx$ both converge or both diverge.



Method 4: The Integral Test

Theorem (Theorem 9.)

Let $\sum a_n$ be a series of positive terms. Let $f(x)$ be a positive, continuous, and decreasing function for all $x \geq N$ for some \underline{N} ; and let $\underline{f(n) = a_n}$ for all n . Then the series $\sum_{n=N}^{\infty} a_n$ and the integral $\int_N^{\infty} f(x)dx$ both converge or both diverge.

Remark. In the case of convergence, the sum of the series and the value of integral will be different.



Q:. Check the convergence of the following series:

$$\sum \frac{n}{n^2+1}.$$



Q:. Check the convergence of the following series:

$$\sum \frac{n}{n^2+1}.$$

Sol. Consider the function $f(x) = \frac{x}{x^2+1}$.



Q:. Check the convergence of the following series:

$$\sum \frac{n}{n^2+1}.$$

Sol. Consider the function $f(x) = \frac{x}{x^2+1}$. Now,

$$f'(x) = \frac{1-x^2}{(x^2+1)^2}.$$



Q:. Check the convergence of the following series:

$$\sum \frac{n}{n^2+1}.$$

Sol. Consider the function $f(x) = \frac{x}{x^2+1}$. Now,

$f'(x) = \frac{1-x^2}{(x^2+1)^2}$. Thus $f'(x) \leq 0$ when $x \geq 1 \implies f(x)$ is decreasing for all $x \geq 1$.



Q:. Check the convergence of the following series:

$$\sum \frac{n}{n^2+1}.$$

Sol. Consider the function $f(x) = \frac{x}{x^2+1}$. Now,

$f'(x) = \frac{1-x^2}{(x^2+1)^2}$. Thus $f'(x) \leq 0$ when $x \geq 1 \implies f(x)$ is decreasing for all $x \geq 1$. Thus $f(x)$ is positive, continuous and decreasing for all $x \geq 1$.



Q:. Check the convergence of the following series:

$$\sum \frac{n}{n^2+1}.$$

Sol. Consider the function $f(x) = \frac{x}{x^2+1}$. Now,

$f'(x) = \frac{1-x^2}{(x^2+1)^2}$. Thus $f'(x) \leq 0$ when $x \geq 1 \implies f(x)$ is decreasing for all $x \geq 1$. Thus $f(x)$ is positive, continuous and decreasing for all $x \geq 1$. Now,

$$\int_1^{\infty} \frac{x}{x^2+1} dx =$$



Q:. Check the convergence of the following series:

$$\sum \frac{n}{n^2+1}.$$

Sol. Consider the function $f(x) = \frac{x}{x^2+1}$. Now,

$f'(x) = \frac{1-x^2}{(x^2+1)^2}$. Thus $f'(x) \leq 0$ when $x \geq 1 \implies f(x)$ is decreasing for all $x \geq 1$. Thus $f(x)$ is positive, continuous and decreasing for all $x \geq 1$. Now,

$$\int_1^{\infty} \frac{x}{x^2+1} dx =$$

$$\frac{1}{2} \lim_{b \rightarrow \infty} [\ln(x^2+1)]_1^b \rightarrow \infty.$$

Hence by integral test the series is divergent.



Q: $\sum ne^{-n^2}$



Q:. $\sum ne^{-n^2}$

Sol. The function $f(x) = xe^{-x^2}$ is positive, continuous and decreasing for all $x \geq 1$.



Q.: $\sum ne^{-n^2}$

Sol. The function $f(x) = xe^{-x^2}$ is positive, continuous and decreasing for all $x \geq 1$. Now, we have

$$\begin{aligned}\int_1^\infty \frac{x}{e^{x^2}} dx &= \frac{1}{2} \int_1^\infty \frac{1}{e^u} du \quad (\text{set } u = x^2) \\ &= \frac{-1}{2} \lim_{b \rightarrow \infty} [e^{-u}]_1^b = \frac{1}{2e}\end{aligned}$$



Q.: $\sum ne^{-n^2}$

Sol. The function $f(x) = xe^{-x^2}$ is positive, continuous and decreasing for all $x \geq 1$. Now, we have

$$\begin{aligned}\int_1^{\infty} \frac{x}{e^{x^2}} dx &= \frac{1}{2} \int_1^{\infty} \frac{1}{e^u} du \quad (\text{set } u = x^2) \\ &= \frac{-1}{2} \lim_{b \rightarrow \infty} [e^{-u}]_1^b = \frac{1}{2e}\end{aligned}$$

Hence by integral test the series $\sum ne^{-n^2}$ is convergent.



Q: $\sum_{n=2}^{\infty} \frac{1}{n \ln n}$.



Q:. $\sum_{n=2} \frac{1}{n \ln n}$.

Sol. The function $f(x) = \frac{1}{x \ln x}$ is positive, continuous and decreasing for all $x \geq 2$.



Q.: $\sum_{n=2}^{\infty} \frac{1}{n \ln n}$.

Sol. The function $f(x) = \frac{1}{x \ln x}$ is positive, continuous and decreasing for all $x \geq 2$. Here the n^{th} term is $f(n) = \frac{1}{(n+1) \ln(n+1)}$. Now, we have

$$\int_2^{\infty} \frac{1}{(x+1) \ln(x+1)} dx \rightarrow \infty.$$



Q.: $\sum_{n=2}^{\infty} \frac{1}{n \ln n}$.

Sol. The function $f(x) = \frac{1}{x \ln x}$ is positive, continuous and decreasing for all $x \geq 2$. Here the n^{th} term is $f(n) = \frac{1}{(n+1) \ln(n+1)}$. Now, we have

$$\int_2^{\infty} \frac{1}{(x+1) \ln(x+1)} dx \rightarrow \infty.$$

(set $\ln(x+1) = t$).



Q.: $\sum_{n=2}^{\infty} \frac{1}{n \ln n}$.

Sol. The function $f(x) = \frac{1}{x \ln x}$ is positive, continuous and decreasing for all $x \geq 2$. Here the n^{th} term is $f(n) = \frac{1}{(n+1) \ln(n+1)}$. Now, we have

$$\int_2^{\infty} \frac{1}{(x+1) \ln(x+1)} dx \rightarrow \infty.$$

(set $\ln(x+1) = t$). Hence by integral test the series $\sum \frac{1}{n \ln n}$ is divergent.



Method 5: p -series test

The p -series $\sum \frac{1}{n^p}$ is convergent for $p > 1$, and divergent for $p \leq 1$, where p is a real constant.



Method 5: p -series test

The p -series $\sum \frac{1}{n^p}$ is convergent for $p > 1$, and divergent for $p \leq 1$, where p is a real constant.

Proof. For $p \leq 0$, the series is trivially divergent



Method 5: p -series test

The p -series $\sum \frac{1}{n^p}$ is convergent for $p > 1$, and divergent for $p \leq 1$, where p is a real constant.

Proof. For $p \leq 0$, the series is trivially divergent (as $\lim a_n \neq 0$).



Method 5: p -series test

The p -series $\sum \frac{1}{n^p}$ is convergent for $p > 1$, and divergent for $p \leq 1$, where p is a real constant.

Proof. For $p \leq 0$, the series is trivially divergent (as $\lim a_n \neq 0$).

Let $p > 0$. Now for $p > 0$, the function $f(x) = \frac{1}{x^p}$ is positive, continuous and decreasing when $x \geq 1$



Method 5: p -series test

The p -series $\sum \frac{1}{n^p}$ is convergent for $p > 1$, and divergent for $p \leq 1$, where p is a real constant.

Proof. For $p \leq 0$, the series is trivially divergent (as $\lim a_n \neq 0$).

Let $p > 0$. Now for $p > 0$, the function $f(x) = \frac{1}{x^p}$ is positive, continuous and decreasing when $x \geq 1$ (verify yourself). Hence, we can apply integral test:



- If $p > 1$,



- If $p > 1$,

$$\int_1^{\infty} f(x) dx = \int_1^{\infty} \frac{1}{x^p} dx = \lim_{b \rightarrow \infty} \left[\frac{x^{-p+1}}{-p+1} \right]_1^b$$



- If $p > 1$,

$$\begin{aligned}\int_1^\infty f(x) dx &= \int_1^\infty \frac{1}{x^p} dx = \lim_{b \rightarrow \infty} \left[\frac{x^{-p+1}}{-p+1} \right]_1^b \\ &= \frac{1}{-p+1} \lim_{b \rightarrow \infty} \left(\frac{1}{b^{p-1}} - 1 \right) = \frac{1}{1-p} (0 - 1) = \frac{1}{p-1}\end{aligned}$$



- If $p > 1$,

$$\begin{aligned}\int_1^\infty f(x) dx &= \int_1^\infty \frac{1}{x^p} dx = \lim_{b \rightarrow \infty} \left[\frac{x^{-p+1}}{-p+1} \right]_1^b \\ &= \frac{1}{-p+1} \lim_{b \rightarrow \infty} \left(\frac{1}{b^{p-1}} - 1 \right) = \frac{1}{1-p} (0 - 1) = \frac{1}{p-1} \\ &\quad (b^{p-1} \rightarrow \infty \text{ as } b \rightarrow \infty \text{ because})\end{aligned}$$



- If $p > 1$,

$$\begin{aligned}\int_1^\infty f(x) dx &= \int_1^\infty \frac{1}{x^p} dx = \lim_{b \rightarrow \infty} \left[\frac{x^{-p+1}}{-p+1} \right]_1^b \\ &= \frac{1}{-p+1} \lim_{b \rightarrow \infty} \left(\frac{1}{b^{p-1}} - 1 \right) = \frac{1}{1-p} (0 - 1) = \frac{1}{p-1} \\ &\quad (b^{p-1} \rightarrow \infty \text{ as } b \rightarrow \infty \text{ because } p-1 > 0).\end{aligned}$$



- If $p > 1$,

$$\begin{aligned}\int_1^\infty f(x) dx &= \int_1^\infty \frac{1}{x^p} dx = \lim_{b \rightarrow \infty} \left[\frac{x^{-p+1}}{-p+1} \right]_1^b \\ &= \frac{1}{-p+1} \lim_{b \rightarrow \infty} \left(\frac{1}{b^{p-1}} - 1 \right) = \frac{1}{1-p} (0 - 1) = \frac{1}{p-1} \\ &\quad (b^{p-1} \rightarrow \infty \text{ as } b \rightarrow \infty \text{ because } p-1 > 0). \text{ Thus, the} \\ &\quad \text{series converges for } p > 1.\end{aligned}$$



- If $p > 1$,

$$\begin{aligned}\int_1^\infty f(x) dx &= \int_1^\infty \frac{1}{x^p} dx = \lim_{b \rightarrow \infty} \left[\frac{x^{-p+1}}{-p+1} \right]_1^b \\ &= \frac{1}{-p+1} \lim_{b \rightarrow \infty} \left(\frac{1}{b^{p-1}} - 1 \right) = \frac{1}{1-p} (0 - 1) = \frac{1}{p-1} \\ &\quad (b^{p-1} \rightarrow \infty \text{ as } b \rightarrow \infty \text{ because } p-1 > 0). \text{ Thus, the series converges for } p > 1.\end{aligned}$$

- If $0 < p < 1$, then

$$\int_1^\infty f(x) dx = \int_1^\infty \frac{1}{x^p} dx = \lim_{b \rightarrow \infty} \left[\frac{x^{-p+1}}{-p+1} \right]_1^b \rightarrow \infty.$$

Hence, the series is divergent.



- If $p > 1$,

$$\begin{aligned}\int_1^\infty f(x) dx &= \int_1^\infty \frac{1}{x^p} dx = \lim_{b \rightarrow \infty} \left[\frac{x^{-p+1}}{-p+1} \right]_1^b \\ &= \frac{1}{-p+1} \lim_{b \rightarrow \infty} \left(\frac{1}{b^{p-1}} - 1 \right) = \frac{1}{1-p} (0 - 1) = \frac{1}{p-1} \\ &\quad (b^{p-1} \rightarrow \infty \text{ as } b \rightarrow \infty \text{ because } p-1 > 0). \text{ Thus, the series converges for } p > 1.\end{aligned}$$

- If $0 < p < 1$, then

$$\int_1^\infty f(x) dx = \int_1^\infty \frac{1}{x^p} dx = \lim_{b \rightarrow \infty} \left[\frac{x^{-p+1}}{-p+1} \right]_1^b \rightarrow \infty.$$

Hence, the series is divergent.

- For $p = 1$, we have



- If $p > 1$,

$$\begin{aligned}\int_1^\infty f(x) dx &= \int_1^\infty \frac{1}{x^p} dx = \lim_{b \rightarrow \infty} \left[\frac{x^{-p+1}}{-p+1} \right]_1^b \\ &= \frac{1}{-p+1} \lim_{b \rightarrow \infty} \left(\frac{1}{b^{p-1}} - 1 \right) = \frac{1}{1-p} (0 - 1) = \frac{1}{p-1} \\ &\quad (b^{p-1} \rightarrow \infty \text{ as } b \rightarrow \infty \text{ because } p-1 > 0). \text{ Thus, the series converges for } p > 1.\end{aligned}$$

- If $0 < p < 1$, then

$$\int_1^\infty f(x) dx = \int_1^\infty \frac{1}{x^p} dx = \lim_{b \rightarrow \infty} \left[\frac{x^{-p+1}}{-p+1} \right]_1^b \rightarrow \infty.$$

Hence, the series is divergent.

- For $p = 1$, we have $\int_1^\infty \frac{1}{x} dx = \lim_{b \rightarrow \infty} [\ln x]_1^b \rightarrow \infty$.
Hence, the series is divergent.



Q:. Which of the following series converge and which diverge?

- $\sum \frac{1}{n}$. (Harmonic series)



Q:. Which of the following series converge and which diverge?

- $\sum \frac{1}{n}$. (Harmonic series)
- $\sum \frac{1}{n^2}$.



Q:. Which of the following series converge and which diverge?

- $\sum \frac{1}{n}$. (Harmonic series)
- $\sum \frac{1}{n^2}$.

Ans. Divergent, Convergent.



Homework. For what $p > 0$, does the series $\sum_2^{\infty} \frac{1}{n(\ln n)^p}$ converge?



Homework. For what $p > 0$, does the series $\sum_2^{\infty} \frac{1}{n(\ln n)^p}$ converge?

Ans. $p > 1$.

Homework. Check the convergence of the following series:

- $\sum \frac{n^2}{e^{n/3}}.$



Homework. For what $p > 0$, does the series $\sum_2^{\infty} \frac{1}{n(\ln n)^p}$ converge?

Ans. $p > 1$.

Homework. Check the convergence of the following series:

- $\sum \frac{n^2}{e^{n/3}}$.
- $\sum \frac{5^n}{4^n + 3}$.



Homework. For what $p > 0$, does the series $\sum_2^{\infty} \frac{1}{n(\ln n)^p}$ converge?

Ans. $p > 1$.

Homework. Check the convergence of the following series:

- $\sum \frac{n^2}{e^{n/3}}.$
- $\sum \frac{5^n}{4^n+3}.$
- $\sum \frac{\ln n}{\sqrt{n}}.$

Ans. Convergent, Divergent, Divergent.



Convergence of series: Examples

Q: $\sum \ln \sqrt{n+1} - \ln \sqrt{n}$



Convergence of series: Examples

Q:. $\sum \ln \sqrt{n+1} - \ln \sqrt{n}$

Sol. Divergent



Convergence of series: Examples

Q:. $\sum \ln \sqrt{n+1} - \ln \sqrt{n}$

Sol. Divergent (telescoping series)



Convergence of series: Examples

Q:. $\sum \ln \sqrt{n+1} - \ln \sqrt{n}$

Sol. Divergent (telescoping series)

Q:. $\sum \left(\frac{1}{2^n} + \frac{(-1)^n}{5^n} \right)$



Convergence of series: Examples

Q:. $\sum \ln \sqrt{n+1} - \ln \sqrt{n}$

Sol. Divergent (telescoping series)

Q:. $\sum \left(\frac{1}{2^n} + \frac{(-1)^n}{5^n} \right)$

Sol. Convergent



Convergence of series: Examples

Q:. $\sum \ln \sqrt{n+1} - \ln \sqrt{n}$

Sol. Divergent (telescoping series)

Q:. $\sum \left(\frac{1}{2^n} + \frac{(-1)^n}{5^n} \right)$

Sol. Convergent (sum of geometric series)



Convergence of series: Examples

Q:. $\sum \ln \sqrt{n+1} - \ln \sqrt{n}$

Sol. Divergent (telescoping series)

Q:. $\sum \left(\frac{1}{2^n} + \frac{(-1)^n}{5^n} \right)$

Sol. Convergent (sum of geometric series)

Q:. $\sum \frac{n}{n+10}$



Convergence of series: Examples

Q:. $\sum \ln \sqrt{n+1} - \ln \sqrt{n}$

Sol. Divergent (telescoping series)

Q:. $\sum \left(\frac{1}{2^n} + \frac{(-1)^n}{5^n} \right)$

Sol. Convergent (sum of geometric series)

Q:. $\sum \frac{n}{n+10}$

Sol. Divergent



Convergence of series: Examples

Q:. $\sum \ln \sqrt{n+1} - \ln \sqrt{n}$

Sol. Divergent (telescoping series)

Q:. $\sum \left(\frac{1}{2^n} + \frac{(-1)^n}{5^n} \right)$

Sol. Convergent (sum of geometric series)

Q:. $\sum \frac{n}{n+10}$

Sol. Divergent (n^{th} term test)



Q.: $\sum \frac{n^2}{e^{n/3}}$ (use Integral test)



Q:. $\sum \frac{n^2}{e^{n/3}}$ (use Integral test)
Sol. Convergent



Q:. $\sum \frac{n^2}{e^{n/3}}$ (use Integral test)

Sol. Convergent

$$f'(x) = \frac{-x(x-6)}{3e^{x/3}} < 0 \text{ for } x > 6.$$



Q:. $\sum \frac{n^2}{e^{n/3}}$ (use Integral test)

Sol. Convergent

$$f'(x) = \frac{-x(x-6)}{3e^{x/3}} < 0 \text{ for } x > 6.$$

$$\int_7^{\infty} f(x) dx = \frac{327}{e^{7/3}} \implies$$



Q.: $\sum \frac{n^2}{e^{n/3}}$ (use Integral test)

Sol. Convergent

$$f'(x) = \frac{-x(x-6)}{3e^{x/3}} < 0 \text{ for } x > 6.$$

$$\int_7^{\infty} f(x) dx = \frac{327}{e^{7/3}} \implies \sum_{n=7}^{\infty} \frac{n^2}{e^{n/3}} \text{ is convergent}$$



Q.: $\sum \frac{n^2}{e^{n/3}}$ (use Integral test)

Sol. Convergent

$$f'(x) = \frac{-x(x-6)}{3e^{x/3}} < 0 \text{ for } x > 6.$$

$$\int_7^{\infty} f(x) dx = \frac{327}{e^{7/3}} \implies \sum_{n=7}^{\infty} \frac{n^2}{e^{n/3}} \text{ is convergent} \implies$$

$$\sum_{n=1}^{\infty} \frac{n^2}{e^{n/3}} \text{ is convergent.}$$



Section 10.4

Comparison Tests



Method 6: The Direct Comparison Test

Theorem (Theorem 10.)

Let $\sum a_n$ be a series of nonnegative terms.



Method 6: The Direct Comparison Test

Theorem (Theorem 10.)

Let $\sum a_n$ be a series of nonnegative terms.

(a) If $a_n \leq c_n$ for all $n \geq N$ and $\sum c_n$ is convergent, then $\sum a_n$ will be convergent.



Method 6: The Direct Comparison Test

Theorem (Theorem 10.)

Let $\sum a_n$ be a series of nonnegative terms.

- (a) If $a_n \leq c_n$ for all $n \geq N$ and $\sum c_n$ is convergent, then $\sum a_n$ will be convergent.*
- (b) If $a_n \geq d_n$ for all $n \geq N$ and $\sum d_n$ is a divergent series of **nonnegative terms**, then $\sum a_n$ will be divergent.*



Q: Investigate the convergence or divergence of $\sum \frac{1}{n^3+5n}$.



Q:. Investigate the convergence or divergence of $\sum \frac{1}{n^3+5n}$.

Sol. Here, $a_n = \frac{1}{n^3+5n}$.



Q:. Investigate the convergence or divergence of $\sum \frac{1}{n^3+5n}$.

Sol. Here, $a_n = \frac{1}{n^3+5n}$. Now, $n^3 + 5n \geq n^3 \implies \frac{1}{n^3+5n} \leq \frac{1}{n^3}$ for all n .



Q:. Investigate the convergence or divergence of $\sum \frac{1}{n^3+5n}$.

Sol. Here, $a_n = \frac{1}{n^3+5n}$. Now, $n^3 + 5n \geq n^3 \implies \frac{1}{n^3+5n} \leq \frac{1}{n^3}$ for all n . Therefore, we have,

$$a_n \leq \frac{1}{n^3} = c_n \text{ (say) for all } n.$$



Q.: Investigate the convergence or divergence of $\sum \frac{1}{n^3+5n}$.

Sol. Here, $a_n = \frac{1}{n^3+5n}$. Now, $n^3 + 5n \geq n^3 \implies \frac{1}{n^3+5n} \leq \frac{1}{n^3}$ for all n . Therefore, we have,

$$a_n \leq \frac{1}{n^3} = c_n \text{ (say) for all } n.$$

Moreover, from p -test, $\sum c_n$ is convergent.



Q:. Investigate the convergence or divergence of $\sum \frac{1}{n^3+5n}$.

Sol. Here, $a_n = \frac{1}{n^3+5n}$. Now, $n^3 + 5n \geq n^3 \implies \frac{1}{n^3+5n} \leq \frac{1}{n^3}$ for all n . Therefore, we have,

$$a_n \leq \frac{1}{n^3} = c_n \text{ (say) for all } n.$$

Moreover, from p -test, $\sum c_n$ is convergent. Therefore, by DCT, $\sum a_n$ is convergent.



Q.: $\sum \frac{5^n + 1}{2^n - 1}$



Q.: $\sum \frac{5^n+1}{2^n-1}$

Sol. Here $a_n = \frac{5^n+1}{2^n-1} \geq \frac{5^n}{2^n-1} \geq \frac{5^n}{2^n} = \left(\frac{5}{2}\right)^n$ for all n .



Q.: $\sum \frac{5^n+1}{2^n-1}$

Sol. Here $a_n = \frac{5^n+1}{2^n-1} \geq \frac{5^n}{2^n-1} \geq \frac{5^n}{2^n} = \left(\frac{5}{2}\right)^n$ for all n .

Since the series $\sum \left(\frac{5}{2}\right)^n$ is a divergent geometric series ($|r| = \frac{5}{2} > 1$),



Q.: $\sum \frac{5^n+1}{2^n-1}$

Sol. Here $a_n = \frac{5^n+1}{2^n-1} \geq \frac{5^n}{2^n-1} \geq \frac{5^n}{2^n} = \left(\frac{5}{2}\right)^n$ for all n .

Since the series $\sum \left(\frac{5}{2}\right)^n$ is a divergent geometric series ($|r| = \frac{5}{2} > 1$), therefore by DCT, $\sum \frac{5^n+1}{2^n-1}$ diverges.



Q.: $\sum \frac{n-1}{n^4+2}$



Q.: $\sum \frac{n-1}{n^4+2}$

Sol. $n^4 \leq n^4 + 2 \implies \frac{1}{n^4} \geq \frac{1}{n^4+2} \implies \frac{n}{n^4} \geq \frac{n}{n^4+2} \implies \frac{1}{n^3} \geq \frac{n}{n^4+2}$
 $\frac{n}{n^4+2} \geq \frac{n-1}{n^4+2}.$



Q.: $\sum \frac{n-1}{n^4+2}$

Sol. $n^4 \leq n^4 + 2 \implies \frac{1}{n^4} \geq \frac{1}{n^4+2} \implies \frac{n}{n^4} \geq \frac{n}{n^4+2} \implies \frac{1}{n^3} \geq \frac{n}{n^4+2} \geq \frac{n-1}{n^4+2}$. Let $c_n = \frac{1}{n^3}$.



Q.: $\sum \frac{n-1}{n^4+2}$

Sol. $n^4 \leq n^4 + 2 \implies \frac{1}{n^4} \geq \frac{1}{n^4+2} \implies \frac{n}{n^4} \geq \frac{n}{n^4+2} \implies \frac{1}{n^3} \geq \frac{n}{n^4+2} \geq \frac{n-1}{n^4+2}$. Let $c_n = \frac{1}{n^3}$.

From p -test, $\sum c_n$ is convergent $\implies \sum a_n$ is convergent.



Q: $\sum \sqrt{\frac{n+4}{n^4+4}}$



Q.: $\sum \sqrt{\frac{n+4}{n^4+4}}$

Sol. We have $n^3 \leq n^4 \implies 4n^3 \leq 4n^4 \implies n^4 + 4n^3 \leq n^4 + 4n^4 = 5n^4 \implies n^4 + 4n^3 \leq 5n^4 + 20 = 5(n^4 + 4)$



Q:. $\sum \sqrt{\frac{n+4}{n^4+4}}$

Sol. We have $n^3 \leq n^4 \implies 4n^3 \leq 4n^4 \implies n^4 + 4n^3 \leq n^4 + 4n^4 = 5n^4 \implies n^4 + 4n^3 \leq 5n^4 + 20 = 5(n^4 + 4)$
 $\implies \frac{n^4 + 4n^3}{n^4 + 4} \leq 5 \implies \frac{n+4}{n^4 + 4} \leq \frac{5}{n^3} \implies \sqrt{\frac{n+4}{n^4 + 4}} \leq \sqrt{\frac{5}{n^3}} \implies$



Q.: $\sum \sqrt{\frac{n+4}{n^4+4}}$

Sol. We have $n^3 \leq n^4 \implies 4n^3 \leq 4n^4 \implies n^4 + 4n^3 \leq n^4 + 4n^4 = 5n^4 \implies n^4 + 4n^3 \leq 5n^4 + 20 = 5(n^4 + 4)$
 $\implies \frac{n^4 + 4n^3}{n^4 + 4} \leq 5 \implies \frac{n+4}{n^4 + 4} \leq \frac{5}{n^3} \implies \sqrt{\frac{n+4}{n^4 + 4}} \leq \sqrt{\frac{5}{n^3}} \implies$
 $c_n = \frac{\sqrt{5}}{n^{3/2}}.$



Q.: $\sum \sqrt{\frac{n+4}{n^4+4}}$

Sol. We have $n^3 \leq n^4 \implies 4n^3 \leq 4n^4 \implies n^4 + 4n^3 \leq n^4 + 4n^4 = 5n^4 \implies n^4 + 4n^3 \leq 5n^4 + 20 = 5(n^4 + 4)$
 $\implies \frac{n^4 + 4n^3}{n^4 + 4} \leq 5 \implies \frac{n+4}{n^4+4} \leq \frac{5}{n^3} \implies \sqrt{\frac{n+4}{n^4+4}} \leq \sqrt{\frac{5}{n^3}} \implies$
 $c_n = \frac{\sqrt{5}}{n^{3/2}}.$

From p -test, $\sum c_n$ is convergent $\implies \sum a_n$ is convergent.



Q.: $\sum \frac{1}{n^3 - 5n}$



Q.: $\sum \frac{1}{n^3-5n}$

Sol. Here, $a_n = \frac{1}{n^3-5n}$. Now, $\frac{1}{n^3-5n} \geq \frac{1}{n^3} \geq 0$ for all $n \geq 3$.
But, from p -test, $\sum \frac{1}{n^3}$ is convergent.



Q.: $\sum \frac{1}{n^3-5n}$

Sol. Here, $a_n = \frac{1}{n^3-5n}$. Now, $\frac{1}{n^3-5n} \geq \frac{1}{n^3} \geq 0$ for all $n \geq 3$.

But, from p -test, $\sum \frac{1}{n^3}$ is convergent. Therefore, by DCT, we can not conclude anything.



Q.: $\sum \frac{\sqrt{n}}{n^2+1}$

Sol. Convergent

Q.: $\sum \frac{n+2^n}{n^2 2^n}$

Sol. Convergent

Q.: $\sum \frac{1}{1+2+3+\dots+n}$

Sol. Convergent



Method 7: The Limit Comparison Test

Theorem (Theorem 11.)

Suppose a_n and b_n are positive for all $n \geq N$.



Method 7: The Limit Comparison Test

Theorem (Theorem 11.)

Suppose a_n and b_n are positive for all $n \geq N$.

(a) If $\lim \frac{a_n}{b_n} = c$ (finite and non zero), then both $\sum a_n$ and $\sum b_n$ converge or diverge together.



Method 7: The Limit Comparison Test

Theorem (Theorem 11.)

Suppose a_n and b_n are positive for all $n \geq N$.

- (a) If $\lim \frac{a_n}{b_n} = c$ (finite and non zero), then both $\sum a_n$ and $\sum b_n$ converge or diverge together.*
- (b) If $\lim \frac{a_n}{b_n} = 0$, and $\sum b_n$ converges, then $\sum a_n$ also converges.*



Method 7: The Limit Comparison Test

Theorem (Theorem 11.)

Suppose a_n and b_n are positive for all $n \geq N$.

- (a) If $\lim \frac{a_n}{b_n} = c$ (finite and non zero), then both $\sum a_n$ and $\sum b_n$ converge or diverge together.*
- (b) If $\lim \frac{a_n}{b_n} = 0$, and $\sum b_n$ converges, then $\sum a_n$ also converges.*
- (c) If $\lim \frac{a_n}{b_n} = \infty$, and $\sum b_n$ diverges, then $\sum a_n$ also diverges.*



Q.: $\sum \frac{1}{n^3 - 5n}$



Q.: $\sum \frac{1}{n^3-5n}$

Sol. Here, $a_n = \frac{1}{n^3-5n}$. What could be b_n ?



Q.: $\sum \frac{1}{n^3-5n}$

Sol. Here, $a_n = \frac{1}{n^3-5n}$. What could be b_n ?

Let $b_n = \frac{1}{n^3}$.



Q.: $\sum \frac{1}{n^3-5n}$

Sol. Here, $a_n = \frac{1}{n^3-5n}$. What could be b_n ?

Let $b_n = \frac{1}{n^3}$.

Now, $\lim \frac{a_n}{b_n} = \lim \frac{n^3}{n^3-5n} = \lim \frac{1}{1-\frac{5}{n^2}} = 1 > 0$.



Q.: $\sum \frac{1}{n^3-5n}$

Sol. Here, $a_n = \frac{1}{n^3-5n}$. What could be b_n ?

Let $b_n = \frac{1}{n^3}$.

Now, $\lim \frac{a_n}{b_n} = \lim \frac{n^3}{n^3-5n} = \lim \frac{1}{1-\frac{5}{n^2}} = 1 > 0$.

From p -test, $\sum \frac{1}{n^3}$ is convergent. Therefore, by LCT, $\sum \frac{1}{n^3-5n}$ is convergent.



Q.: $\sum \frac{n+1}{n^p}$



Q.: $\sum \frac{n+1}{n^p}$

Sol. What could be b_n ?



Q.: $\sum \frac{n+1}{n^p}$

Sol. What could be b_n ?

Here, $a_n = \frac{n+1}{n^p} = \frac{1+\frac{1}{n}}{n^{p-1}}$.



Q.: $\sum \frac{n+1}{n^p}$

Sol. What could be b_n ?

Here, $a_n = \frac{n+1}{n^p} = \frac{1+\frac{1}{n}}{n^{p-1}}$. Let $b_n = \frac{1}{n^{p-1}}$.



Q.: $\sum \frac{n+1}{n^p}$

Sol. What could be b_n ?

Here, $a_n = \frac{n+1}{n^p} = \frac{1+\frac{1}{n}}{n^{p-1}}$. Let $b_n = \frac{1}{n^{p-1}}$.

Now, $\lim \frac{a_n}{b_n} = \lim 1 + \frac{1}{n} = 1 > 0$.



Q.: $\sum \frac{n+1}{n^p}$

Sol. What could be b_n ?

Here, $a_n = \frac{n+1}{n^p} = \frac{1+\frac{1}{n}}{n^{p-1}}$. Let $b_n = \frac{1}{n^{p-1}}$.

Now, $\lim \frac{a_n}{b_n} = \lim 1 + \frac{1}{n} = 1 > 0$.

From p -test, $\sum b_n$ is convergent if $p-1 > 1$ and divergent if $p-1 \leq 1$.



Q: $\sum (\sqrt{n^4 + 1} - \sqrt{n^4 - 1})$



Q.: $\sum(\sqrt{n^4+1} - \sqrt{n^4-1})$

Sol. Here

$$\begin{aligned}a_n &= \sqrt{n^4+1} - \sqrt{n^4-1} \\&= \sqrt{n^4+1} - \sqrt{n^4-1} \cdot \frac{\sqrt{n^4+1} + \sqrt{n^4-1}}{\sqrt{n^4+1} + \sqrt{n^4-1}} \\&= \frac{2}{\sqrt{n^4+1} + \sqrt{n^4-1}}.\end{aligned}$$



Q.: $\sum (\sqrt{n^4 + 1} - \sqrt{n^4 - 1})$

Sol. Here

$$\begin{aligned} a_n &= \sqrt{n^4 + 1} - \sqrt{n^4 - 1} \\ &= \sqrt{n^4 + 1} - \sqrt{n^4 - 1} \cdot \frac{\sqrt{n^4 + 1} + \sqrt{n^4 - 1}}{\sqrt{n^4 + 1} + \sqrt{n^4 - 1}} \\ &= \frac{2}{\sqrt{n^4 + 1} + \sqrt{n^4 - 1}}. \end{aligned}$$

b_n ?



Consider $b_n = \frac{1}{n^2}$.



Consider $b_n = \frac{1}{n^2}$. Note that $\sum b_n$ is convergent. Now



Consider $b_n = \frac{1}{n^2}$. Note that $\sum b_n$ is convergent. Now

$$\lim \frac{a_n}{b_n} = \lim \frac{\frac{2}{\sqrt{n^4+1} + \sqrt{n^4-1}}}{\frac{1}{n^2}} = 1.$$



Consider $b_n = \frac{1}{n^2}$. Note that $\sum b_n$ is convergent. Now

$$\lim \frac{a_n}{b_n} = \lim \frac{\frac{2}{\sqrt{n^4+1} + \sqrt{n^4-1}}}{\frac{1}{n^2}} = 1.$$

Hence $\sum a_n$ is convergent, by LCT (a).



Q: $\sum \tan \frac{1}{n}$



Q:. $\sum \tan \frac{1}{n}$

Sol. Let $b_n = \frac{1}{n} \implies \lim_{n \rightarrow \infty} \frac{\tan \frac{1}{n}}{\frac{1}{n}} = \lim_{n \rightarrow \infty} \frac{1}{\cos \frac{1}{n}} \cdot \frac{\sin \frac{1}{n}}{\frac{1}{n}} =$



Q:. $\sum \tan \frac{1}{n}$

Sol. Let $b_n = \frac{1}{n} \implies \lim_{n \rightarrow \infty} \frac{\tan \frac{1}{n}}{\frac{1}{n}} = \lim_{n \rightarrow \infty} \frac{1}{\cos \frac{1}{n}} \cdot \frac{\sin \frac{1}{n}}{\frac{1}{n}} =$
 $\lim_{x \rightarrow 0} \frac{1}{\cos x} \cdot \frac{\sin x}{x} = 1 \cdot 1 = 1.$



Q:. $\sum \tan \frac{1}{n}$

Sol. Let $b_n = \frac{1}{n} \implies \lim_{n \rightarrow \infty} \frac{\tan \frac{1}{n}}{\frac{1}{n}} = \lim_{n \rightarrow \infty} \frac{1}{\cos \frac{1}{n}} \cdot \frac{\sin \frac{1}{n}}{\frac{1}{n}} =$

$$\lim_{x \rightarrow 0} \frac{1}{\cos x} \cdot \frac{\sin x}{x} = 1 \cdot 1 = 1.$$

Now, $\sum b_n$ is a divergent p -series, hence, $\sum a_n$ is divergent, by LCT.



Q: $\sum \frac{(1+n \ln n)}{n^2+5}.$



Q.: $\sum \frac{(1+n \ln n)}{n^2+5}$.

Sol. Here $a_n = \frac{(1+n \ln n)}{n^2+5}$.



Q.: $\sum \frac{(1+n \ln n)}{n^2+5}$.

Sol. Here $a_n = \frac{(1+n \ln n)}{n^2+5}$. For large n , a_n would behave like $\frac{n \ln n}{n^2} = \frac{\ln n}{n}$. Consider $b_n = \frac{1}{n}$.



Q.: $\sum \frac{(1+n \ln n)}{n^2+5}$.

Sol. Here $a_n = \frac{(1+n \ln n)}{n^2+5}$. For large n , a_n would behave like $\frac{n \ln n}{n^2} = \frac{\ln n}{n}$. Consider $b_n = \frac{1}{n}$. Note that $\sum b_n$ is divergent. Now

$$\lim \frac{a_n}{b_n} = \lim \frac{n^2 \ln n + n}{n^2 + 5} = \infty$$

Hence $\sum a_n$ is divergent, by LCT.



Q:. $\sum \frac{1}{\sqrt{n^3+2}}.$

Sol. Convergent.

Q:. $\sum \frac{2^n-n}{n2^n}$

Sol. Divergent

Q:. $\sum_{n=2}^{\infty} \frac{1}{\sqrt{n} \ln n}.$

Sol. Divergent



Section 10.5

Ratio and Root Tests



Method 8: Ratio Test

Theorem (Theorem 12.)

Let $\sum a_n$ be a series of positive terms. Suppose $\lim \frac{a_{n+1}}{a_n} = r$.



Method 8: Ratio Test

Theorem (Theorem 12.)

Let $\sum a_n$ be a series of positive terms. Suppose $\lim \frac{a_{n+1}}{a_n} = r$.

(a) If $r < 1$, the series $\sum a_n$ is convergent;



Method 8: Ratio Test

Theorem (Theorem 12.)

Let $\sum a_n$ be a series of positive terms. Suppose $\lim \frac{a_{n+1}}{a_n} = r$.

(a) If $r < 1$, the series $\sum a_n$ is convergent;

(b) If $r > 1$, the series $\sum a_n$ is divergent;



Method 8: Ratio Test

Theorem (Theorem 12.)

Let $\sum a_n$ be a series of positive terms. Suppose $\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = r$.

- (a) If $r < 1$, the series $\sum a_n$ is convergent;*
- (b) If $r > 1$, the series $\sum a_n$ is divergent;*
- (c) If $r = 1$, the test is inconclusive.*



Q: $\sum \frac{n}{2^n}.$



Q.: $\sum \frac{n}{2^n}$.

Sol. Here $a_n = \frac{n}{2^n}$. So



Q.: $\sum \frac{n}{2^n}$.

Sol. Here $a_n = \frac{n}{2^n}$. So

$$\frac{a_{n+1}}{a_n} = \frac{\frac{n+1}{2^{n+1}}}{\frac{n}{2^n}} = \frac{n+1}{2^{n+1}} \cdot \frac{2^n}{n}$$
$$\lim \frac{a_{n+1}}{a_n} = \lim \frac{1}{2} \cdot \frac{n+1}{n} = \frac{1}{2} < 1.$$



Q.: $\sum \frac{n}{2^n}$.

Sol. Here $a_n = \frac{n}{2^n}$. So

$$\frac{a_{n+1}}{a_n} = \frac{\frac{n+1}{2^{n+1}}}{\frac{n}{2^n}} = \frac{n+1}{2^{n+1}} \cdot \frac{2^n}{n}$$
$$\lim \frac{a_{n+1}}{a_n} = \lim \frac{1}{2} \cdot \frac{n+1}{n} = \frac{1}{2} < 1.$$

Therefore by ratio test, the series converges.



Q: $\sum e^{-n} n^3.$



Q:. $\sum e^{-n}n^3$.

Sol. Here $a_n = e^{-n}n^3$. So



Q.: $\sum e^{-n}n^3$.

Sol. Here $a_n = e^{-n}n^3$. So

$$\frac{a_{n+1}}{a_n} = \frac{e^{-(n+1)}(n+1)^3}{e^{-n}n^3} = e^{-1} \left(1 + \frac{1}{n}\right)^3.$$

Thus $\lim \frac{a_{n+1}}{a_n} = e^{-1} < 1$. Therefore by ratio test, the series converges.



Q.: $\sum \frac{1}{n^2}$.



Q.: $\sum \frac{1}{n^2}$.

Sol. Here $a_n = \frac{1}{n^2}$. So



Q.: $\sum \frac{1}{n^2}$.

Sol. Here $a_n = \frac{1}{n^2}$. So

$$\frac{a_{n+1}}{a_n} = \frac{n^2}{(n+1)^2}$$
$$\lim \frac{a_{n+1}}{a_n} = 1.$$



Q.: $\sum \frac{1}{n^2}$.

Sol. Here $a_n = \frac{1}{n^2}$. So

$$\frac{a_{n+1}}{a_n} = \frac{n^2}{(n+1)^2}$$
$$\lim \frac{a_{n+1}}{a_n} = 1.$$

Therefore the ratio test yields no conclusion.



Q.: $\sum \frac{3^n}{n^3 2^n}.$



Q.: $\sum \frac{3^n}{n^3 2^n}.$

Sol. Divergent.

Q.: $\sum \frac{(2n+3)(2^n+3)}{3^{n+2}}.$

Sol. Convergent.



Method 9: Root Test or n^{th} Root Test

Theorem (Theorem 13.)

Let $\sum a_n$ be a series of non-negative terms, and suppose that $\lim(a_n)^{\frac{1}{n}} = r$.



Method 9: Root Test or n^{th} Root Test

Theorem (Theorem 13.)

Let $\sum a_n$ be a series of non-negative terms, and suppose that $\lim(a_n)^{\frac{1}{n}} = r$.

(a) If $r < 1$, the series $\sum a_n$ converges;



Method 9: Root Test or n^{th} Root Test

Theorem (Theorem 13.)

Let $\sum a_n$ be a series of non-negative terms, and suppose that $\lim(a_n)^{\frac{1}{n}} = r$.

(a) If $r < 1$, the series $\sum a_n$ converges;

(b) If $r > 1$, the series $\sum a_n$ diverges;



Method 9: Root Test or n^{th} Root Test

Theorem (Theorem 13.)

Let $\sum a_n$ be a series of non-negative terms, and suppose that $\lim(a_n)^{\frac{1}{n}} = r$.

- (a) If $r < 1$, the series $\sum a_n$ converges;*
- (b) If $r > 1$, the series $\sum a_n$ diverges;*
- (c) If $r = 1$, the test is inconclusive.*



Q.: $\sum \left(\frac{2n+4}{5n-1} \right)^n$.



Q.: $\sum \left(\frac{2n+4}{5n-1} \right)^n$.

Sol. Here

$$a_n = \left(\frac{2n+4}{5n-1} \right)^n$$

$$\Rightarrow (a_n)^{1/n} = \frac{2n+4}{5n-1}$$

$$\Rightarrow \lim(a_n)^{1/n} = \frac{2}{5} < 1.$$



Q.: $\sum \left(\frac{2n+4}{5n-1}\right)^n$.

Sol. Here

$$a_n = \left(\frac{2n+4}{5n-1}\right)^n$$

$$\Rightarrow (a_n)^{1/n} = \frac{2n+4}{5n-1}$$

$$\Rightarrow \lim(a_n)^{1/n} = \frac{2}{5} < 1.$$

Therefore the series converges by root test.



Q.: $\sum \frac{(n!)^n}{(n^n)^2}.$



Q.: $\sum \frac{(n!)^n}{(n^n)^2}$.

Sol. Here

$$a_n = \frac{(n!)^n}{(n^n)^2}$$

$$(a_n)^{1/n} = \frac{n!}{n^2}$$

$$\lim (a_n)^{1/n} = \lim \frac{n!}{n^2} = \infty.$$

Therefore the series diverges by root test.



Section 10.6

Alternating Series, Absolute and Conditional Convergence



Alternating Series

A series whose terms are alternately positive and negative is called an **alternating series**.



Alternating Series

A series whose terms are alternately positive and negative is called an **alternating series**.

An alternating series is one of the form $\sum (-1)^{n+1} u_n$ or $\sum (-1)^n u_n$, where $u_n > 0$ for all n .



Examples: Alternating series

- $\sum (-1)^{n+1}$



Examples: Alternating series

- $\sum (-1)^{n+1}$
- $\sum (-1)^n \frac{1}{n \ln n}$



Examples: Alternating series

- $\sum (-1)^{n+1}$
- $\sum (-1)^n \frac{1}{n \ln n}$
- $\sum (-1)^n \ln \left(1 + \frac{1}{n}\right)$



Examples: Alternating series

- $\sum (-1)^{n+1}$
- $\sum (-1)^n \frac{1}{n \ln n}$
- $\sum (-1)^n \ln \left(1 + \frac{1}{n} \right)$
- $\sum \frac{\cos n\pi}{n^2+1}$



Theorem (Theorem 14. Leibniz's Theorem for Alternating Series)

The alternating series $\sum (-1)^{n+1} u_n$ converges if



Theorem (Theorem 14. Leibniz's Theorem for Alternating Series)

The alternating series $\sum (-1)^{n+1} u_n$ converges if

(i) $u_n \geq u_{n+1}$ for all $n \geq N$ for some N ; and



Theorem (Theorem 14. Leibniz's Theorem for Alternating Series)

The alternating series $\sum (-1)^{n+1} u_n$ converges if

- (i) $u_n \geq u_{n+1}$ for all $n \geq N$ for some N ; and*
- (ii) $\lim u_n = 0$.*



Q: $\sum (-1)^{n+1} \frac{1}{(2n-1)!}$



Q:. $\sum (-1)^{n+1} \frac{1}{(2n-1)!}$

Sol. (i) Here $u_n = \frac{1}{(2n-1)!}$. It is easy to verify that $u_n \geq u_{n+1}$ for all $n \geq 1$.



Q:. $\sum (-1)^{n+1} \frac{1}{(2n-1)!}$

Sol. (i) Here $u_n = \frac{1}{(2n-1)!}$. It is easy to verify that $u_n \geq u_{n+1}$ for all $n \geq 1$.

(ii) Also, we have $\lim u_n = 0$.



Q:. $\sum (-1)^{n+1} \frac{1}{(2n-1)!}$

Sol. (i) Here $u_n = \frac{1}{(2n-1)!}$. It is easy to verify that $u_n \geq u_{n+1}$ for all $n \geq 1$.

(ii) Also, we have $\lim u_n = 0$.

Hence by Leibniz's test the series is convergent.



Proof. To prove Leibniz Theorem, we will show that the sequence S_n is convergent, which is proved using the following result:



Proof. To prove Leibniz Theorem, we will show that the sequence S_n is convergent, which is proved using the following result:

Ex. 133 on p. 572: For a sequence $\{S_n\}$, if $\{S_{2n}\}$ and $\{S_{2n+1}\}$ converge to the same number L , then $S_n \rightarrow L$.



Proof. To prove Leibniz Theorem, we will show that the sequence S_n is convergent, which is proved using the following result:

Ex. 133 on p. 572: For a sequence $\{S_n\}$, if $\{S_{2n}\}$ and $\{S_{2n+1}\}$ converge to the same number L , then $S_n \rightarrow L$.

First consider the sequence $\{S_{2n}\}$.



Proof. To prove Leibniz Theorem, we will show that the sequence S_n is convergent, which is proved using the following result:

Ex. 133 on p. 572: For a sequence $\{S_n\}$, if $\{S_{2n}\}$ and $\{S_{2n+1}\}$ converge to the same number L , then $S_n \rightarrow L$.

First consider the sequence $\{S_{2n}\}$. We'll show that S_{2n} converges (by showing that S_{2n} is non-decreasing and bounded from above). **Why ?**



S_{2n} is non-decreasing. We have



S_{2n} is non-decreasing. We have

$$S_{2n+2} = S_{2n} + (u_{2n+1} - u_{2n+2}).$$

Since $u_{2n+1} - u_{2n+2} \geq 0$ Why ?



S_{2n} is non-decreasing. We have

$$S_{2n+2} = S_{2n} + (u_{2n+1} - u_{2n+2}).$$

Since $u_{2n+1} - u_{2n+2} \geq 0$ **Why ?** $\implies S_{2n+2} \geq S_{2n}$. Thus the sequence $\{S_{2n}\}$ is non-decreasing.



S_{2n} is non-decreasing. We have

$$S_{2n+2} = S_{2n} + (u_{2n+1} - u_{2n+2}).$$

Since $u_{2n+1} - u_{2n+2} \geq 0$ **Why ?** $\implies S_{2n+2} \geq S_{2n}$. Thus the sequence $\{S_{2n}\}$ is non-decreasing.

S_{2n} is bounded from above.



S_{2n} is non-decreasing. We have

$$S_{2n+2} = S_{2n} + (u_{2n+1} - u_{2n+2}).$$

Since $u_{2n+1} - u_{2n+2} \geq 0$ **Why ?** $\implies S_{2n+2} \geq S_{2n}$. Thus the sequence $\{S_{2n}\}$ is non-decreasing.

S_{2n} is bounded from above. Arrange S_{2n} as

$$S_{2n} = u_1 - (u_2 - u_3) - \cdots - (u_{2n-2} - u_{2n-1}) - u_{2n} \leq$$



S_{2n} is non-decreasing. We have

$$S_{2n+2} = S_{2n} + (u_{2n+1} - u_{2n+2}).$$

Since $u_{2n+1} - u_{2n+2} \geq 0$ **Why ?** $\implies S_{2n+2} \geq S_{2n}$. Thus the sequence $\{S_{2n}\}$ is non-decreasing.

S_{2n} is bounded from above. Arrange S_{2n} as
$$S_{2n} = u_1 - (u_2 - u_3) - \cdots - (u_{2n-2} - u_{2n-1}) - u_{2n} \leq u_1.$$



S_{2n} is non-decreasing. We have

$$S_{2n+2} = S_{2n} + (u_{2n+1} - u_{2n+2}).$$

Since $u_{2n+1} - u_{2n+2} \geq 0$ **Why ?** $\implies S_{2n+2} \geq S_{2n}$. Thus the sequence $\{S_{2n}\}$ is non-decreasing.

S_{2n} is bounded from above. Arrange S_{2n} as
 $S_{2n} = u_1 - (u_2 - u_3) - \cdots - (u_{2n-2} - u_{2n-1}) - u_{2n} \leq u_1$.

Thus the sequence $\{S_{2n}\}$ is bounded from above.



Therefore it is convergent and has a limit, say L
i.e., $\lim S_{2n} \rightarrow L$.



Therefore it is convergent and has a limit, say L
i.e., $\lim S_{2n} \rightarrow L$.

Now consider the sequence $\{S_{2n+1}\}$. We have



Therefore it is convergent and has a limit, say L
i.e., $\lim S_{2n} \rightarrow L$.

Now consider the sequence $\{S_{2n+1}\}$. We have

$$\begin{aligned} S_{2n+1} &= S_{2n} + u_{2n+1} \\ \Rightarrow \lim S_{2n+1} &= \lim S_{2n} + \lim u_{2n+1}. \end{aligned}$$



Therefore it is convergent and has a limit, say L
i.e., $\lim S_{2n} \rightarrow L$.

Now consider the sequence $\{S_{2n+1}\}$. We have

$$\begin{aligned} S_{2n+1} &= S_{2n} + u_{2n+1} \\ \Rightarrow \lim S_{2n+1} &= \lim S_{2n} + \lim u_{2n+1}. \end{aligned}$$

Therefore, using condition (ii):



Therefore it is convergent and has a limit, say L
i.e., $\lim S_{2n} \rightarrow L$.

Now consider the sequence $\{S_{2n+1}\}$. We have

$$\begin{aligned} S_{2n+1} &= S_{2n} + u_{2n+1} \\ \Rightarrow \lim S_{2n+1} &= \lim S_{2n} + \lim u_{2n+1}. \end{aligned}$$

Therefore, using condition (ii):

$$\lim S_{2n+1} = L + 0 = L.$$



Therefore it is convergent and has a limit, say L
i.e., $\lim S_{2n} \rightarrow L$.

Now consider the sequence $\{S_{2n+1}\}$. We have

$$\begin{aligned} S_{2n+1} &= S_{2n} + u_{2n+1} \\ \Rightarrow \lim S_{2n+1} &= \lim S_{2n} + \lim u_{2n+1}. \end{aligned}$$

Therefore, using condition (ii):

$$\lim S_{2n+1} = L + 0 = L.$$

Thus $S_n \rightarrow L$ and hence $\sum (-1)^{n+1} u_n$ converges to L .



Remark 1

If second condition of Leibniz's test fails, i.e., $\lim u_n \neq 0$, then the series would be divergent.



Remark 1

If second condition of Leibniz's test fails, i.e., $\lim u_n \neq 0$, then the series would be divergent.

Q.: $\sum (-1)^{n+1} \frac{2^n}{n^2}$.



Remark 1

If second condition of Leibniz's test fails, i.e., $\lim u_n \neq 0$, then the series would be divergent.

Q.: $\sum (-1)^{n+1} \frac{2^n}{n^2}$.

Sol. Here $u_n = \frac{2^n}{n^2}$ and $\lim u_n = \infty \implies$ the series is divergent.



Remark 2

The Leibniz's test is sufficient but not necessary for the convergence of an alternating series. There are examples of alternating series for which condition (i) fails but the series converges. Thus, if the first condition of Leibniz's test fails, the result would be inconclusive. In this case, we can not say that the series is divergent as it may be convergent.



Remark 2

The Leibniz's test is sufficient but not necessary for the convergence of an alternating series. There are examples of alternating series for which condition (i) fails but the series converges. Thus, if the first condition of Leibniz's test fails, the result would be inconclusive. In this case, we can not say that the series is divergent as it may be convergent.

Q.: $\sum (-1)^n u_n = -1 + \frac{1}{8} - \frac{1}{9} + \frac{1}{64} - \frac{1}{25} + \dots$



Remark 2

The Leibniz's test is sufficient but not necessary for the convergence of an alternating series. There are examples of alternating series for which condition (i) fails but the series converges. Thus, if the first condition of Leibniz's test fails, the result would be inconclusive. In this case, we can not say that the series is divergent as it may be convergent.

Q.: $\sum (-1)^n u_n = -1 + \frac{1}{8} - \frac{1}{9} + \frac{1}{64} - \frac{1}{25} + \dots$

Sol. Clearly, the series is not monotonically decreasing but still it is convergent.



Remark 2

The Leibniz's test is sufficient but not necessary for the convergence of an alternating series. There are examples of alternating series for which condition (i) fails but the series converges. Thus, if the first condition of Leibniz's test fails, the result would be inconclusive. In this case, we can not say that the series is divergent as it may be convergent.

Q.: $\sum (-1)^n u_n = -1 + \frac{1}{8} - \frac{1}{9} + \frac{1}{64} - \frac{1}{25} + \dots$

Sol. Clearly, the series is not monotonically decreasing but still it is convergent. **how?**



Remark 2

The Leibniz's test is sufficient but not necessary for the convergence of an alternating series. There are examples of alternating series for which condition (i) fails but the series converges. Thus, if the first condition of Leibniz's test fails, the result would be inconclusive. In this case, we can not say that the series is divergent as it may be convergent.

Q.: $\sum (-1)^n u_n = -1 + \frac{1}{8} - \frac{1}{9} + \frac{1}{64} - \frac{1}{25} + \dots$

Sol. Clearly, the series is not monotonically decreasing but still it is convergent. **how?** (difference of geometric series)



Q.: $\sum (-1)^{n+1} \frac{\sqrt{n}}{n+1}.$



Q.: $\sum (-1)^{n+1} \frac{\sqrt{n}}{n+1}$.

Sol. (i) Here $u_n = \frac{\sqrt{n}}{n+1}$. Is $u_n \geq u_{n+1}$?



Q.: $\sum (-1)^{n+1} \frac{\sqrt{n}}{n+1}$.

Sol. (i) Here $u_n = \frac{\sqrt{n}}{n+1}$. Is $u_n \geq u_{n+1}$?

Consider

$$f(x) = \frac{\sqrt{x}}{x+1} \Rightarrow f'(x) = -\frac{x-1}{2\sqrt{x}(x+1)^2} < 0 \forall x > 1.$$

Hence, $f(x)$ is non-increasing for all

$$x > 1 \Rightarrow u_n \geq u_{n+1}, \forall n > 1.$$



Q.: $\sum (-1)^{n+1} \frac{\sqrt{n}}{n+1}$.

Sol. (i) Here $u_n = \frac{\sqrt{n}}{n+1}$. Is $u_n \geq u_{n+1}$?

Consider

$$f(x) = \frac{\sqrt{x}}{x+1} \Rightarrow f'(x) = -\frac{x-1}{2\sqrt{x}(x+1)^2} < 0 \forall x > 1.$$

Hence, $f(x)$ is non-increasing for all $x > 1 \Rightarrow u_n \geq u_{n+1}, \forall n > 1$.

(ii) Also, we have $\lim u_n = 0$ (how?).



Q.: $\sum (-1)^{n+1} \frac{\sqrt{n}}{n+1}$.

Sol. (i) Here $u_n = \frac{\sqrt{n}}{n+1}$. Is $u_n \geq u_{n+1}$?

Consider

$$f(x) = \frac{\sqrt{x}}{x+1} \Rightarrow f'(x) = -\frac{x-1}{2\sqrt{x}(x+1)^2} < 0 \forall x > 1.$$

Hence, $f(x)$ is non-increasing for all

$$x > 1 \Rightarrow u_n \geq u_{n+1}, \forall n > 1.$$

(ii) Also, we have $\lim u_n = 0$ (how?). (Use L'Hôpital Rule)



Q.: $\sum (-1)^{n+1} \frac{\sqrt{n}}{n+1}$.

Sol. (i) Here $u_n = \frac{\sqrt{n}}{n+1}$. Is $u_n \geq u_{n+1}$?

Consider

$$f(x) = \frac{\sqrt{x}}{x+1} \Rightarrow f'(x) = -\frac{x-1}{2\sqrt{x}(x+1)^2} < 0 \forall x > 1.$$

Hence, $f(x)$ is non-increasing for all

$$x > 1 \Rightarrow u_n \geq u_{n+1}, \forall n > 1.$$

(ii) Also, we have $\lim u_n = 0$ (how?). (Use L'Hôpital Rule)

Hence by Leibniz's test the series is convergent.



Q:. $\sum (-1)^n \ln\left(1 + \frac{1}{n}\right).$

Sol. Convergent



Absolute and Conditional Convergence

A series $\sum a_n$ is said to be

- **absolutely convergent** if the series $\sum |a_n|$ is convergent.



Absolute and Conditional Convergence

A series $\sum a_n$ is said to be

- **absolutely convergent** if the series $\sum |a_n|$ is convergent.
- **conditionally convergent** if the series $\sum |a_n|$ is divergent but $\sum a_n$ is convergent.



Example: Absolute and Conditional Convergence

- $\sum (-1)^n \frac{1}{2^n}$



Example: Absolute and Conditional Convergence

- $\sum (-1)^n \frac{1}{2^n}$ (absolutely convergent)



Example: Absolute and Conditional Convergence

- $\sum (-1)^n \frac{1}{2^n}$ (absolutely convergent)
- $\sum (-1)^n \frac{1}{n}$



Example: Absolute and Conditional Convergence

- $\sum (-1)^n \frac{1}{2^n}$ (absolutely convergent)
- $\sum (-1)^n \frac{1}{n}$ (conditionally convergent)



Example: Absolute and Conditional Convergence

- $\sum (-1)^n \frac{1}{2^n}$ (absolutely convergent)
- $\sum (-1)^n \frac{1}{n}$ (conditionally convergent)
- $\sum (-1)^{n+1} \frac{3+n}{5+n}$



Example: Absolute and Conditional Convergence

- $\sum (-1)^n \frac{1}{2^n}$ (absolutely convergent)
- $\sum (-1)^n \frac{1}{n}$ (conditionally convergent)
- $\sum (-1)^{n+1} \frac{3+n}{5+n}$ (divergent)



Theorem (Theorem 16. The Absolute Convergence Test)

If $\sum |a_n|$ converges, then $\sum a_n$ also converges. That is “absolute convergence implies convergence”.



Theorem (Theorem 16. The Absolute Convergence Test)

If $\sum |a_n|$ converges, then $\sum a_n$ also converges. That is “absolute convergence implies convergence”.

Proof. We have

$$-|a_n| \leq a_n \leq |a_n|, \quad \forall n.$$



Theorem (Theorem 16. The Absolute Convergence Test)

If $\sum |a_n|$ converges, then $\sum a_n$ also converges. That is “absolute convergence implies convergence”.

Proof. We have

$$-|a_n| \leq a_n \leq |a_n|, \quad \forall n.$$

$$\implies 0 \leq |a_n| + a_n \leq 2|a_n|.$$



Theorem (Theorem 16. The Absolute Convergence Test)

If $\sum |a_n|$ converges, then $\sum a_n$ also converges. That is “absolute convergence implies convergence”.

Proof. We have

$$-|a_n| \leq a_n \leq |a_n|, \quad \forall n.$$

$$\implies 0 \leq |a_n| + a_n \leq 2|a_n|.$$

Now the series $\sum 2|a_n|$ converges as the series $\sum |a_n|$



Therefore by Direct Comparison Test, the non-negative terms series $\sum(|a_n| + a_n)$ converges.



Therefore by Direct Comparison Test, the non-negative terms series $\sum(|a_n| + a_n)$ converges.

Now we can write

$$\sum a_n = \sum (|a_n| + a_n - |a_n|) = \sum (|a_n| + a_n) - \sum |a_n|.$$



Therefore by Direct Comparison Test, the non-negative terms series $\sum(|a_n| + a_n)$ converges.

Now we can write

$$\sum a_n = \sum (|a_n| + a_n - |a_n|) = \sum (|a_n| + a_n) - \sum |a_n|.$$

Therefore $\sum a_n$, being a difference of two convergent series, converges. □



Steps to check a series for absolute and conditional convergence

- First check the convergence of the series $\sum |a_n|$, i.e., if $\sum |a_n|$ is convergent then $\sum a_n$ is absolutely convergent.



Steps to check a series for absolute and conditional convergence

- First check the convergence of the series $\sum |a_n|$, i.e., if $\sum |a_n|$ is convergent then $\sum a_n$ is absolutely convergent.
- If $\sum |a_n|$ is divergent but $\sum a_n$ is convergent then $\sum a_n$ is conditionally convergent.



Steps to check a series for absolute and conditional convergence

- First check the convergence of the series $\sum |a_n|$, i.e., if $\sum |a_n|$ is convergent then $\sum a_n$ is absolutely convergent.
- If $\sum |a_n|$ is divergent but $\sum a_n$ is convergent then $\sum a_n$ is conditionally convergent.
- Otherwise $\sum a_n$ is divergent.



Q: $\sum (-1)^{n+1} \frac{\sin nx}{n^3}.$



Q:. $\sum (-1)^{n+1} \frac{\sin nx}{n^3}$.

Sol. Here $|a_n| = \frac{|\sin nx|}{n^3}$.



Q:. $\sum (-1)^{n+1} \frac{\sin nx}{n^3}$.

Sol. Here $|a_n| = \frac{|\sin nx|}{n^3}$. Let $b_n = \frac{1}{n^3}$.



Q:. $\sum (-1)^{n+1} \frac{\sin nx}{n^3}$.

Sol. Here $|a_n| = \frac{|\sin nx|}{n^3}$. Let $b_n = \frac{1}{n^3}$. Clearly, $\frac{|\sin nx|}{n^3} \leq \frac{1}{n^3}$ and $\sum \frac{1}{n^3}$ is convergent



Q.: $\sum (-1)^{n+1} \frac{\sin nx}{n^3}$.

Sol. Here $|a_n| = \frac{|\sin nx|}{n^3}$. Let $b_n = \frac{1}{n^3}$. Clearly, $\frac{|\sin nx|}{n^3} \leq \frac{1}{n^3}$ and $\sum \frac{1}{n^3}$ is convergent (from p-series test). Hence $\sum \frac{|\sin nx|}{n^3}$ is convergent.



Q.: $\sum (-1)^{n+1} \frac{\sin nx}{n^3}$.

Sol. Here $|a_n| = \frac{|\sin nx|}{n^3}$. Let $b_n = \frac{1}{n^3}$. Clearly, $\frac{|\sin nx|}{n^3} \leq \frac{1}{n^3}$ and $\sum \frac{1}{n^3}$ is convergent (from p-series test). Hence $\sum \frac{|\sin nx|}{n^3}$ is convergent. This implies that $\sum (-1)^{n+1} \frac{\sin nx}{n^3}$ is absolutely convergent and hence, convergent.



Q: $\sum (-1)^{n+1} \frac{n}{n^2+1}.$



Q: $\sum (-1)^{n+1} \frac{n}{n^2+1}$.

Sol.

- First we check the behavior of $\sum |a_n| = \sum \frac{n}{n^2+1}$.



Q.: $\sum (-1)^{n+1} \frac{n}{n^2+1}$.

Sol.

- First we check the behavior of $\sum |a_n| = \sum \frac{n}{n^2+1}$. Let $b_n = \frac{1}{n}$.

Now $\lim \frac{|a_n|}{b_n} = \lim \frac{n^2}{n^2+1} = 1$.



Q.: $\sum (-1)^{n+1} \frac{n}{n^2+1}$.

Sol.

- First we check the behavior of $\sum |a_n| = \sum \frac{n}{n^2+1}$. Let $b_n = \frac{1}{n}$.

Now $\lim \frac{|a_n|}{b_n} = \lim \frac{n^2}{n^2+1} = 1$.

Since $\sum \frac{1}{n}$ is divergent so $\sum \frac{n}{n^2+1}$ is also divergent (by LCT).



- Now we check the behavior of $\sum a_n = \sum (-1)^{n+1} \frac{n}{n^2+1}$ by using Leibniz's test.



- Now we check the behavior of $\sum a_n = \sum (-1)^{n+1} \frac{n}{n^2+1}$ by using Leibniz's test.

(i) Here, $u_{n+1} - u_n = \frac{-(n^2+n)+1}{(n^2+1)[(n+1)^2+1]} < 0$ for all n .



- Now we check the behavior of $\sum a_n = \sum (-1)^{n+1} \frac{n}{n^2+1}$ by using Leibniz's test.

(i) Here, $u_{n+1} - u_n = \frac{-(n^2+n)+1}{(n^2+1)[(n+1)^2+1]} < 0$ for all n .

(ii) Also we have $\lim_{n \rightarrow \infty} \frac{n}{n^2+1} = \lim_{n \rightarrow \infty} \frac{1}{n+1/n} = 0$



- Now we check the behavior of $\sum a_n = \sum (-1)^{n+1} \frac{n}{n^2+1}$ by using Leibniz's test.

(i) Here, $u_{n+1} - u_n = \frac{-(n^2+n)+1}{(n^2+1)[(n+1)^2+1]} < 0$ for all n .

(ii) Also we have $\lim \frac{n}{n^2+1} = \lim \frac{1}{n+1/n} = 0$

Therefore by Leibniz's test $\sum (-1)^{n+1} \frac{n}{n^2+1}$ is convergent.



- Now we check the behavior of $\sum a_n = \sum (-1)^{n+1} \frac{n}{n^2+1}$ by using Leibniz's test.

(i) Here, $u_{n+1} - u_n = \frac{-(n^2+n)+1}{(n^2+1)[(n+1)^2+1]} < 0$ for all n .

(ii) Also we have $\lim_{n \rightarrow \infty} \frac{n}{n^2+1} = \lim_{n \rightarrow \infty} \frac{1}{n+1/n} = 0$

Therefore by Leibniz's test $\sum (-1)^{n+1} \frac{n}{n^2+1}$ is convergent.

Hence $\sum (-1)^{n+1} \frac{n}{n^2+1}$ is conditionally convergent.



Q: $\sum (-1)^{n+1} \frac{(2n)!}{2^n n! n}$



Q:. $\sum (-1)^{n+1} \frac{(2n)!}{2^n n! n}$

Sol. $\lim \frac{(2n)!}{2^n n! n} = \lim \frac{(n+1)(n+2)\dots(2n)}{2^n n} = \lim \frac{(n+1)(n+2)\dots(n+(n-1))}{2^{n-1}}$



Q:. $\sum (-1)^{n+1} \frac{(2n)!}{2^n n! n}$

Sol. $\lim \frac{(2n)!}{2^n n! n} = \lim \frac{(n+1)(n+2)\dots(2n)}{2^n n} = \lim \frac{(n+1)(n+2)\dots(n+(n-1))}{2^{n-1}}$

$> \lim \left(\frac{n+1}{2}\right)^{n-1} = \infty \neq 0$



Q:. $\sum (-1)^{n+1} \frac{(2n)!}{2^n n! n}$

Sol. $\lim \frac{(2n)!}{2^n n! n} = \lim \frac{(n+1)(n+2)\dots(2n)}{2^n n} = \lim \frac{(n+1)(n+2)\dots(n+(n-1))}{2^{n-1}}$

$> \lim \left(\frac{n+1}{2}\right)^{n-1} = \infty \neq 0$

By n^{th} term test, $\sum (-1)^{n+1} \frac{(2n)!}{2^n n! n}$ is divergent.



Q: $\sum (-1)^n \frac{10^n}{(n+1)!}$



Q:. $\sum (-1)^n \frac{10^n}{(n+1)!}$

Sol. Here $|a_n| = \frac{10^n}{(n+1)!}$.



Q:. $\sum (-1)^n \frac{10^n}{(n+1)!}$

Sol. Here $|a_n| = \frac{10^n}{(n+1)!}$.

Now, $\lim \left| \frac{a_{n+1}}{a_n} \right| = \lim \frac{10}{n+2} = 0 < 1$.



Q.: $\sum (-1)^n \frac{10^n}{(n+1)!}$

Sol. Here $|a_n| = \frac{10^n}{(n+1)!}$.

Now, $\lim \left| \frac{a_{n+1}}{a_n} \right| = \lim \frac{10}{n+2} = 0 < 1$. By ratio test, $\sum (-1)^n \frac{10^n}{(n+1)!}$ is absolutely convergent and hence, convergent.



Q.: $\sum (-1)^n (\sqrt{n+1} - \sqrt{n})$



Q:. $\sum (-1)^n (\sqrt{n+1} - \sqrt{n})$

Sol. Now,

$$\sqrt{n+1} - \sqrt{n} = (\sqrt{n+1} - \sqrt{n}) \cdot \frac{\sqrt{n+1} + \sqrt{n}}{\sqrt{n+1} + \sqrt{n}} = \frac{1}{\sqrt{n+1} + \sqrt{n}}.$$



Q:. $\sum (-1)^n (\sqrt{n+1} - \sqrt{n})$

Sol. Now,

$$\sqrt{n+1} - \sqrt{n} = (\sqrt{n+1} - \sqrt{n}) \cdot \frac{\sqrt{n+1} + \sqrt{n}}{\sqrt{n+1} + \sqrt{n}} = \frac{1}{\sqrt{n+1} + \sqrt{n}}.$$

Here $\sum |a_n| = \sum \frac{1}{\sqrt{n+1} + \sqrt{n}}.$



Q:. $\sum (-1)^n (\sqrt{n+1} - \sqrt{n})$

Sol. Now,

$$\sqrt{n+1} - \sqrt{n} = (\sqrt{n+1} - \sqrt{n}) \cdot \frac{\sqrt{n+1} + \sqrt{n}}{\sqrt{n+1} + \sqrt{n}} = \frac{1}{\sqrt{n+1} + \sqrt{n}}.$$

Here $\sum |a_n| = \sum \frac{1}{\sqrt{n+1} + \sqrt{n}}$. Let $b_n = \frac{1}{\sqrt{n}}$.



Q:. $\sum (-1)^n (\sqrt{n+1} - \sqrt{n})$

Sol. Now,

$$\sqrt{n+1} - \sqrt{n} = (\sqrt{n+1} - \sqrt{n}) \cdot \frac{\sqrt{n+1} + \sqrt{n}}{\sqrt{n+1} + \sqrt{n}} = \frac{1}{\sqrt{n+1} + \sqrt{n}}.$$

Here $\sum |a_n| = \sum \frac{1}{\sqrt{n+1} + \sqrt{n}}$. Let $b_n = \frac{1}{\sqrt{n}}$.

Now, $\lim \frac{|a_n|}{b_n} = \lim \frac{\sqrt{n}}{\sqrt{n+1} + \sqrt{n}} = \lim \frac{1}{\sqrt{1 + \frac{1}{n}} + 1} = \frac{1}{2} \neq 0$.



Q:. $\sum (-1)^n (\sqrt{n+1} - \sqrt{n})$

Sol. Now,

$$\sqrt{n+1} - \sqrt{n} = (\sqrt{n+1} - \sqrt{n}) \cdot \frac{\sqrt{n+1} + \sqrt{n}}{\sqrt{n+1} + \sqrt{n}} = \frac{1}{\sqrt{n+1} + \sqrt{n}}.$$

Here $\sum |a_n| = \sum \frac{1}{\sqrt{n+1} + \sqrt{n}}$. Let $b_n = \frac{1}{\sqrt{n}}$.

Now, $\lim \frac{|a_n|}{b_n} = \lim \frac{\sqrt{n}}{\sqrt{n+1} + \sqrt{n}} = \lim \frac{1}{\sqrt{1 + \frac{1}{n}} + 1} = \frac{1}{2} \neq 0$.

Here, $\sum b_n$ is a divergent p -series.



Q.: $\sum (-1)^n (\sqrt{n+1} - \sqrt{n})$

Sol. Now,

$$\sqrt{n+1} - \sqrt{n} = (\sqrt{n+1} - \sqrt{n}) \cdot \frac{\sqrt{n+1} + \sqrt{n}}{\sqrt{n+1} + \sqrt{n}} = \frac{1}{\sqrt{n+1} + \sqrt{n}}.$$

Here $\sum |a_n| = \sum \frac{1}{\sqrt{n+1} + \sqrt{n}}$. Let $b_n = \frac{1}{\sqrt{n}}$.

Now, $\lim_{n \rightarrow \infty} \frac{|a_n|}{b_n} = \lim_{n \rightarrow \infty} \frac{\sqrt{n}}{\sqrt{n+1} + \sqrt{n}} = \lim_{n \rightarrow \infty} \frac{1}{\sqrt{1 + \frac{1}{n}} + 1} = \frac{1}{2} \neq 0$.

Here, $\sum b_n$ is a divergent p -series. Hence, $\sum |a_n|$ is divergent implies $\sum (-1)^n (\sqrt{n+1} - \sqrt{n})$ is not absolutely convergent.



Again, $\{\frac{1}{\sqrt{n+1} + \sqrt{n}}\}$ is a decreasing sequence of positive terms, i.e., $u_n \geq u_{n+1} \forall n$.



Again, $\{\frac{1}{\sqrt{n+1} + \sqrt{n}}\}$ is a decreasing sequence of positive terms, i.e., $u_n \geq u_{n+1} \forall n$.

Also, $\lim u_n \rightarrow 0$.



Again, $\{\frac{1}{\sqrt{n+1} + \sqrt{n}}\}$ is a decreasing sequence of positive terms, i.e., $u_n \geq u_{n+1} \forall n$.

Also, $\lim u_n \rightarrow 0$.

Hence, using Leibniz's test, $\sum (-1)^n (\sqrt{n+1} - \sqrt{n})$ is conditionally convergent.



Q.: $\sum \frac{\cos n\pi}{n \sqrt{n}}.$

Sol. Convergent.

Q.: $\sum (-1)^n \frac{\ln n}{n - \ln n}.$

Sol. Conditionally convergent.



Summary for checking the Convergence of a Series

- If $\lim a_n \nrightarrow 0$, the series diverges.



Summary for checking the Convergence of a Series

- If $\lim a_n \nrightarrow 0$, the series diverges.
- If $\lim a_n = 0$, check whether it is a known series (like geometric series, p-series etc.).



Summary for checking the Convergence of a Series

- If $\lim a_n \nrightarrow 0$, the series diverges.
- If $\lim a_n = 0$, check whether it is a known series (like geometric series, p-series etc.).
- For nonnegative (positive) terms series try integral test, comparison tests, ratio test, root test etc.



Summary for checking the Convergence of a Series

- If $\lim a_n \nrightarrow 0$, the series diverges.
- If $\lim a_n = 0$, check whether it is a known series (like geometric series, p-series etc.).
- For nonnegative (positive) terms series try integral test, comparison tests, ratio test, root test etc.
- For series with some negative terms (not necessarily alternating series) see whether $\sum |a_n|$ converges. If yes, so does $\sum a_n$.



Summary for checking the Convergence of a Series

- If $\lim a_n \nrightarrow 0$, the series diverges.
- If $\lim a_n = 0$, check whether it is a known series (like geometric series, p-series etc.).
- For nonnegative (positive) terms series try integral test, comparison tests, ratio test, root test etc.
- For series with some negative terms (not necessarily alternating series) see whether $\sum |a_n|$ converges. If yes, so does $\sum a_n$.
- For alternating series apply Leibniz's test.



Section 10.7

Power Series



Think of the series where its terms are functions of the variable x .



Think of the series where its terms are functions of the variable x .

$$\frac{1}{3-x} =$$



Think of the series where its terms are functions of the variable x .

$$\frac{1}{3-x} = \sum (x-2)^n$$



Think of the series where its terms are functions of the variable x .

$$\frac{1}{3-x} = \sum (x-2)^n$$

- A **power series about $x = a$** is a series of the form $\sum_{n=0}^{\infty} a_n(x-a)^n$ in which the center a and coefficients a_n are constants.



Think of the series where its terms are functions of the variable x .

$$\frac{1}{3-x} = \sum (x-2)^n$$

- A **power series about $x = a$** is a series of the form $\sum_{n=0}^{\infty} a_n(x-a)^n$ in which the center a and coefficients a_n are constants.
- A **power series about $x = 0$** is $\sum_{n=0}^{\infty} a_n x^n$.



Examples

- $\sum_{n=0}^{\infty} x^n$ (geometric series).



Examples

- $\sum_{n=0}^{\infty} x^n$ (geometric series).
- $\sum_{n=0}^{\infty} \frac{x^n}{n!}$ (exponential series).



Examples

- $\sum_{n=0}^{\infty} x^n$ (geometric series).
- $\sum_{n=0}^{\infty} \frac{x^n}{n!}$ (exponential series).
- $\sum (-1)^{n+1} \frac{x^n}{n}$ (logarithmic series).



Convergence of Power series

For what values of x a power series converge.



Convergence of Power series

For what values of x a power series converge.

$$\sum (x - 2)^n$$



Convergence of Power series

For what values of x a power series converge.

$\sum (x-2)^n$ it is a geometric series which converges whenever



Convergence of Power series

For what values of x a power series converge.

$\sum (x-2)^n$ it is a geometric series which converges whenever $|r| = |x-2| < 1$, i.e., for each x , $1 < x < 3$, the series converges to $\frac{1}{3-x}$.



Ratio Test (in general form)

Let $\sum a_n$ be any series and $\lim \left| \frac{a_{n+1}}{a_n} \right| = r$.



Ratio Test (in general form)

Let $\sum a_n$ be any series and $\lim \left| \frac{a_{n+1}}{a_n} \right| = r$.

(a) If $r < 1$, the series $\sum a_n$ is absolutely convergent;



Ratio Test (in general form)

Let $\sum a_n$ be any series and $\lim \left| \frac{a_{n+1}}{a_n} \right| = r$.

- (a) If $r < 1$, the series $\sum a_n$ is absolutely convergent;
- (b) If $r > 1$, the series $\sum a_n$ is divergent;



Ratio Test (in general form)

Let $\sum a_n$ be any series and $\lim \left| \frac{a_{n+1}}{a_n} \right| = r$.

- (a) If $r < 1$, the series $\sum a_n$ is absolutely convergent;
- (b) If $r > 1$, the series $\sum a_n$ is divergent;
- (c) If $r = 1$, the test is inconclusive.



Root Test or n^{th} Root Test (in general form)

Let $\sum a_n$ be any series and suppose that $\lim(|a_n|)^{\frac{1}{n}} = r$.



Root Test or n^{th} Root Test (in general form)

Let $\sum a_n$ be any series and suppose that $\lim(|a_n|)^{\frac{1}{n}} = r$.

(a) If $r < 1$, the series $\sum a_n$ converges absolutely;



Root Test or n^{th} Root Test (in general form)

Let $\sum a_n$ be any series and suppose that $\lim(|a_n|)^{\frac{1}{n}} = r$.

- (a) If $r < 1$, the series $\sum a_n$ converges absolutely;
- (b) If $r > 1$, the series $\sum a_n$ diverges;



Root Test or n^{th} Root Test (in general form)

Let $\sum a_n$ be any series and suppose that $\lim(|a_n|)^{\frac{1}{n}} = r$.

- (a) If $r < 1$, the series $\sum a_n$ converges absolutely;
- (b) If $r > 1$, the series $\sum a_n$ diverges;
- (c) If $r = 1$, the test is inconclusive.



Testing a Power Series for Convergence

Step-1. Testing for absolute convergence: Use Ratio (or Root) test to find an open interval where the series converges absolutely as follows:



Testing a Power Series for Convergence

Step-1. Testing for absolute convergence: Use Ratio (or Root) test to find an open interval where the series converges absolutely as follows:

$$\begin{aligned}\lim \left| \frac{u_{n+1}}{u_n} \right| &= \lim \left| \frac{a_{n+1}}{a_n} (x - a) \right| \\ &= |x - a| \lim \left| \frac{a_{n+1}}{a_n} \right| \\ &= \frac{|x - a|}{R}.\end{aligned}$$



Testing a Power Series for Convergence

Now for convergence, we must have (ratio test)

$$\lim \left| \frac{u_{n+1}}{u_n} \right| < 1$$

$$\frac{|x - a|}{R} < 1$$

$$|x - a| < R$$

$$x \in (a - R, a + R).$$



Testing a Power Series for Convergence

Step-2. Testing at the end points: If R is finite, then test for absolute (or conditional) convergence at both end points $x = a - R$ and $x = a + R$ (by substituting $x = a - R$ and $x = a + R$ respectively in the given power series).



Testing a Power Series for Convergence

Step-2. Testing at the end points: If R is finite, then test for absolute (or conditional) convergence at both end points $x = a - R$ and $x = a + R$ (by substituting $x = a - R$ and $x = a + R$ respectively in the given power series).

Step-3. Make a conclusion based on Step 1 and Step 2.



Radius of Convergence

The interval $(a - R, a + R)$ is called the **interval of absolute convergence** of the power series



Radius of Convergence

The interval $(a - R, a + R)$ is called the **interval of absolute convergence** of the power series and the **half length** of the interval (that is R) is called the **radius of convergence** of the power series.



Radius of Convergence

The interval $(a - R, a + R)$ is called the **interval of absolute convergence** of the power series and the **half length** of the interval (that is R) is called the **radius of convergence** of the power series.

Remark.



Radius of Convergence

The interval $(a - R, a + R)$ is called the **interval of absolute convergence** of the power series and the **half length** of the interval (that is R) is called the **radius of convergence** of the power series.

Remark.

- If series converges only at center then we say $R = 0$.



Radius of Convergence

The interval $(a - R, a + R)$ is called the **interval of absolute convergence** of the power series and the **half length** of the interval (that is R) is called the **radius of convergence** of the power series.

Remark.

- If series converges only at center then we say $R = 0$.
- If series converges for all x then we say $R = \infty$.



Formula for Radius of Convergence

For the power series $\sum_{n=0}^{\infty} a_n(x-a)^n$ the radius of convergence is given by the formulae



Formula for Radius of Convergence

For the power series $\sum_{n=0}^{\infty} a_n(x-a)^n$ the radius of convergence is given by the formulae

$$\frac{1}{R} = \lim \left| \frac{a_{n+1}}{a_n} \right|.$$



Formula for Radius of Convergence

For the power series $\sum_{n=0}^{\infty} a_n(x-a)^n$ the radius of convergence is given by the formulae

$$\frac{1}{R} = \lim \left| \frac{a_{n+1}}{a_n} \right|.$$

or

$$\frac{1}{R} = \lim |a_n|^{1/n}.$$



Convergence of Power series

For a power series $\sum_{n=0}^{\infty} a_n(x-a)^n$, exactly one of the following is true:

- The series converge only at the center, i.e., $x = a$



Convergence of Power series

For a power series $\sum_{n=0}^{\infty} a_n(x-a)^n$, exactly one of the following is true:

- The series converge only at the center, i.e., $x = a$
- The series converges absolutely for all real numbers x



Convergence of Power series

For a power series $\sum_{n=0}^{\infty} a_n(x-a)^n$, exactly one of the following is true:

- The series converge only at the center, i.e., $x = a$
- The series converges absolutely for all real numbers x
- The series converges absolutely over a finite interval, $|x-a| < R$, i.e., $(a-R, a+R)$ such that $R > 0$.



Convergence of Power series

For a power series $\sum_{n=0}^{\infty} a_n(x-a)^n$, exactly one of the following is true:

- The series converge only at the center, i.e., $x = a$
- The series converges absolutely for all real numbers x
- The series converges absolutely over a finite interval, $|x-a| < R$, i.e., $(a-R, a+R)$ such that $R > 0$. The series may converge absolutely, converge conditionally or diverge, at the end points $x = a-R$ and $x = a+R$.



Q.: $\sum_{n=0}^{\infty} \frac{x^n}{2^n(n+1)^2}.$



Q: $\sum_{n=0}^{\infty} \frac{x^n}{2^n(n+1)^2}.$

Step-1. Testing for absolute convergence
(Ratio Test):



Q.: $\sum_{n=0}^{\infty} \frac{x^n}{2^n(n+1)^2}.$

Step-1. Testing for absolute convergence

(Ratio Test): $\lim \left| \frac{u_{n+1}}{u_n} \right| = \lim \left| \frac{(x)^{n+1}}{2^{n+1}(n+2)^2} \cdot \frac{2^n(n+1)^2}{x^n} \right| =$
 $\lim \left(\frac{n+1}{n+2} \right)^2 \frac{|x|}{2} = \frac{|x|}{2}.$



Q.: $\sum_{n=0}^{\infty} \frac{x^n}{2^n(n+1)^2}.$

Step-1. Testing for absolute convergence

(Ratio Test): $\lim \left| \frac{u_{n+1}}{u_n} \right| = \lim \left| \frac{(x)^{n+1}}{2^{n+1}(n+2)^2} \cdot \frac{2^n(n+1)^2}{x^n} \right| =$
 $\lim \left(\frac{n+1}{n+2} \right)^2 \frac{|x|}{2} = \frac{|x|}{2}.$

For the absolute convergence,



Q.: $\sum_{n=0}^{\infty} \frac{x^n}{2^n(n+1)^2}.$

Step-1. Testing for absolute convergence

(Ratio Test): $\lim \left| \frac{u_{n+1}}{u_n} \right| = \lim \left| \frac{(x)^{n+1}}{2^{n+1}(n+2)^2} \cdot \frac{2^n(n+1)^2}{x^n} \right| =$
 $\lim \left(\frac{n+1}{n+2} \right)^2 \frac{|x|}{2} = \frac{|x|}{2}.$

For the absolute convergence, we must have $\lim \left| \frac{u_{n+1}}{u_n} \right| < 1$
i.e., $|x| < 2$ or $x \in (-2, 2)$. Thus the given series is absolutely convergent for all $x \in (-2, 2)$.



Step-2. Testing at the end points $x = -2$ and $x = 2$:



Step-2. Testing at the end points $x = -2$ and $x = 2$:

- At $x = 2$, the given series becomes $\sum_{n=1}^{\infty} \frac{1}{(n+1)^2}$



Step-2. Testing at the end points $x = -2$ and $x = 2$:

- At $x = 2$, the given series becomes $\sum_{n=1}^{\infty} \frac{1}{(n+1)^2}$ which is convergent p-series.



Step-2. Testing at the end points $x = -2$ and $x = 2$:

- At $x = 2$, the given series becomes $\sum_{n=1}^{\infty} \frac{1}{(n+1)^2}$ which is convergent p-series.
- At $x = -2$, the given series becomes $\sum_{n=1}^{\infty} \frac{(-1)^n}{(n+1)^2}$



Step-2. Testing at the end points $x = -2$ and $x = 2$:

- At $x = 2$, the given series becomes $\sum_{n=1}^{\infty} \frac{1}{(n+1)^2}$ which is convergent p-series.
- At $x = -2$, the given series becomes $\sum_{n=1}^{\infty} \frac{(-1)^n}{(n+1)^2}$ which is absolutely convergent.



Step-3. Conclusion: For the given series
(i) $R = 2$.



Step-3. Conclusion: For the given series

(i) $R = 2$.

(ii) For $-2 \leq x \leq 2 \rightarrow$ absolutely convergent.



Step-3. Conclusion: For the given series

- (i) $R = 2$.
- (ii) For $-2 \leq x \leq 2 \rightarrow$ absolutely convergent.
- (iii) there are no values for which the series converges conditionally.



Q.: $\sum_{n=0}^{\infty} \frac{nx^n}{n+2}$



Q: $\sum_{n=0}^{\infty} \frac{nx^n}{n+2}$

Step-1. Testing for absolute convergence
(Ratio Test):



Q: $\sum_{n=0}^{\infty} \frac{nx^n}{n+2}$

Step-1. Testing for absolute convergence

(Ratio Test): $\lim \left| \frac{u_{n+1}}{u_n} \right| = \lim \left| \frac{(x)^{n+1}(n+1)}{n+3} \cdot \frac{n+2}{nx^n} \right| =$
 $|x| \lim \frac{(n+1)(n+2)}{n(n+3)} = |x|.$



Q.: $\sum_{n=0}^{\infty} \frac{nx^n}{n+2}$

Step-1. Testing for absolute convergence

(Ratio Test): $\lim \left| \frac{u_{n+1}}{u_n} \right| = \lim \left| \frac{(x)^{n+1}(n+1)}{n+3} \cdot \frac{n+2}{nx^n} \right| =$
 $|x| \lim \frac{(n+1)(n+2)}{n(n+3)} = |x|.$

For the absolute convergence, we must have $\lim \left| \frac{u_{n+1}}{u_n} \right| < 1$
i.e., $|x| < 1$ or $x \in (-1, 1)$. Thus the given series is absolutely convergent for all $x \in (-1, 1)$.



Step-2. Testing at the end points $x = -1$ and $x = 1$:



Step-2. Testing at the end points $x = -1$ and $x = 1$:

- At $x = 1$, the given series becomes $\sum_{n=0}^{\infty} \frac{n}{n+2}$



Step-2. Testing at the end points $x = -1$ and $x = 1$:

- At $x = 1$, the given series becomes $\sum_{n=0}^{\infty} \frac{n}{n+2}$ which is divergent



Step-2. Testing at the end points $x = -1$ and $x = 1$:

- At $x = 1$, the given series becomes $\sum_{n=0}^{\infty} \frac{n}{n+2}$ which is divergent (n^{th} term test).



Step-2. Testing at the end points $x = -1$ and $x = 1$:

- At $x = 1$, the given series becomes $\sum_{n=0}^{\infty} \frac{n}{n+2}$ which is divergent (n^{th} term test).
- At $x = -1$, the given series becomes $\sum_{n=0}^{\infty} (-1)^n \frac{n}{n+2}$ which is again divergent.



Step-3. Conclusion: For the given series
(i) $R = 1$.



Step-3. Conclusion: For the given series

- (i) $R = 1$.
- (ii) For $-1 < x < 1$, it is absolutely convergent.



Step-3. Conclusion: For the given series

- (i) $R = 1$.
- (ii) For $-1 < x < 1$, it is absolutely convergent.
- (iii) there are no values for which the series converges conditionally



Q.: $\sum_{n=1}^{\infty} \frac{(3x+1)^{n+1}}{2n+2}$



Q: $\sum_{n=1}^{\infty} \frac{(3x+1)^{n+1}}{2n+2}$

Step-1. Testing for absolute convergence
(Ratio Test):



Q: $\sum_{n=1}^{\infty} \frac{(3x+1)^{n+1}}{2n+2}$

Step-1. Testing for absolute convergence

(Ratio Test): $\lim \left| \frac{u_{n+1}}{u_n} \right| = \lim \left| \frac{(3x+1)^{n+2}}{2n+4} \cdot \frac{2n+2}{(3x+1)^{n+1}} \right| =$
 $|3x+1| \lim \frac{(2n+2)}{(2n+4)} = |3x+1|.$



Q.: $\sum_{n=1}^{\infty} \frac{(3x+1)^{n+1}}{2n+2}$

Step-1. Testing for absolute convergence

(Ratio Test): $\lim \left| \frac{u_{n+1}}{u_n} \right| = \lim \left| \frac{(3x+1)^{n+2}}{2n+4} \cdot \frac{2n+2}{(3x+1)^{n+1}} \right| =$
 $|3x+1| \lim \frac{(2n+2)}{(2n+4)} = |3x+1|.$

For the absolute convergence, we must have $\lim \left| \frac{u_{n+1}}{u_n} \right| < 1$
i.e., $|3x+1| < 1$ or $x \in (-\frac{2}{3}, 0).$



Q: $\sum_{n=1}^{\infty} \frac{(3x+1)^{n+1}}{2n+2}$

Step-1. Testing for absolute convergence

(Ratio Test): $\lim \left| \frac{u_{n+1}}{u_n} \right| = \lim \left| \frac{(3x+1)^{n+2}}{2n+4} \cdot \frac{2n+2}{(3x+1)^{n+1}} \right| =$
 $|3x+1| \lim \frac{(2n+2)}{(2n+4)} = |3x+1|.$

For the absolute convergence, we must have $\lim \left| \frac{u_{n+1}}{u_n} \right| < 1$
i.e., $|3x+1| < 1$ or $x \in (-\frac{2}{3}, 0)$. Thus the given series is absolutely convergent for all $x \in (-\frac{2}{3}, 0)$.



Step-2. Testing at the end points $x = -\frac{2}{3}$ and $x = 0$:



Step-2. Testing at the end points $x = -\frac{2}{3}$ and $x = 0$:

- At $x = -\frac{2}{3}$, the given series becomes
$$\sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{2n+2}$$



Step-2. Testing at the end points $x = -\frac{2}{3}$ and $x = 0$:

- At $x = -\frac{2}{3}$, the given series becomes $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{2n+2}$ which is a



Step-2. Testing at the end points $x = -\frac{2}{3}$ and $x = 0$:

- At $x = -\frac{2}{3}$, the given series becomes $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{2n+2}$ which is a conditionally convergent series.



Step-2. Testing at the end points $x = -\frac{2}{3}$ and $x = 0$:

- At $x = -\frac{2}{3}$, the given series becomes $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{2n+2}$ which is a conditionally convergent series.
- At $x = 0$, the given series becomes $\sum_{n=1}^{\infty} \frac{1}{2n+2}$ which is a divergent series



Step-2. Testing at the end points $x = -\frac{2}{3}$ and $x = 0$:

- At $x = -\frac{2}{3}$, the given series becomes $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{2n+2}$ which is a conditionally convergent series.
- At $x = 0$, the given series becomes $\sum_{n=1}^{\infty} \frac{1}{2n+2}$ which is a divergent series (Why)



Step-2. Testing at the end points $x = -\frac{2}{3}$ and $x = 0$:

- At $x = -\frac{2}{3}$, the given series becomes $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{2n+2}$ which is a conditionally convergent series.
- At $x = 0$, the given series becomes $\sum_{n=1}^{\infty} \frac{1}{2n+2}$ which is a divergent series (Why) (from DCT).



Step-3. Conclusion: For the given series

(i) $R = \frac{1}{3}$



Step-3. Conclusion: For the given series

(i) $R = \frac{1}{3}$ ($|x - a| < R$).



Step-3. Conclusion: For the given series

(i) $R = \frac{1}{3}$ ($|x - a| < R$).

(ii) For $-\frac{2}{3} < x < 0$, it is absolutely convergent.



Step-3. Conclusion: For the given series

- (i) $R = \frac{1}{3}$ ($|x - a| < R$).
- (ii) For $-\frac{2}{3} < x < 0$, it is absolutely convergent.
- (iii) It converges conditionally for $x = -\frac{2}{3}$



Q: $\sum_{n=1}^{\infty} n!(x+10)^n.$



Q:. $\sum_{n=1}^{\infty} n!(x+10)^n.$

Sol. Here

$$\lim \left| \frac{u_{n+1}}{u_n} \right| = \lim \left| \frac{(n+1)!(x+10)^{n+1}}{n!(x+10)^n} \right| = \lim (n+1)|x+10|.$$



Q:. $\sum_{n=1}^{\infty} n!(x+10)^n.$

Sol. Here

$$\lim \left| \frac{u_{n+1}}{u_n} \right| = \lim \left| \frac{(n+1)!(x+10)^{n+1}}{n!(x+10)^n} \right| = \lim (n+1)|x+10|.$$

Now $\lim (n+1)|x+10| < 1$ only if



Q.: $\sum_{n=1}^{\infty} n!(x+10)^n.$

Sol. Here

$$\lim \left| \frac{u_{n+1}}{u_n} \right| = \lim \left| \frac{(n+1)!(x+10)^{n+1}}{n!(x+10)^n} \right| = \lim (n+1)|x+10|.$$

Now $\lim (n+1)|x+10| < 1$ only if $|x+10| = 0$, i.e., when $x = -10$.



Q.: $\sum_{n=1}^{\infty} n!(x+10)^n.$

Sol. Here

$$\lim \left| \frac{u_{n+1}}{u_n} \right| = \lim \left| \frac{(n+1)!(x+10)^{n+1}}{n!(x+10)^n} \right| = \lim (n+1)|x+10|.$$

Now $\lim (n+1)|x+10| < 1$ only if $|x+10| = 0$, i.e., when $x = -10$. Thus, $R = 0$ and the series converges only at $x = -10$.



Q: Find the radius of convergence $\sum_{n=1}^{\infty} \left(\frac{n}{n+1}\right)^{n^2} x^n$



Q:. Find the radius of convergence $\sum_{n=1}^{\infty} \left(\frac{n}{n+1}\right)^{n^2} x^n$

Sol. $\lim |u_n|^{\frac{1}{n}} = \lim \left(\left(\frac{n}{n+1}\right)^{n^2} |x|^n \right)^{\frac{1}{n}} = |x| \lim \left(\frac{n}{n+1}\right)^n$



Q:. Find the radius of convergence $\sum_{n=1}^{\infty} \left(\frac{n}{n+1}\right)^{n^2} x^n$

Sol. $\lim |u_n|^{\frac{1}{n}} = \lim \left(\left(\frac{n}{n+1}\right)^{n^2} |x|^n \right)^{\frac{1}{n}} = |x| \lim \left(\frac{n}{n+1}\right)^n = |x| \frac{1}{e}.$



Q:. Find the radius of convergence $\sum_{n=1}^{\infty} \left(\frac{n}{n+1}\right)^{n^2} x^n$

Sol. $\lim |u_n|^{\frac{1}{n}} = \lim \left(\left(\frac{n}{n+1}\right)^{n^2} |x|^n \right)^{\frac{1}{n}} = |x| \lim \left(\frac{n}{n+1}\right)^n = |x| \frac{1}{e}.$

For the absolute convergence, we must have
 $\lim |u_n|^{\frac{1}{n}} < 1 \implies |x| < e \implies R = e.$



Questions

Discuss the nature of convergence of the following power series:

1. $\sum_{n=0}^{\infty} \frac{(x-3)^{2n}}{3^n}$

2. $\sum \frac{x^n}{n!}$

Ans 1. (i) $R = \sqrt{3}$.

(ii) For $3 - \sqrt{3} < x < 3 + \sqrt{3} \rightarrow$ absolutely convergent.

(iii) For $3 - \sqrt{3} < x < 3 - \sqrt{3} \rightarrow$ convergent.

(iv) No point of conditional convergence.

Ans 2. Absolutely convergent for all values of x .



Section 10.8

Taylor and Maclaurin Series



Q:. Can we expand an infinitely differentiable function (such as $f(x) = \sin x$ or $f(x) = e^x$) into a power series $\sum a_n(x-a)^n$ that converge to the correct function value $f(x)$ for all x in some open interval $(a-R, a+R)$, where $R > 0$ or $R = \infty$.



Q:. Can we expand an infinitely differentiable function (such as $f(x) = \sin x$ or $f(x) = e^x$) into a power series $\sum a_n(x-a)^n$ that converge to the correct function value $f(x)$ for all x in some open interval $(a-R, a+R)$, where $R > 0$ or $R = \infty$. To proceed further let us assume that an infinitely differentiable function f on an interval $(a-R, a+R)$ can be represented by a power series $\sum a_n(x-a)^n$ on that interval. Our aim is to determine the coefficients a_n .



Taylor Series

Let $f(x)$ be a function with derivatives of all orders in an interval containing a as an interior point.



Taylor Series

Let $f(x)$ be a function with derivatives of all orders in an interval containing a as an interior point. Then the **Taylor series** generated by $f(x)$ at a is given by



Taylor Series

Let $f(x)$ be a function with derivatives of all orders in an interval containing a as an interior point. Then the **Taylor series** generated by $f(x)$ at a is given by

$$\sum_{n=0}^{\infty} a_n(x-a)^n, \text{ where } a_n = \frac{f^{(n)}(a)}{n!}.$$



Taylor Series

Let $f(x)$ be a function with derivatives of all orders in an interval containing a as an interior point. Then the **Taylor series** generated by $f(x)$ at a is given by

$$\sum_{n=0}^{\infty} a_n(x-a)^n, \text{ where } a_n = \frac{f^{(n)}(a)}{n!}.$$

That is the Taylor series is

$$f(a) + \frac{f'(a)}{1!}(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \cdots + \frac{f^{(n)}(a)}{n!}(x-a)^n + \cdots$$



Maclaurin Series

The **Maclaurin series** generated by $f(x)$ is given by

$$\sum_{n=0}^{\infty} a_n x^n, \text{ where } a_n = \frac{f^{(n)}(0)}{n!}.$$

That is the Maclaurin series is

$$f(0) + \frac{f'(0)}{1!}x + \frac{f''(0)}{2!}x^2 + \dots + \frac{f^{(n)}(0)}{n!}x^n + \dots$$



Maclaurin Series

The **Maclaurin series** generated by $f(x)$ is given by

$$\sum_{n=0}^{\infty} a_n x^n, \text{ where } a_n = \frac{f^{(n)}(0)}{n!}.$$

That is the Maclaurin series is

$$f(0) + \frac{f'(0)}{1!}x + \frac{f''(0)}{2!}x^2 + \dots + \frac{f^{(n)}(0)}{n!}x^n + \dots$$

The Maclaurin series is the Taylor series generated by f at $x = 0$.



Q:. Determine the Maclaurin series of $\cos x$.



Q.: Determine the Maclaurin series of $\cos x$.

Sol. Here

$$f(x) = \cos x \quad \Rightarrow f(0) = 1$$

$$f'(x) = -\sin x \quad \Rightarrow f'(0) = 0$$

$$f''(x) = -\cos x \quad \Rightarrow f''(0) = -1$$

$$f'''(x) = \sin x \quad \Rightarrow f'''(0) = 0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots$$



Maclaurin series is given as

$$f(0) + \frac{f'(0)}{1!}x + \frac{f''(0)}{2!}x^2 + \dots + \frac{f^{(n)}(0)}{n!}x^n + \dots$$



Maclaurin series is given as

$$f(0) + \frac{f'(0)}{1!}x + \frac{f''(0)}{2!}x^2 + \dots + \frac{f^{(n)}(0)}{n!}x^n + \dots$$

Thus, the required series is

$$1 - \frac{1}{2!}x^2 + \frac{1}{4!}x^4 + \dots = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)!}x^{2n}$$



Q:30. Determine the Taylor series of 2^x at $x = 1$.



Q:30. Determine the Taylor series of 2^x at $x = 1$.

Sol. Here

$$f(x) = 2^x \Rightarrow f(1) = 2$$

$$f'(x) = 2^x \ln 2 \Rightarrow f'(1) = 2 \ln 2$$

$$f''(x) = 2^x (\ln 2)^2 \Rightarrow f''(1) = 2 (\ln 2)^2$$

$$f'''(x) = 2^x (\ln 2)^3 \Rightarrow f'''(1) = 2 (\ln 2)^3$$

$$f^{iv}(x) = 2^x (\ln 2)^4 \Rightarrow f^{iv}(1) = 2 (\ln 2)^4$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots$$

$$f^{(n)}(x) = 2^x (\ln 2)^n \Rightarrow f^{(n)}(1) = 2 (\ln 2)^n$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots$$



The Taylor series is

$$f(a) + \frac{f'(a)}{1!}(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \cdots + \frac{f^{(n)}(a)}{n!}(x-a)^n + \cdots$$



The Taylor series is

$$f(a) + \frac{f'(a)}{1!}(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \cdots + \frac{f^{(n)}(a)}{n!}(x-a)^n + \cdots$$

Thus, the required series is

$$2 + 2\ln 2(x-1) + \frac{2(\ln 2)^2}{2!}(x-1)^2 + \cdots + \frac{2(\ln 2)^n}{n!}(x-1)^n + \cdots$$



Taylor Polynomial

Let $f(x)$ be a function with derivatives of order k for $k = 1, 2, \dots, N$ in an interval containing a as an interior point.



Taylor Polynomial

Let $f(x)$ be a function with derivatives of order k for $k = 1, 2, \dots, N$ in an interval containing a as an interior point. Then for any integer n from 0 to N , the **Taylor polynomial of order n** generated by $f(x)$ at $x = a$ is given by



Taylor Polynomial

Let $f(x)$ be a function with derivatives of order k for $k = 1, 2, \dots, N$ in an interval containing a as an interior point. Then for any integer n from 0 to N , the **Taylor polynomial of order n** generated by $f(x)$ at $x = a$ is given by

$$P_n(x) = f(a) + \frac{f'(a)}{1!}(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \dots + \frac{f^{(n)}(a)}{n!}(x-a)^n$$



Taylor Polynomial

Let $f(x)$ be a function with derivatives of order k for $k = 1, 2, \dots, N$ in an interval containing a as an interior point. Then for any integer n from 0 to N , the **Taylor polynomial of order n** generated by $f(x)$ at $x = a$ is given by

$$P_n(x) = f(a) + \frac{f'(a)}{1!}(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \dots + \frac{f^{(n)}(a)}{n!}(x-a)^n$$



Q:. Find the Taylor polynomial of order 3 generated by $f(x) = \frac{1}{x}$ at $a = 2$.



Q:. Find the Taylor polynomial of order 3 generated by $f(x) = \frac{1}{x}$ at $a = 2$.

Sol.

$$f(x) = x^{-1} \quad \Rightarrow f(2) = 1/2$$

$$f'(x) = -x^{-2} \quad \Rightarrow f'(2) = -1/4$$

$$f''(x) = 2x^{-3} \quad \Rightarrow f''(2) = 1/4$$

$$f'''(x) = -6x^{-4} \quad \Rightarrow f'''(2) = -3/8$$



Q:. Find the Taylor polynomial of order 3 generated by $f(x) = \frac{1}{x}$ at $a = 2$.

Sol.

$$f(x) = x^{-1} \quad \Rightarrow f(2) = 1/2$$

$$f'(x) = -x^{-2} \quad \Rightarrow f'(2) = -1/4$$

$$f''(x) = 2x^{-3} \quad \Rightarrow f''(2) = 1/4$$

$$f'''(x) = -6x^{-4} \quad \Rightarrow f'''(2) = -3/8$$

$$P_3(x) = \frac{1}{2} - \frac{1}{4}(x-2) + \frac{1}{8}(x-2)^2 - \frac{1}{16}(x-2)^3$$

