

MATHEMATICS-II (MATH F112)

Dr. Krishnendra Shekhawat

BITS PILANI
Department of Mathematics



Section 4.6

Constructing Special Bases



Basis



Basis

A subset B of a vector space V is said to be a **basis** of V if



Basis

A subset B of a vector space V is said to be a **basis** of V if

- $\text{span}(B) = V$,



Basis

A subset B of a vector space V is said to be a **basis** of V if

- $\text{span}(B) = V$, i.e., B spans V and



Basis

A subset B of a vector space V is said to be a **basis** of V if

- $\text{span}(B) = V$, i.e., B spans V and
- B is LI.



Basis

A subset B of a vector space V is said to be a **basis** of V if

- $\text{span}(B) = V$, i.e., B spans V and
- B is LI.

The **dimension** of a vector space V



Basis

A subset B of a vector space V is said to be a **basis** of V if

- $\text{span}(B) = V$, i.e., B spans V and
- B is LI.

The **dimension** of a vector space V is the number of vectors in a basis of V



Basis

A subset B of a vector space V is said to be a **basis** of V if

- $\text{span}(B) = V$, i.e., B spans V and
- B is LI.

The **dimension** of a vector space V is the number of vectors in a basis of V and it is denoted by $\dim(V)$.



Examples



Examples

The subset $B = \{[1, 0], [0, 1]\}$ is a basis of \mathbb{R}^2



Examples

The subset $B = \{[1, 0], [0, 1]\}$ is a basis of \mathbb{R}^2 as $\text{span}(B) = \mathbb{R}^2$ and



Examples

The subset $B = \{[1, 0], [0, 1]\}$ is a basis of \mathbb{R}^2 as $\text{span}(B) = \mathbb{R}^2$ and B is LI .



Examples

The subset $B = \{[1, 0], [0, 1]\}$ is a basis of \mathbb{R}^2 as $\text{span}(B) = \mathbb{R}^2$ and B is LI. The subset B is called the **standard basis** of \mathbb{R}^2 .



Examples

The subset $B = \{[1, 0], [0, 1]\}$ is a basis of \mathbb{R}^2 as $\text{span}(B) = \mathbb{R}^2$ and B is LI. The subset B is called the **standard basis** of \mathbb{R}^2 . Here, $\dim(\mathbb{R}^2) = 2$.



Examples

The subset $B = \{[1, 0], [0, 1]\}$ is a basis of \mathbb{R}^2 as $\text{span}(B) = \mathbb{R}^2$ and B is LI. The subset B is called the **standard basis** of \mathbb{R}^2 . Here, $\dim(\mathbb{R}^2) = 2$.

The subset $B = \{1, x, x^2, \dots, x^n\}$ is a basis of P_n



Examples

The subset $B = \{[1, 0], [0, 1]\}$ is a basis of \mathbb{R}^2 as $\text{span}(B) = \mathbb{R}^2$ and B is LI. The subset B is called the **standard basis** of \mathbb{R}^2 . Here, $\dim(\mathbb{R}^2) = 2$.

The subset $B = \{1, x, x^2, \dots, x^n\}$ is a basis of P_n as B is LI (why)



Examples

The subset $B = \{[1, 0], [0, 1]\}$ is a basis of \mathbb{R}^2 as $\text{span}(B) = \mathbb{R}^2$ and B is LI. The subset B is called the **standard basis** of \mathbb{R}^2 . Here, $\dim(\mathbb{R}^2) = 2$.

The subset $B = \{1, x, x^2, \dots, x^n\}$ is a basis of P_n as B is LI (why) and $\text{span}(B) = P_n$ (why).



Examples

The subset $B = \{[1, 0], [0, 1]\}$ is a basis of \mathbb{R}^2 as $\text{span}(B) = \mathbb{R}^2$ and B is LI. The subset B is called the **standard basis** of \mathbb{R}^2 . Here, $\dim(\mathbb{R}^2) = 2$.

The subset $B = \{1, x, x^2, \dots, x^n\}$ is a basis of P_n as B is LI (why) and $\text{span}(B) = P_n$ (why). Here, $\dim(P_n) = n + 1$.



Q:. Find a basis and dimension for
 $W = \{p \in P_5 | p(2) = p(3) = 0\}.$



Q:. Find a basis and dimension for
 $W = \{p \in P_5 | p(2) = p(3) = 0\}.$

Sol. Here W can be expressed as



Q:. Find a basis and dimension for
 $W = \{p \in P_5 | p(2) = p(3) = 0\}.$

Sol. Here W can be expressed as
 $W = \{(x-2)(x-3)(ax^3 + bx^2 + cx + d) | a, b, c, d \in \mathbb{R}\} \implies$



Q:. Find a basis and dimension for
 $W = \{p \in P_5 | p(2) = p(3) = 0\}.$

Sol. Here W can be expressed as

$$W = \{(x-2)(x-3)(ax^3 + bx^2 + cx + d) | a, b, c, d \in \mathbb{R}\} \implies$$

$$B = \{(x-2)(x-3), x(x-2)(x-3), x^2(x-2)(x-3), x^3(x-2)(x-3)\}$$



Q:. Find a basis and dimension for
 $W = \{p \in P_5 | p(2) = p(3) = 0\}.$

Sol. Here W can be expressed as

$$W = \{(x-2)(x-3)(ax^3 + bx^2 + cx + d) | a, b, c, d \in \mathbb{R}\} \implies$$

$$B = \{(x-2)(x-3), x(x-2)(x-3), x^2(x-2)(x-3), x^3(x-2)(x-3)\} \text{ spans } W.$$



Q:. Find a basis and dimension for
 $W = \{p \in P_5 | p(2) = p(3) = 0\}.$

Sol. Here W can be expressed as

$$W = \{(x-2)(x-3)(ax^3 + bx^2 + cx + d) | a, b, c, d \in \mathbb{R}\} \implies$$

$B = \{(x-2)(x-3), x(x-2)(x-3), x^2(x-2)(x-3), x^3(x-2)(x-3)\}$ spans W . Also B is LI



Q:. Find a basis and dimension for
 $W = \{p \in P_5 | p(2) = p(3) = 0\}.$

Sol. Here W can be expressed as

$$W = \{(x-2)(x-3)(ax^3 + bx^2 + cx + d) | a, b, c, d \in \mathbb{R}\} \implies$$

$B = \{(x-2)(x-3), x(x-2)(x-3), x^2(x-2)(x-3), x^3(x-2)(x-3)\}$ spans W . Also B is LI (why).



Q:. Find a basis and dimension for
 $W = \{p \in P_5 | p(2) = p(3) = 0\}.$

Sol. Here W can be expressed as

$$W = \{(x-2)(x-3)(ax^3 + bx^2 + cx + d) | a, b, c, d \in \mathbb{R}\} \implies$$

$B = \{(x-2)(x-3), x(x-2)(x-3), x^2(x-2)(x-3), x^3(x-2)(x-3)\}$ spans W . Also B is LI (why).

Hence, B is a basis for W



Q:. Find a basis and dimension for
 $W = \{p \in P_5 | p(2) = p(3) = 0\}.$

Sol. Here W can be expressed as

$$W = \{(x-2)(x-3)(ax^3 + bx^2 + cx + d) | a, b, c, d \in \mathbb{R}\} \implies$$

$B = \{(x-2)(x-3), x(x-2)(x-3), x^2(x-2)(x-3), x^3(x-2)(x-3)\}$ spans W . Also B is LI (why).

Hence, B is a basis for W and $\dim(B) = 4$.



Q:.. Let $W = \{X \in \mathbb{R}^5 : AX = 0\}$,



Q:. Let $W = \{X \in \mathbb{R}^5 : AX = 0\}$, where

$$A = \begin{bmatrix} 1 & 2 & 1 & 0 & -1 \\ 2 & -1 & 0 & 1 & 3 \\ 1 & -3 & -1 & 1 & 4 \\ 2 & 9 & 4 & -1 & -7 \end{bmatrix}$$

Find a basis for W .



Q.: Let $W = \{X \in \mathbb{R}^5 : AX = 0\}$, where

$$A = \begin{bmatrix} 1 & 2 & 1 & 0 & -1 \\ 2 & -1 & 0 & 1 & 3 \\ 1 & -3 & -1 & 1 & 4 \\ 2 & 9 & 4 & -1 & -7 \end{bmatrix}$$

Find a basis for W .

Sol.

$$\{[-1/5, -2/5, 1, 0, 0]^T, [-2/5, 1/5, 0, 1, 0]^T, [-1, 1, 0, 0, 1]^T\}$$



Finding a Basis for a Spanning Set



Finding a Basis for a Spanning Set

Q.: Let $S = \{[1, 2], [3, 0], [0, 2]\} \subseteq \mathbb{R}^2$.



Finding a Basis for a Spanning Set

Q.: Let $S = \{[1, 2], [3, 0], [0, 2]\} \subseteq \mathbb{R}^2$. Find a basis for $\text{span}(S)$.



Finding a Basis for a Spanning Set

Q.: Let $S = \{[1, 2], [3, 0], [0, 2]\} \subseteq \mathbb{R}^2$. Find a basis for $\text{span}(S)$.

Sol. $\text{span}(S) = \{x[1, 2] + y[3, 0] + z[0, 2] | x, y, z \in \mathbb{R}\} \implies$



Finding a Basis for a Spanning Set

Q.: Let $S = \{[1, 2], [3, 0], [0, 2]\} \subseteq \mathbb{R}^2$. Find a basis for $\text{span}(S)$.

Sol. $\text{span}(S) = \{x[1, 2] + y[3, 0] + z[0, 2] | x, y, z \in \mathbb{R}\} \implies$
 $\text{span}(S) = \{[x + 3y, 2x + 2z] | x, y, z \in \mathbb{R}\} \implies$



Finding a Basis for a Spanning Set

Q:. Let $S = \{[1, 2], [3, 0], [0, 2]\} \subseteq \mathbb{R}^2$. Find a basis for $\text{span}(S)$.

Sol. $\text{span}(S) = \{x[1, 2] + y[3, 0] + z[0, 2] | x, y, z \in \mathbb{R}\} \implies$
 $\text{span}(S) = \{[x + 3y, 2x + 2z] | x, y, z \in \mathbb{R}\} \implies$
 $\text{span}(S) = \{(x + 3y)[1, 0] + (x + z)[0, 2] | x, y, z \in \mathbb{R}\} \implies$



Finding a Basis for a Spanning Set

Q:. Let $S = \{[1, 2], [3, 0], [0, 2]\} \subseteq \mathbb{R}^2$. Find a basis for $\text{span}(S)$.

Sol. $\text{span}(S) = \{x[1, 2] + y[3, 0] + z[0, 2] | x, y, z \in \mathbb{R}\} \implies$
 $\text{span}(S) = \{[x + 3y, 2x + 2z] | x, y, z \in \mathbb{R}\} \implies$
 $\text{span}(S) = \{(x + 3y)[1, 0] + (x + z)[0, 2] | x, y, z \in \mathbb{R}\} \implies$
 $\text{span}(S) = \text{span}([1, 0], [0, 2]) = \text{span}(B).$



Finding a Basis for a Spanning Set

Q:. Let $S = \{[1, 2], [3, 0], [0, 2]\} \subseteq \mathbb{R}^2$. Find a basis for $\text{span}(S)$.

Sol. $\text{span}(S) = \{x[1, 2] + y[3, 0] + z[0, 2] | x, y, z \in \mathbb{R}\} \implies$
 $\text{span}(S) = \{[x + 3y, 2x + 2z] | x, y, z \in \mathbb{R}\} \implies$
 $\text{span}(S) = \{(x + 3y)[1, 0] + (x + z)[0, 2] | x, y, z \in \mathbb{R}\} \implies$
 $\text{span}(S) = \text{span}([1, 0], [0, 2]) = \text{span}(B).$

Since, B is LI,



Finding a Basis for a Spanning Set

Q:. Let $S = \{[1, 2], [3, 0], [0, 2]\} \subseteq \mathbb{R}^2$. Find a basis for $\text{span}(S)$.

Sol. $\text{span}(S) = \{x[1, 2] + y[3, 0] + z[0, 2] | x, y, z \in \mathbb{R}\} \implies$
 $\text{span}(S) = \{[x + 3y, 2x + 2z] | x, y, z \in \mathbb{R}\} \implies$
 $\text{span}(S) = \{(x + 3y)[1, 0] + (x + z)[0, 2] | x, y, z \in \mathbb{R}\} \implies$
 $\text{span}(S) = \text{span}([1, 0], [0, 2]) = \text{span}(B).$

Since, B is LI, B forms a basis for $\text{span}(S)$.



Conclusion



Conclusion

In order to find a basis for $\text{span}(S)$, we need a subset B of $\text{span}(S)$ such that



Conclusion

In order to find a basis for $\text{span}(S)$, we need a subset B of $\text{span}(S)$ such that

- $\text{span}(S) = \text{span}(B)$;



Conclusion

In order to find a basis for $\text{span}(S)$, we need a subset B of $\text{span}(S)$ such that

- $\text{span}(S) = \text{span}(B)$;
- B is LI.



Conclusion

In order to find a basis for $\text{span}(S)$, we need a subset B of $\text{span}(S)$ such that

- $\text{span}(S) = \text{span}(B)$;
- B is LI.

Since $[1, 2] = \frac{1}{3}[3, 0] + [0, 2]$



Conclusion

In order to find a basis for $\text{span}(S)$, we need a subset B of $\text{span}(S)$ such that

- $\text{span}(S) = \text{span}(B)$;
- B is LI.

Since $[1, 2] = \frac{1}{3}[3, 0] + [0, 2]$ implies $\text{span}(S) = \text{span}(B)$



Conclusion

In order to find a basis for $\text{span}(S)$, we need a subset B of $\text{span}(S)$ such that

- $\text{span}(S) = \text{span}(B)$;
- B is LI.

Since $[1, 2] = \frac{1}{3}[3, 0] + [0, 2]$ implies $\text{span}(S) = \text{span}(B)$ (why)



Conclusion

In order to find a basis for $\text{span}(S)$, we need a subset B of $\text{span}(S)$ such that

- $\text{span}(S) = \text{span}(B)$;
- B is LI.

Since $[1, 2] = \frac{1}{3}[3, 0] + [0, 2]$ implies $\text{span}(S) = \text{span}(B)$ (why) where $B = \{[3, 0], [0, 2]\}$.



Conclusion

In order to find a basis for $\text{span}(S)$, we need a subset B of $\text{span}(S)$ such that

- $\text{span}(S) = \text{span}(B)$;
- B is LI.

Since $[1, 2] = \frac{1}{3}[3, 0] + [0, 2]$ implies $\text{span}(S) = \text{span}(B)$ (why) where $B = \{[3, 0], [0, 2]\}$. Also, B is LI



Conclusion

In order to find a basis for $\text{span}(S)$, we need a subset B of $\text{span}(S)$ such that

- $\text{span}(S) = \text{span}(B)$;
- B is LI.

Since $[1, 2] = \frac{1}{3}[3, 0] + [0, 2]$ implies $\text{span}(S) = \text{span}(B)$ (why) where $B = \{[3, 0], [0, 2]\}$. Also, B is LI (why).



Conclusion

In order to find a basis for $\text{span}(S)$, we need a subset B of $\text{span}(S)$ such that

- $\text{span}(S) = \text{span}(B)$;
- B is LI.

Since $[1, 2] = \frac{1}{3}[3, 0] + [0, 2]$ implies $\text{span}(S) = \text{span}(B)$ (why) where $B = \{[3, 0], [0, 2]\}$. Also, B is LI (why). Hence, B is a basis of $\text{span}(S)$.



Conclusion

In order to find a basis for $\text{span}(S)$, we need a subset B of $\text{span}(S)$ such that

- $\text{span}(S) = \text{span}(B)$;
- B is LI.

Since $[1, 2] = \frac{1}{3}[3, 0] + [0, 2]$ implies $\text{span}(S) = \text{span}(B)$ (why) where $B = \{[3, 0], [0, 2]\}$. Also, B is LI (why). Hence, B is a basis of $\text{span}(S)$.

Q:. Does there always exists a basis for $\text{span}(S)$.



Recall



Recall

Theorem: Let V be a finite dimensional vector space such that $\dim(V) = n$.



Recall

Theorem: Let V be a finite dimensional vector space such that $\dim(V) = n$. Suppose S is a finite subset of V that spans V . Then $|S| \geq n$.



Recall

Theorem: Let V be a finite dimensional vector space such that $\dim(V) = n$. Suppose S is a finite subset of V that spans V . Then $|S| \geq n$. Moreover, $|S| = n$ if and only if S is a basis of V .



Simplified Span Method to find a Basis for $\text{span}(S)$



Simplified Span Method to find a Basis for $\text{span}(S)$

Let $S \subseteq \mathbb{R}^n$ containing k vectors.



Simplified Span Method to find a Basis for $\text{span}(S)$

Let $S \subseteq \mathbb{R}^n$ containing k vectors.

- Construct a matrix A of order $k \times n$ by using vectors of S



Simplified Span Method to find a Basis for $\text{span}(S)$

Let $S \subseteq \mathbb{R}^n$ containing k vectors.

- Construct a matrix A of order $k \times n$ by using vectors of S as rows of A .



Simplified Span Method to find a Basis for $\text{span}(S)$

Let $S \subseteq \mathbb{R}^n$ containing k vectors.

- Construct a matrix A of order $k \times n$ by using vectors of S as rows of A .
- Compute $C = \text{RREF}(A)$.



Simplified Span Method to find a Basis for $\text{span}(S)$

Let $S \subseteq \mathbb{R}^n$ containing k vectors.

- Construct a matrix A of order $k \times n$ by using vectors of S as rows of A .
- Compute $C = \text{RREF}(A)$.
- Non-zero rows of C forms a basis for $\text{span}(S)$.



Example

Q:. Let $S = \{[1, 2, 3, -1, 0], [3, 6, 8, -2, 0],$
 $[-1, -1, -3, 1, 1], [-2, -3, -5, 1, 1]\}$ be a subset of \mathbb{R}^5 .



Example

Q:. Let $S = \{[1, 2, 3, -1, 0], [3, 6, 8, -2, 0],$
 $[-1, -1, -3, 1, 1], [-2, -3, -5, 1, 1]\}$ be a subset of \mathbb{R}^5 .
Find a basis for $\text{span}(S)$.



Example

Q:. Let $S = \{[1, 2, 3, -1, 0], [3, 6, 8, -2, 0],$
 $[-1, -1, -3, 1, 1], [-2, -3, -5, 1, 1]\}$ be a subset of \mathbb{R}^5 .
Find a basis for $\text{span}(S)$.

Sol. Step 1.



Example

Q.: Let $S = \{[1, 2, 3, -1, 0], [3, 6, 8, -2, 0], [-1, -1, -3, 1, 1], [-2, -3, -5, 1, 1]\}$ be a subset of \mathbb{R}^5 . Find a basis for $\text{span}(S)$.

Sol. Step 1.

$$A = \begin{bmatrix} 1 & 2 & 3 & -1 & 0 \\ 3 & 6 & 8 & -2 & 0 \\ -1 & -1 & -3 & 1 & 1 \\ -2 & -3 & -5 & 1 & 1 \end{bmatrix}$$



Step 2.



Step 2.

$$C = \text{RREF}(A) = \begin{bmatrix} \boxed{1} & 0 & 0 & 2 & -2 \\ 0 & \boxed{1} & 0 & 0 & 1 \\ 0 & 0 & \boxed{1} & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$



Step 2.

$$C = \text{RREF}(A) = \begin{bmatrix} \boxed{1} & 0 & 0 & 2 & -2 \\ 0 & \boxed{1} & 0 & 0 & 1 \\ 0 & 0 & \boxed{1} & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Step 3.



Step 2.

$$C = \text{RREF}(A) = \begin{bmatrix} \boxed{1} & 0 & 0 & 2 & -2 \\ 0 & \boxed{1} & 0 & 0 & 1 \\ 0 & 0 & \boxed{1} & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Step 3.

Corresponding to non-zero rows,



Step 2.

$$C = \text{RREF}(A) = \begin{bmatrix} \boxed{1} & 0 & 0 & 2 & -2 \\ 0 & \boxed{1} & 0 & 0 & 1 \\ 0 & 0 & \boxed{1} & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Step 3.

Corresponding to non-zero rows,

$$B = \{[1, 0, 0, 2, -2], [0, 1, 0, 0, 1], [0, 0, 1, -1, 0]\}$$



Step 2.

$$C = \text{RREF}(A) = \begin{bmatrix} \boxed{1} & 0 & 0 & 2 & -2 \\ 0 & \boxed{1} & 0 & 0 & 1 \\ 0 & 0 & \boxed{1} & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Step 3.

Corresponding to non-zero rows,

$B = \{[1, 0, 0, 2, -2], [0, 1, 0, 0, 1], [0, 0, 1, -1, 0]\}$ is a basis for $\text{span}(S)$.



Exercise

Q:. Let

$S = \{x^3 - 3x^2 + 2, 2x^3 - 7x^2 + x - 3, 4x^3 - 13x^2 + x + 5\}$ be a subset of P_3 . Use Simplified Span Method to find a basis for $\text{span}(S)$.



Exercise

Q:. Let

$S = \{x^3 - 3x^2 + 2, 2x^3 - 7x^2 + x - 3, 4x^3 - 13x^2 + x + 5\}$ be a subset of P_3 . Use Simplified Span Method to find a basis for $\text{span}(S)$.

Sol. $B = \{x^3 - 3x, x^2 - x, 1\}$



Next is to reduce a spanning set to a basis



Next is to reduce a spanning set to a basis by eliminating redundant vectors



Next is to reduce a spanning set to a basis by eliminating redundant vectors without changing the form of the original vectors.



Next is to reduce a spanning set to a basis by eliminating redundant vectors without changing the form of the original vectors. How?



Next is to reduce a spanning set to a basis by eliminating redundant vectors without changing the form of the original vectors. How?

Theorem: If S is a spanning set for a finite dimensional vector space V , then there is a set $B \subseteq S$ that is a basis for V .



Independence Test Method to find a Basis for $\text{span}(S)$



Independence Test Method to find a Basis for $\text{span}(S)$

Let $S \subseteq \mathbb{R}^n$ containing k vectors.



Independence Test Method to find a Basis for $\text{span}(S)$

Let $S \subseteq \mathbb{R}^n$ containing k vectors.

- Construct a matrix A of order $n \times k$ by using vectors of S



Independence Test Method to find a Basis for $\text{span}(S)$

Let $S \subseteq \mathbb{R}^n$ containing k vectors.

- Construct a matrix A of order $n \times k$ by using vectors of S as columns of A .



Independence Test Method to find a Basis for $\text{span}(S)$

Let $S \subseteq \mathbb{R}^n$ containing k vectors.

- Construct a matrix A of order $n \times k$ by using vectors of S as columns of A .
- Compute $C = \text{RREF}(A)$.



Independence Test Method to find a Basis for $\text{span}(S)$

Let $S \subseteq \mathbb{R}^n$ containing k vectors.

- Construct a matrix A of order $n \times k$ by using vectors of S as columns of A .
- Compute $C = \text{RREF}(A)$.
- Vectors corresponding to pivot columns of C forms a basis for $\text{span}(S)$.



Example

Q:. Let $S = \{[1, 2, -2, 1], [-3, 0, -4, 3], [2, 1, 1, -1], [-3, 3, -9, 6], [9, 3, 7, -6]\}$ be a subset of \mathbb{R}^4 .



Example

Q:. Let $S = \{[1, 2, -2, 1], [-3, 0, -4, 3], [2, 1, 1, -1], [-3, 3, -9, 6], [9, 3, 7, -6]\}$ be a subset of \mathbb{R}^4 . Find a basis for $\text{span}(S)$.



Example

Q:. Let $S = \{[1, 2, -2, 1], [-3, 0, -4, 3], [2, 1, 1, -1], [-3, 3, -9, 6], [9, 3, 7, -6]\}$ be a subset of \mathbb{R}^4 . Find a basis for $\text{span}(S)$.

Sol. Step 1.



Example

Q.: Let $S = \{[1, 2, -2, 1], [-3, 0, -4, 3], [2, 1, 1, -1], [-3, 3, -9, 6], [9, 3, 7, -6]\}$ be a subset of \mathbb{R}^4 . Find a basis for $\text{span}(S)$.

Sol. Step 1.

$$A = \begin{bmatrix} 1 & -3 & 2 & -3 & 9 \\ 2 & 0 & 1 & 3 & 3 \\ -2 & -4 & 1 & -9 & 7 \\ 1 & 3 & -1 & 6 & -6 \end{bmatrix}$$



Step 2.



Step 2.

$$C = \text{RREF}(A) = \begin{bmatrix} \boxed{1} & 0 & 1/2 & 3/2 & 3/2 \\ 0 & \boxed{1} & -1/2 & 3/2 & -5/2 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$



Step 2.

$$C = \text{RREF}(A) = \begin{bmatrix} \boxed{1} & 0 & 1/2 & 3/2 & 3/2 \\ 0 & \boxed{1} & -1/2 & 3/2 & -5/2 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Step 3.



Step 2.

$$C = \text{RREF}(A) = \begin{bmatrix} \boxed{1} & 0 & 1/2 & 3/2 & 3/2 \\ 0 & \boxed{1} & -1/2 & 3/2 & -5/2 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Step 3.

Corresponding to pivot columns,



Step 2.

$$C = \text{RREF}(A) = \begin{bmatrix} \boxed{1} & 0 & 1/2 & 3/2 & 3/2 \\ 0 & \boxed{1} & -1/2 & 3/2 & -5/2 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Step 3.

Corresponding to pivot columns,

$B = \{[1, 2, -2, 1], [-3, 0, -4, 3]\}$ forms a basis for $\text{span}(S)$.



Exercise

Q:. Let $S = \{x^3 - 3x^2 + 1, 2x^2 + x, 2x^3 + 3x + 2, 4x - 5\}$ be a subset of P_3 . Use Independence Test Method to find a basis for $\text{span}(S)$.



Exercise

Q.: Let $S = \{x^3 - 3x^2 + 1, 2x^2 + x, 2x^3 + 3x + 2, 4x - 5\}$ be a subset of P_3 . Use Independence Test Method to find a basis for $\text{span}(S)$.

Sol. $B = \{x^3 - 3x^2 + 1, 2x^2 + x, 4x - 5\}$



Exercise

Q:. Let $S = \{x^3 - 3x^2 + 1, 2x^2 + x, 2x^3 + 3x + 2, 4x - 5\}$ be a subset of P_3 . Use Independence Test Method to find a basis for $\text{span}(S)$.

Sol. $B = \{x^3 - 3x^2 + 1, 2x^2 + x, 4x - 5\}$

Q:. Does Simplified Span Method give a basis B for $\text{span}(S)$ such that $B \subset S$.



Exercise

Q.: Let $S = \{x^3 - 3x^2 + 1, 2x^2 + x, 2x^3 + 3x + 2, 4x - 5\}$ be a subset of P_3 . Use Independence Test Method to find a basis for $\text{span}(S)$.

Sol. $B = \{x^3 - 3x^2 + 1, 2x^2 + x, 4x - 5\}$

Q.: Does Simplified Span Method give a basis B for $\text{span}(S)$ such that $B \subset S$.

Sol. No. Counterexample

$$\begin{bmatrix} 1 & 0 & 0 \\ 2 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$



Exercise

Q.: Let $S = \{x^3 - 3x^2 + 1, 2x^2 + x, 2x^3 + 3x + 2, 4x - 5\}$ be a subset of P_3 . Use Independence Test Method to find a basis for $\text{span}(S)$.

Sol. $B = \{x^3 - 3x^2 + 1, 2x^2 + x, 4x - 5\}$

Q.: Does Simplified Span Method give a basis B for $\text{span}(S)$ such that $B \subset S$.

Sol. No. Counterexample

$$\begin{bmatrix} 1 & 0 & 0 \\ 2 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$



Example

Q:. Let $T = \{[1, 0, 1, 0], [-1, 1, -1, 0]\}$ be a LI subset of \mathbb{R}^4 .



Example

Q:. Let $T = \{[1, 0, 1, 0], [-1, 1, -1, 0]\}$ be a LI subset of \mathbb{R}^4 . Extend T to form a basis for \mathbb{R}^4 .



Example

Q:. Let $T = \{[1, 0, 1, 0], [-1, 1, -1, 0]\}$ be a LI subset of \mathbb{R}^4 . Extend T to form a basis for \mathbb{R}^4 .

Why T is not a basis for \mathbb{R}^4 .



Example

Q:. Let $T = \{[1, 0, 1, 0], [-1, 1, -1, 0]\}$ be a LI subset of \mathbb{R}^4 . Extend T to form a basis for \mathbb{R}^4 .

Why T is not a basis for \mathbb{R}^4 . How can we extend T to form a basis for \mathbb{R}^4 .



Example

Q:. Let $T = \{[1, 0, 1, 0], [-1, 1, -1, 0]\}$ be a LI subset of \mathbb{R}^4 . Extend T to form a basis for \mathbb{R}^4 .

Why T is not a basis for \mathbb{R}^4 . How can we extend T to form a basis for \mathbb{R}^4 .

Sol. $\text{span}(T) = \{a[1, 0, 1, 0] + b[-1, 1, -1, 0] | a, b \in \mathbb{R}\} \implies$



Example

Q.: Let $T = \{[1, 0, 1, 0], [-1, 1, -1, 0]\}$ be a LI subset of \mathbb{R}^4 . Extend T to form a basis for \mathbb{R}^4 .

Why T is not a basis for \mathbb{R}^4 . How can we extend T to form a basis for \mathbb{R}^4 .

Sol. $\text{span}(T) = \{a[1, 0, 1, 0] + b[-1, 1, -1, 0] | a, b \in \mathbb{R}\} \implies$
 $\text{span}(T) = \{[a - b, b, a - b, 0] | a, b \in \mathbb{R}\}.$



Example

Q:. Let $T = \{[1, 0, 1, 0], [-1, 1, -1, 0]\}$ be a LI subset of \mathbb{R}^4 . Extend T to form a basis for \mathbb{R}^4 .

Why T is not a basis for \mathbb{R}^4 . How can we extend T to form a basis for \mathbb{R}^4 .

Sol. $\text{span}(T) = \{a[1, 0, 1, 0] + b[-1, 1, -1, 0] | a, b \in \mathbb{R}\} \implies$
 $\text{span}(T) = \{[a - b, b, a - b, 0] | a, b \in \mathbb{R}\}.$

Now $[0, 0, 0, 1] \notin \text{span}(T) \implies$



Example

Q:. Let $T = \{[1, 0, 1, 0], [-1, 1, -1, 0]\}$ be a LI subset of \mathbb{R}^4 . Extend T to form a basis for \mathbb{R}^4 .

Why T is not a basis for \mathbb{R}^4 . How can we extend T to form a basis for \mathbb{R}^4 .

Sol. $\text{span}(T) = \{a[1, 0, 1, 0] + b[-1, 1, -1, 0] | a, b \in \mathbb{R}\} \implies$
 $\text{span}(T) = \{[a - b, b, a - b, 0] | a, b \in \mathbb{R}\}.$

Now $[0, 0, 0, 1] \notin \text{span}(T) \implies$

$S = \{[1, 0, 1, 0], [-1, 1, -1, 0], [0, 0, 0, 1]\}$ is a LI a subset of \mathbb{R}^4 .



Similarly, we can verify that $[0, 0, 1, 0] \notin \text{span}(S) \implies$



Similarly, we can verify that $[0, 0, 1, 0] \notin \text{span}(S) \implies$
 $B = \{[1, 0, 1, 0], [-1, 1, -1, 0], [0, 0, 0, 1], [0, 0, 1, 0]\}$ is a LI
a subset of \mathbb{R}^4 .



Similarly, we can verify that $[0, 0, 1, 0] \notin \text{span}(S) \implies$
 $B = \{[1, 0, 1, 0], [-1, 1, -1, 0], [0, 0, 0, 1], [0, 0, 1, 0]\}$ is a LI
a subset of \mathbb{R}^4 .

Hence, B is a basis of \mathbb{R}^4 containing T .



Similarly, we can verify that $[0, 0, 1, 0] \notin \text{span}(S) \implies$
 $B = \{[1, 0, 1, 0], [-1, 1, -1, 0], [0, 0, 0, 1], [0, 0, 1, 0]\}$ is a LI
a subset of \mathbb{R}^4 .

Hence, B is a basis of \mathbb{R}^4 containing T .

Another approach



Similarly, we can verify that $[0, 0, 1, 0] \notin \text{span}(S) \implies B = \{[1, 0, 1, 0], [-1, 1, -1, 0], [0, 0, 0, 1], [0, 0, 1, 0]\}$ is a LI subset of \mathbb{R}^4 .

Hence, B is a basis of \mathbb{R}^4 containing T .

Another approach

Sol. Let $T = \{v_1, v_2\}$.



Similarly, we can verify that $[0, 0, 1, 0] \notin \text{span}(S) \implies B = \{[1, 0, 1, 0], [-1, 1, -1, 0], [0, 0, 0, 1], [0, 0, 1, 0]\}$ is a LI subset of \mathbb{R}^4 .

Hence, B is a basis of \mathbb{R}^4 containing T .

Another approach

Sol. Let $T = \{v_1, v_2\}$. We know that $A = \{e_1, e_2, e_3, e_4\}$ is a standard basis for \mathbb{R}^4 .



Similarly, we can verify that $[0, 0, 1, 0] \notin \text{span}(S) \implies B = \{[1, 0, 1, 0], [-1, 1, -1, 0], [0, 0, 0, 1], [0, 0, 1, 0]\}$ is a LI subset of \mathbb{R}^4 .

Hence, B is a basis of \mathbb{R}^4 containing T .

Another approach

Sol. Let $T = \{v_1, v_2\}$. We know that $A = \{e_1, e_2, e_3, e_4\}$ is a standard basis for \mathbb{R}^4 . Let $S = \{v_1, v_2, e_1, e_2, e_3, e_4\}$.



Similarly, we can verify that $[0, 0, 1, 0] \notin \text{span}(S) \implies B = \{[1, 0, 1, 0], [-1, 1, -1, 0], [0, 0, 0, 1], [0, 0, 1, 0]\}$ is a LI subset of \mathbb{R}^4 .

Hence, B is a basis of \mathbb{R}^4 containing T .

Another approach

Sol. Let $T = \{v_1, v_2\}$. We know that $A = \{e_1, e_2, e_3, e_4\}$ is a standard basis for \mathbb{R}^4 . Let $S = \{v_1, v_2, e_1, e_2, e_3, e_4\}$.

Now A spans $\mathbb{R}^4 \implies$



Similarly, we can verify that $[0, 0, 1, 0] \notin \text{span}(S) \implies B = \{[1, 0, 1, 0], [-1, 1, -1, 0], [0, 0, 0, 1], [0, 0, 1, 0]\}$ is a LI subset of \mathbb{R}^4 .

Hence, B is a basis of \mathbb{R}^4 containing T .

Another approach

Sol. Let $T = \{v_1, v_2\}$. We know that $A = \{e_1, e_2, e_3, e_4\}$ is a standard basis for \mathbb{R}^4 . Let $S = \{v_1, v_2, e_1, e_2, e_3, e_4\}$.

Now A spans $\mathbb{R}^4 \implies S$ spans \mathbb{R}^4 , i.e., $\mathbb{R}^4 = \text{span}(S)$.



Using Independence Test Method we have



Using Independence Test Method we have

$$C = \begin{bmatrix} 1 & -1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 1 & -1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \Rightarrow$$



Using Independence Test Method we have

$$C = \begin{bmatrix} 1 & -1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 1 & -1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \Rightarrow$$

$$\text{RREF}(C) = \begin{bmatrix} \boxed{1} & 0 & 0 & 1 & 1 & 0 \\ 0 & \boxed{1} & 0 & 1 & 0 & 0 \\ 0 & 0 & \boxed{1} & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & \boxed{1} \end{bmatrix} \Rightarrow$$



Using Independence Test Method we have

$$C = \begin{bmatrix} 1 & -1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 1 & -1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \Rightarrow$$

$$\text{RREF}(C) = \begin{bmatrix} \boxed{1} & 0 & 0 & 1 & 1 & 0 \\ 0 & \boxed{1} & 0 & 1 & 0 & 0 \\ 0 & 0 & \boxed{1} & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & \boxed{1} \end{bmatrix} \Rightarrow$$

$B = \{v_1, v_2, e_1, e_4\}$ is a basis of \mathbb{R}^4 containing T .



Theorem: Every LI subset of a finite dimensional vector space V can be extended to form a basis of V .



Theorem: Every LI subset of a finite dimensional vector space V can be extended to form a basis of V .

Recall



Theorem: Every LI subset of a finite dimensional vector space V can be extended to form a basis of V .

Recall

Theorem: Let V be a finite dimensional vector space such that $\dim(V) = n$.



Theorem: Every LI subset of a finite dimensional vector space V can be extended to form a basis of V .

Recall

Theorem: Let V be a finite dimensional vector space such that $\dim(V) = n$. Suppose T is a linearly independent subset of V . Then T is finite and $|T| \leq n$.



Theorem: Every LI subset of a finite dimensional vector space V can be extended to form a basis of V .

Recall

Theorem: Let V be a finite dimensional vector space such that $\dim(V) = n$. Suppose T is a linearly independent subset of V . Then T is finite and $|T| \leq n$. Moreover, $|T| = n$ if and only if T is a basis for V .



Theorem: Every LI subset of a finite dimensional vector space V can be extended to form a basis of V .

Recall

Theorem: Let V be a finite dimensional vector space such that $\dim(V) = n$. Suppose T is a linearly independent subset of V . Then T is finite and $|T| \leq n$. Moreover, $|T| = n$ if and only if T is a basis for V .

Theorem: Let W be a subspace of a finite dimensional vector space V .



Theorem: Every LI subset of a finite dimensional vector space V can be extended to form a basis of V .

Recall

Theorem: Let V be a finite dimensional vector space such that $\dim(V) = n$. Suppose T is a linearly independent subset of V . Then T is finite and $|T| \leq n$. Moreover, $|T| = n$ if and only if T is a basis for V .

Theorem: Let W be a subspace of a finite dimensional vector space V . Then W is also finite dimensional and $\dim(W) \leq \dim(V)$.



Theorem: Every LI subset of a finite dimensional vector space V can be extended to form a basis of V .

Recall

Theorem: Let V be a finite dimensional vector space such that $\dim(V) = n$. Suppose T is a linearly independent subset of V . Then T is finite and $|T| \leq n$. Moreover, $|T| = n$ if and only if T is a basis for V .

Theorem: Let W be a subspace of a finite dimensional vector space V . Then W is also finite dimensional and $\dim(W) \leq \dim(V)$. $\dim(W) = \dim(V)$ if and only if $W = V$.



Method for finding a Basis by Enlarging a LI Subset



Method for finding a Basis by Enlarging a LI Subset

Let $T = \{t_1, \dots, t_k\}$ be a LI subset of a finite dimensional vector space V .



Method for finding a Basis by Enlarging a LI Subset

Let $T = \{t_1, \dots, t_k\}$ be a LI subset of a finite dimensional vector space V .

- Find a spanning set $A = \{a_1, \dots, a_n\}$ for V .



Method for finding a Basis by Enlarging a LI Subset

Let $T = \{t_1, \dots, t_k\}$ be a LI subset of a finite dimensional vector space V .

- Find a spanning set $A = \{a_1, \dots, a_n\}$ for V .
- Form the ordered spanning set



Method for finding a Basis by Enlarging a LI Subset

Let $T = \{t_1, \dots, t_k\}$ be a LI subset of a finite dimensional vector space V .

- Find a spanning set $A = \{a_1, \dots, a_n\}$ for V .
- Form the ordered spanning set $S = \{t_1, \dots, t_k, a_1, \dots, a_n\}$ for V .



Method for finding a Basis by Enlarging a LI Subset

Let $T = \{t_1, \dots, t_k\}$ be a LI subset of a finite dimensional vector space V .

- Find a spanning set $A = \{a_1, \dots, a_n\}$ for V .
- Form the ordered spanning set $S = \{t_1, \dots, t_k, a_1, \dots, a_n\}$ for V .
- Use Independence Test Method to produce a subset B of S .



Method for finding a Basis by Enlarging a LI Subset

Let $T = \{t_1, \dots, t_k\}$ be a LI subset of a finite dimensional vector space V .

- Find a spanning set $A = \{a_1, \dots, a_n\}$ for V .
- Form the ordered spanning set $S = \{t_1, \dots, t_k, a_1, \dots, a_n\}$ for V .
- Use Independence Test Method to produce a subset B of S . Then B is a basis for V containing T .



Example

Q.: Let $T = \{x^3 - x^2, x^4 - 3x^3 + 5x^2 - x\}$ be a LI subset of P_4 .



Example

Q:. Let $T = \{x^3 - x^2, x^4 - 3x^3 + 5x^2 - x\}$ be a LI subset of P_4 . Extend T to form a basis for P_4 .



Example

Q:. Let $T = \{x^3 - x^2, x^4 - 3x^3 + 5x^2 - x\}$ be a LI subset of P_4 . Extend T to form a basis for P_4 .

Sol. We first convert polynomials in T to vectors in \mathbb{R}^5 ,



Example

Q:. Let $T = \{x^3 - x^2, x^4 - 3x^3 + 5x^2 - x\}$ be a LI subset of P_4 . Extend T to form a basis for P_4 .

Sol. We first convert polynomials in T to vectors in \mathbb{R}^5 , i.e., $T = \{[0, 1, -1, 0, 0], [1, -3, 5, -1, 0]\}$.



Example

Q:. Let $T = \{x^3 - x^2, x^4 - 3x^3 + 5x^2 - x\}$ be a LI subset of P_4 . Extend T to form a basis for P_4 .

Sol. We first convert polynomials in T to vectors in \mathbb{R}^5 , i.e., $T = \{[0, 1, -1, 0, 0], [1, -3, 5, -1, 0]\}$. Let $T = \{v_1, v_2\}$.



Example

Q:. Let $T = \{x^3 - x^2, x^4 - 3x^3 + 5x^2 - x\}$ be a LI subset of P_4 . Extend T to form a basis for P_4 .

Sol. We first convert polynomials in T to vectors in \mathbb{R}^5 , i.e., $T = \{[0, 1, -1, 0, 0], [1, -3, 5, -1, 0]\}$. Let $T = \{v_1, v_2\}$.

Step 1.



Example

Q.: Let $T = \{x^3 - x^2, x^4 - 3x^3 + 5x^2 - x\}$ be a LI subset of P_4 . Extend T to form a basis for P_4 .

Sol. We first convert polynomials in T to vectors in \mathbb{R}^5 , i.e., $T = \{[0, 1, -1, 0, 0], [1, -3, 5, -1, 0]\}$. Let $T = \{v_1, v_2\}$.

Step 1. $A = \{e_1, e_2, e_3, e_4, e_5\}$ is a standard basis for \mathbb{R}^5 ,



Example

Q.: Let $T = \{x^3 - x^2, x^4 - 3x^3 + 5x^2 - x\}$ be a LI subset of P_4 . Extend T to form a basis for P_4 .

Sol. We first convert polynomials in T to vectors in \mathbb{R}^5 , i.e., $T = \{[0, 1, -1, 0, 0], [1, -3, 5, -1, 0]\}$. Let $T = \{v_1, v_2\}$.

Step 1. $A = \{e_1, e_2, e_3, e_4, e_5\}$ is a standard basis for \mathbb{R}^5 , hence it is a spanning set for \mathbb{R}^5 .



Example

Q.: Let $T = \{x^3 - x^2, x^4 - 3x^3 + 5x^2 - x\}$ be a LI subset of P_4 . Extend T to form a basis for P_4 .

Sol. We first convert polynomials in T to vectors in \mathbb{R}^5 , i.e., $T = \{[0, 1, -1, 0, 0], [1, -3, 5, -1, 0]\}$. Let $T = \{v_1, v_2\}$.

Step 1. $A = \{e_1, e_2, e_3, e_4, e_5\}$ is a standard basis for \mathbb{R}^5 , hence it is a spanning set for \mathbb{R}^5 .

Step 2.



Example

Q.: Let $T = \{x^3 - x^2, x^4 - 3x^3 + 5x^2 - x\}$ be a LI subset of P_4 . Extend T to form a basis for P_4 .

Sol. We first convert polynomials in T to vectors in \mathbb{R}^5 , i.e., $T = \{[0, 1, -1, 0, 0], [1, -3, 5, -1, 0]\}$. Let $T = \{v_1, v_2\}$.

Step 1. $A = \{e_1, e_2, e_3, e_4, e_5\}$ is a standard basis for \mathbb{R}^5 , hence it is a spanning set for \mathbb{R}^5 .

Step 2. The ordered spanning set is



Example

Q.: Let $T = \{x^3 - x^2, x^4 - 3x^3 + 5x^2 - x\}$ be a LI subset of P_4 . Extend T to form a basis for P_4 .

Sol. We first convert polynomials in T to vectors in \mathbb{R}^5 , i.e., $T = \{[0, 1, -1, 0, 0], [1, -3, 5, -1, 0]\}$. Let $T = \{v_1, v_2\}$.

Step 1. $A = \{e_1, e_2, e_3, e_4, e_5\}$ is a standard basis for \mathbb{R}^5 , hence it is a spanning set for \mathbb{R}^5 .

Step 2. The ordered spanning set is $S = \{v_1, v_2, e_1, e_2, e_3, e_4, e_5\}$.



Step 3.



Step 3. Using Independence Test Method we have



Step 3. Using Independence Test Method we have

$$C = \begin{bmatrix} 0 & 1 & 1 & 0 & 0 & 0 & 0 \\ 1 & -3 & 0 & 1 & 0 & 0 & 0 \\ -1 & 5 & 0 & 0 & 1 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$



Step 3. Using Independence Test Method we have

$$C = \begin{bmatrix} 0 & 1 & 1 & 0 & 0 & 0 & 0 \\ 1 & -3 & 0 & 1 & 0 & 0 & 0 \\ -1 & 5 & 0 & 0 & 1 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Step 4.



Step 3. Using Independence Test Method we have

$$C = \begin{bmatrix} 0 & 1 & 1 & 0 & 0 & 0 & 0 \\ 1 & -3 & 0 & 1 & 0 & 0 & 0 \\ -1 & 5 & 0 & 0 & 1 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\text{Step 4. RREF}(C) = \begin{bmatrix} \boxed{1} & 0 & 0 & 0 & -1 & -5 & 0 \\ 0 & \boxed{1} & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & \boxed{1} & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & \boxed{1} & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \boxed{1} \end{bmatrix}$$



Step 5.



Step 5. Since columns I, II, III, IV and VII are pivot columns,



Step 5. Since columns I, II, III, IV and VII are pivot columns, by independent test method, the set $B = \{v_1, v_2, e_1, e_2, e_5\}$ is the basis of \mathbb{R}^5 ,



Step 5. Since columns I, II, III, IV and VII are pivot columns, by independent test method, the set $B = \{v_1, v_2, e_1, e_2, e_5\}$ is the basis of \mathbb{R}^5 ,

i.e., $B = \{x^3 - x^2, x^4 - 3x^3 + 5x^2 - x, x^4, x^3, 1\}$ is the basis of P_4 containing T .



Exercise

Q:. Let $T = \left\{ \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & -1 \end{bmatrix} \right\}$ be a LI subset of M_{32} .



Exercise

Q:. Let $T = \left\{ \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & -1 \end{bmatrix} \right\}$ be a LI subset of M_{32} . Extend T to form a basis for M_{32} .



Exercise

Q:. Let $T = \left\{ \begin{bmatrix} 1 & -1 \\ -1 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & -1 \\ -1 & 1 \end{bmatrix} \right\}$ be a LI subset of M_{32} . Extend T to form a basis for M_{32} .

Sol.

$$B = \left\{ \begin{bmatrix} 1 & -1 \\ -1 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & -1 \\ -1 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 1 & 0 \end{bmatrix} \right\}$$

