

# MATHEMATICS-I (MATH F111)

**Dr. Krishnendra Shekhawat**

**BITS PILANI**  
**Department of Mathematics**



# Points of intersection



## Point of intersection of two curves

Consider two curves  $r = f(\theta)$  and  $r = g(\theta)$ .



## Point of intersection of two curves

Consider two curves  $r = f(\theta)$  and  $r = g(\theta)$ . A point  $(r, \theta)$  which lies on both curves is called a **point of intersection** of the curves  $r = f(\theta)$  and  $r = g(\theta)$ .



## Point of intersection of two curves

Consider two curves  $r = f(\theta)$  and  $r = g(\theta)$ . A point  $(r, \theta)$  which lies on both curves is called a **point of intersection** of the curves  $r = f(\theta)$  and  $r = g(\theta)$ .

**Q:.** How to compute the points of intersection of two curves?



# Method to find the points of intersection

**Q:.** Find the points of intersection of  $r = 1 + \cos \theta$  and  $r = 1 - \cos \theta$ .



# Method to find the points of intersection

**Q:.** Find the points of intersection of  $r = 1 + \cos \theta$  and  $r = 1 - \cos \theta$ .

**Step 1 :** Solve the simultaneous equations by eliminating one variable (preferably  $r$ ).



# Method to find the points of intersection

**Q:.** Find the points of intersection of  $r = 1 + \cos \theta$  and  $r = 1 - \cos \theta$ .

**Step 1 :** Solve the simultaneous equations by eliminating one variable (preferably  $r$ ).

Putting both values of  $r$  equal and solve for  $\theta$ :

$$1 + \cos \theta = 1 - \cos \theta,$$

which gives

$$\cos \theta = 0 \text{ and so } \theta = \frac{\pi}{2}, \frac{3\pi}{2}.$$





Now substitute  $\theta = \frac{\pi}{2}$  in first equation to get  $r = 1$  (you can substitute in second equation also).



Now substitute  $\theta = \frac{\pi}{2}$  in first equation to get  $r = 1$  (you can substitute in second equation also). Thus one point of intersection is  $(1, \frac{\pi}{2})$ .



Now substitute  $\theta = \frac{\pi}{2}$  in first equation to get  $r = 1$  (you can substitute in second equation also). Thus one point of intersection is  $\left(1, \frac{\pi}{2}\right)$ .

Now substitute  $\theta = \frac{3\pi}{2}$  in first equation to get  $r = 1$ . Thus the second point of intersection is  $\left(1, \frac{3\pi}{2}\right)$ .



Now substitute  $\theta = \frac{\pi}{2}$  in first equation to get  $r = 1$  (you can substitute in second equation also). Thus one point of intersection is  $(1, \frac{\pi}{2})$ .

Now substitute  $\theta = \frac{3\pi}{2}$  in first equation to get  $r = 1$ . Thus the second point of intersection is  $(1, \frac{3\pi}{2})$ .

**Q:.** Are there more points of intersection?



Now substitute  $\theta = \frac{\pi}{2}$  in first equation to get  $r = 1$  (you can substitute in second equation also). Thus one point of intersection is  $(1, \frac{\pi}{2})$ .

Now substitute  $\theta = \frac{3\pi}{2}$  in first equation to get  $r = 1$ . Thus the second point of intersection is  $(1, \frac{3\pi}{2})$ .

**Q:.** Are there more points of intersection? If yes, how to find them?




**Step 2 :** Plot the graphs simultaneously and see the missing points of intersection in Step 1.



**Step 2 :** Plot the graphs simultaneously and see the missing points of intersection in Step 1.

Step 1 **may not** give all points of intersection  
(because of **deceptive points**)







One more point of intersection  $(0,0)$  is found by graphing.



One more point of intersection  $(0,0)$  is found by graphing.

Thus all points of intersection are  $(0,0), \left(1, \frac{\pi}{2}\right), \left(1, \frac{3\pi}{2}\right)$ .



## Deceptive Point

A point  $(r, \theta)$  is said to be **deceptive** if it lies on the curve but does not satisfy the equation of the curve.



## Deceptive Point

A point  $(r, \theta)$  is said to be **deceptive** if it lies on the curve but does not satisfy the equation of the curve.

Another Example?



## Deceptive Point

A point  $(r, \theta)$  is said to be **deceptive** if it lies on the curve but does not satisfy the equation of the curve.

Another Example?

The point  $(0, 0)$  lies on the curve  $r = 2\cos\theta$  but does not satisfy the equation.



**Q:.** How do you verify that  $(0,0)$  is a deceptive point for the curve  $r = 2\cos\theta$ ?



**Q:.** How do you verify that  $(0,0)$  is a deceptive point for the curve  $r = 2\cos\theta$ ?

**Sol.** Another representation  $(0, \frac{\pi}{2})$  of  $(0,0)$  satisfies the equation so does lie on the curve and hence is a deceptive point.



**Q:.** Find the points of intersection of  $r = \sin \theta$  and  $r = \cos \theta$ .





**Q:.** Find the points of intersection of  $r = \sin \theta$  and  $r = \cos \theta$ .

**Sol. Step 1** Putting both values of  $r$  equal and solve for  $\theta$ :



**Q:.** Find the points of intersection of  $r = \sin \theta$  and  $r = \cos \theta$ .

**Sol. Step 1** Putting both values of  $r$  equal and solve for  $\theta$ :

$$\sin \theta = \cos \theta$$

$$\Rightarrow \theta = \frac{\pi}{4}, \frac{5\pi}{4}.$$




**Q:.** Find the points of intersection of  $r = \sin \theta$  and  $r = \cos \theta$ .


**Sol. Step 1** Putting both values of  $r$  equal and solve for  $\theta$ :

$$\begin{aligned}\sin \theta &= \cos \theta \\ \Rightarrow \theta &= \frac{\pi}{4}, \frac{5\pi}{4}.\end{aligned}$$

**Step 2.** Plot the graphs simultaneously to find the missing points of intersection in Step 1








The points of intersection are

$$\left(1/\sqrt{2}, \frac{\pi}{4}\right) \equiv \left(-1/\sqrt{2}, \frac{5\pi}{4}\right), (0,0)$$



**Q:.** Find the points of intersection of  $r = 1$  and  $r = 2 \sin 2\theta$ .



**Step 1: The points of intersection by solving the equations.**

$$\left(1, \frac{7\pi}{12}\right), \left(1, \frac{11\pi}{12}\right), \left(1, \frac{19\pi}{12}\right), \left(1, \frac{23\pi}{12}\right),$$

**Step 2: The points of intersection found by graphing and symmetry.**

$$\left(1, \frac{7\pi}{12}\right), \left(1, \frac{11\pi}{12}\right), \left(1, \frac{19\pi}{12}\right), \left(1, \frac{23\pi}{12}\right).$$



# Section 11.5

## Areas and Lengths in Polar Coordinates






Q:.. How to compute the area enclosed by a curve?



Q:.. How to compute the area enclosed by a curve?

Q:.. Precisely, how to compute the area enclosed by a curve  $r = f(\theta)$  and by the rays  $\theta = \alpha$  and  $\theta = \beta$ ?





To derive a formula for the area  $A$  of region  $OTS$ , we approximate the region with  $n$  nonoverlapping fan shaped circular sectors based on a partition  $P$  of angle  $TOS$ . The typical sector has radius  $r_k$  and central angle  $\Delta\theta_k$ .



To derive a formula for the area  $A$  of region  $OTS$ , we approximate the region with  $n$  nonoverlapping fan shaped circular sectors based on a partition  $P$  of angle  $TOS$ . The typical sector has radius  $r_k$  and central angle  $\Delta\theta_k$ .

Therefore the circular sector (with an angle  $\Delta\theta_k$ ) is a part of circle of radius  $r_k$ . Thus the area of this sector is

$$A_k = \frac{\Delta\theta_k}{2\pi}(\pi r_k^2) = \frac{1}{2}r_k^2\Delta\theta_k.$$



Then the area of region  $OTS$  is approximately

$$\sum_{k=1}^n A_k = \sum_{k=1}^n \frac{1}{2} r_k^2 \Delta\theta_k.$$



Then the area of region  $OTS$  is approximately

$$\sum_{k=1}^n A_k = \sum_{k=1}^n \frac{1}{2} r_k^2 \Delta\theta_k.$$

Thus the area  $A$  of  $OTS$  can be given by taking  $n \rightarrow \infty$ .

Thus



Then the area of region  $OTS$  is approximately

$$\sum_{k=1}^n A_k = \sum_{k=1}^n \frac{1}{2} r_k^2 \Delta\theta_k.$$

Thus the area  $A$  of  $OTS$  can be given by taking  $n \rightarrow \infty$ .

Thus

$$A = \lim_{n \rightarrow \infty} \sum_{k=1}^n A_k = \lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{1}{2} r_k^2 \Delta\theta_k = \frac{1}{2} \int_{\alpha}^{\beta} r^2 d\theta.$$





Then the area  $A$  of fan shaped region  $OTS$  between the origin and the curve  $r = f(\theta)$  and also between the rays  $\theta = \alpha$  and  $\theta = \beta$  is given by (assuming  $\alpha \leq \beta$ )

$$A = \frac{1}{2} \int_{\alpha}^{\beta} r^2 d\theta.$$



## Example


Find the area of the region inside the lemniscate  
 $r^2 = 2a^2 \cos 2\theta$ ,  $a > 0$ .



## Example

Find the area of the region inside the lemniscate  
 $r^2 = 2a^2 \cos 2\theta$ ,  $a > 0$ .

Sol.



Using symmetries, the required area is given by



Using symmetries, the required area is given by

$$A = 4 \times \text{Area in first quadrant}$$

$$= 4 \times \frac{1}{2} \int_0^{\frac{\pi}{4}} r^2 d\theta$$

$$= 2 \int_0^{\frac{\pi}{4}} 2a^2 \cos 2\theta d\theta$$

$$= 2a^2.$$



Find the area of the region inside the circle  $r = 2 \cos \theta$ .



Find the area of the region inside the circle  $r = 2 \cos \theta$ .

**Sol.**

$$\begin{aligned} A &= \frac{1}{2} \int_0^{2\pi} r^2 d\theta = 2 \times \frac{1}{2} \int_0^{\pi} 4 \cos^2 \theta d\theta \\ &= 2 \int_0^{\pi} 1 + \cos 2\theta d\theta \\ &= 2\pi. \end{aligned}$$



Find the area of the region inside the circle  $r = 2 \cos \theta$ .


**Sol.**

$$\begin{aligned} A &= \frac{1}{2} \int_0^{2\pi} r^2 d\theta = 2 \times \frac{1}{2} \int_0^{\pi} 4 \cos^2 \theta d\theta \\ &= 2 \int_0^{\pi} 1 + \cos 2\theta d\theta \\ &= 2\pi. \end{aligned}$$

**Note:**  $r = 2 \cos \theta$  is a circle with center at  $(1, 0)$  and radius 1. Hence, its actual area is  $\pi$ .







**Figure:**  $r = 2\cos\theta$



The flaw happens because we have not traced the curve. When we trace the curve we find that the whole circle traces between  $0$  to  $\pi$ , i.e., it traces twice from  $0$  to  $2\pi$ , which leads us to the area twice as the actual area.



The flaw happens because we have not traced the curve. When we trace the curve we find that the whole circle traces between  $0$  to  $\pi$ , i.e., it traces twice from  $0$  to  $2\pi$ , which leads us to the area twice as the actual area.

$A = 2 \times \text{Area in first quadrant}$

$$\begin{aligned} &= 2 \times \frac{1}{2} \int_0^{\pi/2} r^2 d\theta = \int_0^{\pi/2} 4 \cos^2 \theta d\theta \\ &= 2 \int_0^{\pi/2} 1 + \cos 2\theta d\theta \\ &= \pi. \end{aligned}$$



# Steps to Evaluate Area

- Plot the polar graphs correctly and clearly.



# Steps to Evaluate Area

- Plot the polar graphs correctly and clearly.
- Label the relevant curves by their equations.



# Steps to Evaluate Area

- Plot the polar graphs correctly and clearly.
- Label the relevant curves by their equations.
- Shade the area required by marking its angular spread and radial spread. You may require to find intersections or tangents at pole.



# Steps to Evaluate Area


- Plot the polar graphs correctly and clearly.
- Label the relevant curves by their equations.
- Shade the area required by marking its angular spread and radial spread. You may require to find intersections or tangents at pole.
- See that area is covered exactly once by radial segment. Justify symmetries if used.



Q: Find the area of the region inside  $r^2 = -\sin \theta$ .







Using symmetries, the required area is given by



Using symmetries, the required area is given by

$$A = 4 \times \text{Area lying between } \pi \text{ and } 3\pi/2$$

$$= 4 \times \frac{1}{2} \int_{\pi}^{\frac{3\pi}{2}} r^2 d\theta$$


$$= 2 \int_{\pi}^{\frac{3\pi}{2}} -\sin \theta d\theta$$

$$= 2.$$



Q: Find the area of the region inside  $r^2 = \sin 2\theta$ .





Using symmetries, the required area is given by



Using symmetries, the required area is given by

$$A = 4 \times \text{Area lying between } 0 \text{ and } \pi/4$$


$$= 4 \times \frac{1}{2} \int_0^{\pi/4} r^2 d\theta$$

$$= 2 \int_0^{\pi/4} \sin 2\theta d\theta$$

$$= 1.$$



# Area Shared by Two Curves





# Area Shared by Two Curves

Area of the region  $0 \leq r_1(\theta) \leq r \leq r_2(\theta)$ ,  $\alpha \leq \theta \leq \beta$  is given by


$$A = \frac{1}{2} \int_{\alpha}^{\beta} (r_2^2 - r_1^2) d\theta.$$



**Q:.** Find the area of the region inside the circle  
 $r = 3a \cos \theta$  and outside the cardioid  
 $r = a(1 + \cos \theta)$ ,  $a > 0$ .



**Q:.** Find the area of the region inside the circle  $r = 3a \cos \theta$  and outside the cardioid  $r = a(1 + \cos \theta)$ ,  $a > 0$ .



(i) The points of intersections are  $\theta = -\frac{\pi}{3}, \frac{\pi}{3}$ .



- (i) The points of intersections are  $\theta = -\frac{\pi}{3}, \frac{\pi}{3}$ .
- (ii) Both the curves are symmetric about  $x$ -axis.



- (i) The points of intersections are  $\theta = -\frac{\pi}{3}, \frac{\pi}{3}$ .
- (ii) Both the curves are symmetric about  $x$ -axis.
- (iii) The graph also gives the point of intersection  $(0,0)$ .



- (i) The points of intersections are  $\theta = -\frac{\pi}{3}, \frac{\pi}{3}$ .
- (ii) Both the curves are symmetric about  $x$ -axis.
- (iii) The graph also gives the point of intersection  $(0,0)$ .
- (iv) Required area is the shaded area.



$A = 2 \times \text{Area in first quadrant}$


$$\begin{aligned} &= 2 \times \frac{1}{2} \int_0^{\frac{\pi}{3}} [(3a \cos \theta)^2 - a^2(1 + \cos \theta)^2] d\theta \\ &= \pi a^2. \end{aligned}$$





**Q:.** Find the area of the region inside the lemniscate  $r^2 = 6 \cos 2\theta$  and outside the circle  $r = \sqrt{3}$ .





(i) The point of intersection is  $\theta = \frac{\pi}{6}$  (in the first quadrant).



- (i) The point of intersection is  $\theta = \frac{\pi}{6}$  (in the first quadrant).
- (ii) Both the curves are symmetric about  $x$ -axis.



- (i) The point of intersection is  $\theta = \frac{\pi}{6}$  (in the first quadrant).
- (ii) Both the curves are symmetric about  $x$ -axis.
- (iii) Required area is the shaded area.



Here  $A = 4 \times$  area in first quadrant, i.e.,



Here  $A = 4 \times$  area in first quadrant, i.e.,

$$A = 4 \times \frac{1}{2} \int_0^{\frac{\pi}{6}} ((6 \cos 2\theta) - 3) d\theta$$




Here  $A = 4 \times$  area in first quadrant, i.e.,

$$\begin{aligned} A &= 4 \times \frac{1}{2} \int_0^{\frac{\pi}{6}} ((6 \cos 2\theta) - 3) d\theta \\ &= 3 \sqrt{3} - \pi. \end{aligned}$$





**Q:.** Find the area of the region shared by the cardioids  $r = 2(1 + \cos \theta)$  and  $r = 2(1 - \cos \theta)$ .



(i) The point of intersections are  $\theta = 0, \frac{\pi}{2}, \frac{3\pi}{2}$ .



- (i) The point of intersections are  $\theta = 0, \frac{\pi}{2}, \frac{3\pi}{2}$ .
- (ii) Both the curves are symmetric about  $x$ -axis.



- (i) The point of intersections are  $\theta = 0, \frac{\pi}{2}, \frac{3\pi}{2}$ .
- (ii) Both the curves are symmetric about  $x$ -axis.
- (iii) Required area is the shaded area.



# Method 1



Method 1

$$A = \left[ 2 \times \frac{1}{2} \int_0^{\frac{\pi}{2}} (2(1 - \cos \theta))^2 d\theta \right] + \left[ 2 \times \frac{1}{2} \int_{\frac{\pi}{2}}^{\pi} (2(1 + \cos \theta))^2 d\theta \right]$$



Method 1

$$\begin{aligned} A &= \left[ 2 \times \frac{1}{2} \int_0^{\frac{\pi}{2}} (2(1 - \cos \theta))^2 d\theta \right] + \left[ 2 \times \frac{1}{2} \int_{\frac{\pi}{2}}^{\pi} (2(1 + \cos \theta))^2 d\theta \right] \\ &= 6\pi - 16. \end{aligned}$$



Method 1

$$\begin{aligned} A &= \left[ 2 \times \frac{1}{2} \int_0^{\frac{\pi}{2}} (2(1 - \cos \theta))^2 d\theta \right] + \left[ 2 \times \frac{1}{2} \int_{\frac{\pi}{2}}^{\pi} (2(1 + \cos \theta))^2 d\theta \right] \\ &= 6\pi - 16. \end{aligned}$$

Method 2





Method 1

$$\begin{aligned} A &= \left[ 2 \times \frac{1}{2} \int_0^{\frac{\pi}{2}} (2(1 - \cos \theta))^2 d\theta \right] + \left[ 2 \times \frac{1}{2} \int_{\frac{\pi}{2}}^{\pi} (2(1 + \cos \theta))^2 d\theta \right] \\ &= 6\pi - 16. \end{aligned}$$

Method 2

$$A = 4 \times \left[ \frac{1}{2} \int_0^{\frac{\pi}{2}} (2(1 - \cos \theta))^2 d\theta \right]$$



## Note:

To apply Method 2, it is required to show that  $r = 2(1 + \cos \theta)$  and  $r = 2(1 - \cos \theta)$  are mirror images of each other.



## Note:

To apply Method 2, it is required to show that  $r = 2(1 + \cos \theta)$  and  $r = 2(1 - \cos \theta)$  are mirror images of each other.

**Sol.** We can write  $r = 1 - \cos \theta = 1 + \cos(\theta - \pi)$ .



## Note:

To apply Method 2, it is required to show that  $r = 2(1 + \cos \theta)$  and  $r = 2(1 - \cos \theta)$  are mirror images of each other.

**Sol.** We can write  $r = 1 - \cos \theta = 1 + \cos(\theta - \pi)$ .

Thus the curve  $r = 1 - \cos \theta$  is obtained from  $r = 1 + \cos \theta$  by replacing  $\theta$  by  $\theta + \pi$ . Therefore, to obtain the curve of  $r = 1 - \cos \theta$ , we just need to rotate the curve of  $r = 1 + \cos \theta$  by an angle  $\pi$ .




**Q:.** Find the area of the region inside the cardioid  $r = 1 + \cos \theta$  and outside the circle  $r = \cos \theta$ .



**Q:.** Find the area of the region inside the cardioid  $r = 1 + \cos \theta$  and outside the circle  $r = \cos \theta$ .

**Sol.**



$$A = 2 \times \frac{1}{2} \int_0^\pi [(1 + \cos \theta)^2 - (\cos \theta)^2] d\theta$$



$$A = 2 \times \frac{1}{2} \int_0^\pi [(1 + \cos \theta)^2 - (\cos \theta)^2] d\theta$$

Q: Is the above expression correct?





$$A = 2 \times \frac{1}{2} \int_0^\pi [(1 + \cos \theta)^2 - (\cos \theta)^2] d\theta$$

**Q:.** Is the above expression correct? No?



$$A = 2 \times \frac{1}{2} \int_0^\pi [(1 + \cos \theta)^2 - (\cos \theta)^2] d\theta$$

**Q:.** Is the above expression correct? No? Why?



$$A = 2 \times \frac{1}{2} \int_0^\pi [(1 + \cos \theta)^2 - (\cos \theta)^2] d\theta$$

**Q:.** Is the above expression correct? No? Why?

**Sol.** If we consider  $\theta$  from  $0$  to  $2\pi$ , then the circle  $r = \cos \theta$  would be traced twice.



# Method 1



## Method 1

Let  $A_1$  and  $A_2$  are the areas in first and second quadrant respectively. Then



## Method 1

Let  $A_1$  and  $A_2$  are the areas in first and second quadrant respectively. Then

$$\begin{aligned} A_1 &= \frac{1}{2} \int_0^{\pi/2} [(1 + \cos \theta)^2 - (\cos \theta)^2] d\theta \\ &= \frac{1}{2} \left( \frac{\pi}{2} + 2 \right). \end{aligned}$$

$$\begin{aligned} A_2 &= \frac{1}{2} \int_{\pi/2}^{\pi} (1 + \cos \theta)^2 d\theta \\ &= \frac{1}{2} \left( \frac{3\pi}{2} - 2 \right). \end{aligned}$$



Thus the required area

$$\begin{aligned} A &= 2(A_1 + A_2) \\ &= \left(\frac{\pi}{2} + 2\right) + \left(\frac{3\pi}{4} - 2\right) \\ &= \frac{5\pi}{4}. \end{aligned}$$



## Method 2





Method 2

$$A_1 = 2 \times \frac{1}{2} \int_0^\pi [(1 + \cos \theta)^2] d\theta = \frac{3\pi}{2}$$



Method 2

$$A_1 = 2 \times \frac{1}{2} \int_0^\pi [(1 + \cos \theta)^2] d\theta = \frac{3\pi}{2}$$

$$A_2 = \text{Area of circle } r = \cos \theta = \frac{\pi}{4}$$



Method 2

$$A_1 = 2 \times \frac{1}{2} \int_0^\pi [(1 + \cos \theta)^2] d\theta = \frac{3\pi}{2}$$

$$A_2 = \text{Area of circle } r = \cos \theta = \frac{\pi}{4}$$

$$\text{Thus the required area} = A_1 - A_2 = \frac{5\pi}{4}$$



# Length of a Polar Curve

Let  $r = f(\theta)$  has a continuous first derivative for  $\alpha \leq \theta \leq \beta$  and the point  $P(r, \theta)$  traces the curve exactly once as  $\theta$  varies from  $\alpha$  to  $\beta$ . Then the length of the curve from  $\theta = \alpha$  to  $\theta = \beta$  is given by

$$L = \int_{\alpha}^{\beta} \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta.$$



Q:.. Find the length of the spiral  $r = \theta^2$ ,  $0 \leq \theta \leq \sqrt{5}$ .



**Q:.** Find the length of the spiral  $r = \theta^2$ ,  $0 \leq \theta \leq \sqrt{5}$ .

**Sol.** The required length is

$$\begin{aligned} L &= \int_0^{\sqrt{5}} \sqrt{(\theta^2)^2 + (2\theta)^2} d\theta \\ &= \int_0^{\sqrt{5}} \sqrt{\theta^4 + 4\theta^2} d\theta \\ &= \int_0^{\sqrt{5}} |\theta| \sqrt{\theta^2 + 4} d\theta \\ &= \int_0^{\sqrt{5}} \theta \sqrt{\theta^2 + 4} d\theta \end{aligned}$$



Substitute  $u = \theta^2 + 4$

$$\begin{aligned} L &= \frac{1}{2} \int_4^9 \sqrt{u} \, du \\ &= \frac{19}{3}. \end{aligned}$$



Q:.. Find the length of cardioid

$$r = 1 + \cos \theta.$$





**Q:.** Find the length of cardioid

$$r = 1 + \cos \theta.$$

**Sol.**

$$\begin{aligned} L &= \int_0^{2\pi} \sqrt{(1 + \cos \theta)^2 + (-\sin \theta)^2} d\theta \\ &= 2 \int_0^{\pi} \sqrt{2 + 2 \cos \theta} d\theta \\ &= 4 \int_0^{\pi} \sqrt{(1 + \cos \theta)/2} d\theta \\ &= 8 \end{aligned}$$



Q: Find the length of the curve

$$r = a \sin^2 \frac{\theta}{2}, \quad 0 \leq \theta \leq \pi, \quad a > 0.$$



**Q:.** Find the length of the curve

$$r = a \sin^2 \frac{\theta}{2}, \quad 0 \leq \theta \leq \pi, \quad a > 0.$$

**Sol.** Note that the curve is a cardioid. The required length is

$$\begin{aligned} L &= \int_0^\pi \sqrt{\left(a \sin^2 \frac{\theta}{2}\right)^2 + \left(a \sin \frac{\theta}{2} \cos \frac{\theta}{2}\right)^2} d\theta \\ &= \int_0^\pi \left|a \sin \frac{\theta}{2}\right| d\theta \end{aligned}$$



$$\begin{aligned} L &= \int_0^{\pi} a \sin \frac{\theta}{2} d\theta \\ &= -2a \left( \cos \frac{\theta}{2} \right)_0^{\pi} \\ &= 2a. \end{aligned}$$



Q.: Find the length of the parabolic segment

$$r = \frac{6}{1 + \cos \theta}, \quad 0 \leq \theta \leq \frac{\pi}{2}.$$



**Q:.** Find the length of the parabolic segment

$$r = \frac{6}{1 + \cos \theta}, \quad 0 \leq \theta \leq \frac{\pi}{2}.$$

**Sol.** Better to convert  $r = \frac{6}{1 + \cos \theta} = 3 \sec^2 \frac{\theta}{2}$ . Thus the required length is



**Q:.** Find the length of the parabolic segment

$$r = \frac{6}{1 + \cos \theta}, \quad 0 \leq \theta \leq \frac{\pi}{2}.$$

**Sol.** Better to convert  $r = \frac{6}{1 + \cos \theta} = 3 \sec^2 \frac{\theta}{2}$ . Thus the required length is

$$\begin{aligned} L &= \int_0^{\frac{\pi}{2}} \sqrt{\left(3 \sec^2 \frac{\theta}{2}\right)^2 + \left(3 \sec^2 \frac{\theta}{2} \tan \frac{\theta}{2}\right)^2} d\theta \\ &= 3 \int_0^{\frac{\pi}{2}} \sec^2 \frac{\theta}{2} \sqrt{1 + \tan^2 \frac{\theta}{2}} d\theta \end{aligned}$$



Put  $\tan \frac{\theta}{2} = t$  so that  $\sec^2 \frac{\theta}{2} d\theta = 2dt$  and so

$$\begin{aligned} L &= 6 \int_0^1 \sqrt{1+t^2} dt \\ &= 6 \left[ \frac{t}{2} \sqrt{1+t^2} + \frac{1}{2} \ln(t + \sqrt{1+t^2}) \right]_0^1 \\ &= 3[ \sqrt{2} + \ln(1 + \sqrt{2}) ]. \end{aligned}$$

