

# Mathematics-II (MATH F112)

## Linear Algebra

Jitender Kumar

Department of Mathematics  
Birla Institute of Technology and Science Pilani  
Pilani-333031



# Chapter: 4 (Finite Dimensional vector space)

- 1 Introduction to Vector Spaces
- 2 Subspaces
- 3 Span
- 4 Linear Independence
- 5 Basis and Dimension
- 6 Constructing Special Basis



**Vector Space:** A nonempty set  $\mathcal{V}$  together with two operations **vector addition** (denoted as  $\oplus$ ) and **scalar multiplication** (denoted as  $\odot$ ) is said to be a **(real) vector space** if for every  $\mathbf{u}, \mathbf{v}, \mathbf{w}$  in  $\mathcal{V}$  and for every  $a, b \in \mathbb{R}$  the following properties hold:



**Vector Space:** A nonempty set  $\mathcal{V}$  together with two operations **vector addition** (denoted as  $\oplus$ ) and **scalar multiplication** (denoted as  $\odot$ ) is said to be a (real) vector space if for every  $\mathbf{u}, \mathbf{v}, \mathbf{w}$  in  $\mathcal{V}$  and for every  $a, b \in \mathbb{R}$  the following properties hold:

1  $\mathbf{u} \oplus \mathbf{v} \in \mathcal{V}$  (Closed under vector addition)



**Vector Space:** A nonempty set  $\mathcal{V}$  together with two operations **vector addition** (denoted as  $\oplus$ ) and **scalar multiplication** (denoted as  $\odot$ ) is said to be a **(real) vector space** if for every  $\mathbf{u}, \mathbf{v}, \mathbf{w}$  in  $\mathcal{V}$  and for every  $a, b \in \mathbb{R}$  the following properties hold:

1  $\mathbf{u} \oplus \mathbf{v} \in \mathcal{V}$  (Closed under vector addition)

2  $\mathbf{u} \oplus \mathbf{v} = \mathbf{v} \oplus \mathbf{u}$  (Commutativity)



**Vector Space:** A nonempty set  $\mathcal{V}$  together with two operations **vector addition** (denoted as  $\oplus$ ) and **scalar multiplication** (denoted as  $\odot$ ) is said to be a **(real) vector space** if for every  $\mathbf{u}, \mathbf{v}, \mathbf{w}$  in  $\mathcal{V}$  and for every  $a, b \in \mathbb{R}$  the following properties hold:

1  $\mathbf{u} \oplus \mathbf{v} \in \mathcal{V}$  (Closed under vector addition)

2  $\mathbf{u} \oplus \mathbf{v} = \mathbf{v} \oplus \mathbf{u}$  (Commutativity)

3  $(\mathbf{u} \oplus \mathbf{v}) \oplus \mathbf{w} = \mathbf{u} \oplus (\mathbf{v} \oplus \mathbf{w})$  (Associativity)



**Vector Space:** A nonempty set  $\mathcal{V}$  together with two operations **vector addition** (denoted as  $\oplus$ ) and **scalar multiplication** (denoted as  $\odot$ ) is said to be a (real) vector space if for every  $\mathbf{u}, \mathbf{v}, \mathbf{w}$  in  $\mathcal{V}$  and for every  $a, b \in \mathbb{R}$  the following properties hold:

- 1  $\mathbf{u} \oplus \mathbf{v} \in \mathcal{V}$  (Closed under vector addition)
- 2  $\mathbf{u} \oplus \mathbf{v} = \mathbf{v} \oplus \mathbf{u}$  (Commutativity)
- 3  $(\mathbf{u} \oplus \mathbf{v}) \oplus \mathbf{w} = \mathbf{u} \oplus (\mathbf{v} \oplus \mathbf{w})$  (Associativity)
- 4 There exists an element  $0 \in \mathcal{V}$ , called a **zero vector**, such that  $\mathbf{u} \oplus 0 = \mathbf{u}$  (Existence of additive identity)



- 5 For each  $\mathbf{u} \in \mathcal{V}$ , there is an element  $-\mathbf{u} \in \mathcal{V}$  such that  $\mathbf{u} \oplus (-\mathbf{u}) = 0$  (Existence of additive inverse)





- 5 For each  $\mathbf{u} \in \mathcal{V}$ , there is an element  $-\mathbf{u} \in \mathcal{V}$  such that  $\mathbf{u} \oplus (-\mathbf{u}) = 0$  (Existence of additive inverse)
- 6  $a \odot \mathbf{u} \in \mathcal{V}$  (Closed under scalar multiplication)



- 5 For each  $\mathbf{u} \in \mathcal{V}$ , there is an element  $-\mathbf{u} \in \mathcal{V}$  such that  $\mathbf{u} \oplus (-\mathbf{u}) = 0$  (Existence of additive inverse)
- 6  $a \odot \mathbf{u} \in \mathcal{V}$  (Closed under scalar multiplication)
- 7  $a \odot (\mathbf{u} \oplus \mathbf{v}) = (a \odot \mathbf{u}) \oplus (a \odot \mathbf{v})$  (Distributivity)



- 5 For each  $\mathbf{u} \in \mathcal{V}$ , there is an element  $-\mathbf{u} \in \mathcal{V}$  such that  $\mathbf{u} \oplus (-\mathbf{u}) = 0$  (Existence of additive inverse)
- 6  $a \odot \mathbf{u} \in \mathcal{V}$  (Closed under scalar multiplication)
- 7  $a \odot (\mathbf{u} \oplus \mathbf{v}) = (a \odot \mathbf{u}) \oplus (a \odot \mathbf{v})$  (Distributivity)
- 8  $(a + b) \odot \mathbf{u} = a \odot \mathbf{u} \oplus b \odot \mathbf{u}$  (Distributivity)



- 5 For each  $\mathbf{u} \in \mathcal{V}$ , there is an element  $-\mathbf{u} \in \mathcal{V}$  such that  $\mathbf{u} \oplus (-\mathbf{u}) = 0$  (Existence of additive inverse)
- 6  $a \odot \mathbf{u} \in \mathcal{V}$  (Closed under scalar multiplication)
- 7  $a \odot (\mathbf{u} \oplus \mathbf{v}) = (a \odot \mathbf{u}) \oplus (a \odot \mathbf{v})$  (Distributivity)
- 8  $(a + b) \odot \mathbf{u} = a \odot \mathbf{u} \oplus b \odot \mathbf{u}$  (Distributivity)
- 9  $(ab) \odot \mathbf{u} = a \odot (b \odot \mathbf{u})$



- 5 For each  $\mathbf{u} \in \mathcal{V}$ , there is an element  $-\mathbf{u} \in \mathcal{V}$  such that  $\mathbf{u} \oplus (-\mathbf{u}) = \mathbf{0}$  (Existence of additive inverse)
- 6  $a \odot \mathbf{u} \in \mathcal{V}$  (Closed under scalar multiplication)
- 7  $a \odot (\mathbf{u} \oplus \mathbf{v}) = (a \odot \mathbf{u}) \oplus (a \odot \mathbf{v})$  (Distributivity)
- 8  $(a + b) \odot \mathbf{u} = a \odot \mathbf{u} \oplus b \odot \mathbf{u}$  (Distributivity)
- 9  $(ab) \odot \mathbf{u} = a \odot (b \odot \mathbf{u})$
- 10  $1 \odot \mathbf{u} = \mathbf{u}$ .



**Note that** the set  $\mathcal{V} = \{0\}$  is a vector space with respect to



**Note that** the set  $\mathcal{V} = \{0\}$  is a vector space with respect to

- **vector addition**  $0 \oplus 0 = 0$
- **scalar multiplication**  $a \odot 0 = 0$  for all  $a \in \mathbb{R}$



**Note that** the set  $\mathcal{V} = \{0\}$  is a vector space with respect to

- **vector addition**  $0 \oplus 0 = 0$
- **scalar multiplication**  $a \odot 0 = 0$  for all  $a \in \mathbb{R}$

The vector space  $\mathcal{V} = \{0\}$  is called the **trivial vector space**.





**Example 1:** The set  $\mathbb{R}$  of real numbers is a **vector space** with respect to the following operations:

- $\mathbf{u} \oplus \mathbf{v} = \mathbf{u} + \mathbf{v}$  (**vector addition**)



**Example 1:** The set  $\mathbb{R}$  of real numbers is a **vector space** with respect to the following operations:

- $\mathbf{u} \oplus \mathbf{v} = \mathbf{u} + \mathbf{v}$  (**vector addition**)
- $a \odot \mathbf{u} = a\mathbf{u}$  (**scalar multiplication**)

for all  $a, \mathbf{u}, \mathbf{v} \in \mathbb{R}$ .



**Example 1:** The set  $\mathbb{R}$  of real numbers is a **vector space** with respect to the following operations:

- $\mathbf{u} \oplus \mathbf{v} = \mathbf{u} + \mathbf{v}$  (**vector addition**)
- $a \odot \mathbf{u} = a\mathbf{u}$  (**scalar multiplication**)

for all  $a, \mathbf{u}, \mathbf{v} \in \mathbb{R}$ .

**Question:** Does the set  $\mathbb{R}^+$  of positive real numbers form a vector space under the above defined vector addition and scalar multiplication?



**Example 2:** The set  $\mathbb{R}^+$  of a positive real numbers is a **vector space** with respect to the following operations:

•  $\mathbf{u} \oplus \mathbf{v} = \mathbf{u} \cdot \mathbf{v}$  (**vector addition**)



**Example 2:** The set  $\mathbb{R}^+$  of a positive real numbers is a **vector space** with respect to the following operations:

- $\mathbf{u} \oplus \mathbf{v} = \mathbf{u} \cdot \mathbf{v}$  (**vector addition**)
- $a \odot \mathbf{u} = \mathbf{u}^a$  (**scalar multiplication**)

for all  $a \in \mathbb{R}$  and  $\mathbf{u}, \mathbf{v} \in \mathbb{R}^+$ .



**Example 3:** The set  $\mathbb{R}^2 = \{[x_1, x_2] \mid x_1, x_2 \in \mathbb{R}\}$  is a **vector space** with respect to the following operations:

- $[x_1, x_2] \oplus [y_1, y_2] = [x_1 + y_1, x_2 + y_2]$  (**vector addition**)



**Example 3:** The set  $\mathbb{R}^2 = \{[x_1, x_2] \mid x_1, x_2 \in \mathbb{R}\}$  is a **vector space** with respect to the following operations:

- $[x_1, x_2] \oplus [y_1, y_2] = [x_1 + y_1, x_2 + y_2]$  (**vector addition**)
- $a \odot [x_1, x_2] = [ax_1, ax_2]$  (**scalar multiplication**)

for all  $a \in \mathbb{R}$  and  $[x_1, x_2], [y_1, y_2] \in \mathbb{R}^2$ .



**Example 3:** The set  $\mathbb{R}^2 = \{[x_1, x_2] \mid x_1, x_2 \in \mathbb{R}\}$  is a **vector space** with respect to the following operations:

- $[x_1, x_2] \oplus [y_1, y_2] = [x_1 + y_1, x_2 + y_2]$  (**vector addition**)
- $a \odot [x_1, x_2] = [ax_1, ax_2]$  (**scalar multiplication**)

for all  $a \in \mathbb{R}$  and  $[x_1, x_2], [y_1, y_2] \in \mathbb{R}^2$ .

**Question:** Does  $\mathbb{R}^2$  form a vector space under the above defined vector addition and





**Example 3:** The set  $\mathbb{R}^2 = \{[x_1, x_2] \mid x_1, x_2 \in \mathbb{R}\}$  is a **vector space** with respect to the following operations:

- $[x_1, x_2] \oplus [y_1, y_2] = [x_1 + y_1, x_2 + y_2]$  (**vector addition**)
- $a \odot [x_1, x_2] = [ax_1, ax_2]$  (**scalar multiplication**)

for all  $a \in \mathbb{R}$  and  $[x_1, x_2], [y_1, y_2] \in \mathbb{R}^2$ .

**Question:** Does  $\mathbb{R}^2$  form a vector space under the above defined vector addition and the following scalar multiplication

$$a \odot [x_1, x_2] = [0, ax_2]$$

for all  $a \in \mathbb{R}$  and  $[x_1, x_2] \in \mathbb{R}^2$ .



**Soln. of Example 3:** Let  $\mathbf{u} = [x_1, x_2]$ ,  $\mathbf{v} = [y_1, y_2]$  and  $\mathbf{w} = [z_1, z_2] \in \mathbb{R}^2$  and  $a, b \in \mathbb{R}$ .

1 Closure Property:

$$\mathbf{u} \oplus \mathbf{v}$$



**Soln. of Example 3:** Let  $\mathbf{u} = [x_1, x_2]$ ,  $\mathbf{v} = [y_1, y_2]$  and  $\mathbf{w} = [z_1, z_2] \in \mathbb{R}^2$  and  $a, b \in \mathbb{R}$ .

1 **Closure Property:**

$$\mathbf{u} \oplus \mathbf{v} = [x_1, x_2] \oplus [y_1, y_2]$$



**Soln. of Example 3:** Let  $\mathbf{u} = [x_1, x_2]$ ,  $\mathbf{v} = [y_1, y_2]$  and  $\mathbf{w} = [z_1, z_2] \in \mathbb{R}^2$  and  $a, b \in \mathbb{R}$ .

1 **Closure Property:**

$$\mathbf{u} \oplus \mathbf{v} = [x_1, x_2] \oplus [y_1, y_2] = [x_1 + y_1, x_2 + y_2]$$



**Soln. of Example 3:** Let  $\mathbf{u} = [x_1, x_2]$ ,  $\mathbf{v} = [y_1, y_2]$  and  $\mathbf{w} = [z_1, z_2] \in \mathbb{R}^2$  and  $a, b \in \mathbb{R}$ .

1 **Closure Property:**

$$\mathbf{u} \oplus \mathbf{v} = [x_1, x_2] \oplus [y_1, y_2] = [x_1 + y_1, x_2 + y_2] \in \mathbb{R}^2.$$



**Soln. of Example 3:** Let  $\mathbf{u} = [x_1, x_2]$ ,  $\mathbf{v} = [y_1, y_2]$  and  $\mathbf{w} = [z_1, z_2] \in \mathbb{R}^2$  and  $a, b \in \mathbb{R}$ .

1 Closure Property:

$$\mathbf{u} \oplus \mathbf{v} = [x_1, x_2] \oplus [y_1, y_2] = [x_1 + y_1, x_2 + y_2] \in \mathbb{R}^2.$$

2 Commutative Property:

$$\mathbf{u} \oplus \mathbf{v}$$



**Soln. of Example 3:** Let  $\mathbf{u} = [x_1, x_2]$ ,  $\mathbf{v} = [y_1, y_2]$  and  $\mathbf{w} = [z_1, z_2] \in \mathbb{R}^2$  and  $a, b \in \mathbb{R}$ .

1 **Closure Property:**

$$\mathbf{u} \oplus \mathbf{v} = [x_1, x_2] \oplus [y_1, y_2] = [x_1 + y_1, x_2 + y_2] \in \mathbb{R}^2.$$

2 **Commutative Property:**

$$\mathbf{u} \oplus \mathbf{v} = [x_1 + y_1, x_2 + y_2]$$



**Soln. of Example 3:** Let  $\mathbf{u} = [x_1, x_2]$ ,  $\mathbf{v} = [y_1, y_2]$  and  $\mathbf{w} = [z_1, z_2] \in \mathbb{R}^2$  and  $a, b \in \mathbb{R}$ .

1 Closure Property:

$$\mathbf{u} \oplus \mathbf{v} = [x_1, x_2] \oplus [y_1, y_2] = [x_1 + y_1, x_2 + y_2] \in \mathbb{R}^2.$$

2 Commutative Property:

$$\mathbf{u} \oplus \mathbf{v} = [x_1 + y_1, x_2 + y_2] = [y_1 + x_1, y_2 + x_2]$$





**Soln. of Example 3:** Let  $\mathbf{u} = [x_1, x_2]$ ,  $\mathbf{v} = [y_1, y_2]$  and  $\mathbf{w} = [z_1, z_2] \in \mathbb{R}^2$  and  $a, b \in \mathbb{R}$ .

1 Closure Property:

$$\mathbf{u} \oplus \mathbf{v} = [x_1, x_2] \oplus [y_1, y_2] = [x_1 + y_1, x_2 + y_2] \in \mathbb{R}^2.$$

2 Commutative Property:

$$\mathbf{u} \oplus \mathbf{v} = [x_1 + y_1, x_2 + y_2] = [y_1 + x_1, y_2 + x_2]$$

(commutativity of  $\mathbb{R}$  under addition)



**Soln. of Example 3:** Let  $\mathbf{u} = [x_1, x_2]$ ,  $\mathbf{v} = [y_1, y_2]$  and  $\mathbf{w} = [z_1, z_2] \in \mathbb{R}^2$  and  $a, b \in \mathbb{R}$ .

1 **Closure Property:**

$$\mathbf{u} \oplus \mathbf{v} = [x_1, x_2] \oplus [y_1, y_2] = [x_1 + y_1, x_2 + y_2] \in \mathbb{R}^2.$$

2 **Commutative Property:**

$$\begin{aligned}\mathbf{u} \oplus \mathbf{v} &= [x_1 + y_1, x_2 + y_2] = [y_1 + x_1, y_2 + x_2] \\ &\quad \text{(commutativity of } \mathbb{R} \text{ under addition)} \\ &= [y_1, y_2] \oplus [x_1, x_2] =\end{aligned}$$



**Soln. of Example 3:** Let  $\mathbf{u} = [x_1, x_2]$ ,  $\mathbf{v} = [y_1, y_2]$  and  $\mathbf{w} = [z_1, z_2] \in \mathbb{R}^2$  and  $a, b \in \mathbb{R}$ .

1 **Closure Property:**

$$\mathbf{u} \oplus \mathbf{v} = [x_1, x_2] \oplus [y_1, y_2] = [x_1 + y_1, x_2 + y_2] \in \mathbb{R}^2.$$

2 **Commutative Property:**

$$\begin{aligned}\mathbf{u} \oplus \mathbf{v} &= [x_1 + y_1, x_2 + y_2] = [y_1 + x_1, y_2 + x_2] \\ &\quad \text{(commutativity of } \mathbb{R} \text{ under addition)} \\ &= [y_1, y_2] \oplus [x_1, x_2] = \mathbf{v} \oplus \mathbf{u}\end{aligned}$$



**Soln. of Example 3:** Let  $\mathbf{u} = [x_1, x_2]$ ,  $\mathbf{v} = [y_1, y_2]$  and  $\mathbf{w} = [z_1, z_2] \in \mathbb{R}^2$  and  $a, b \in \mathbb{R}$ .

1 **Closure Property:**

$$\mathbf{u} \oplus \mathbf{v} = [x_1, x_2] \oplus [y_1, y_2] = [x_1 + y_1, x_2 + y_2] \in \mathbb{R}^2.$$

2 **Commutative Property:**

$$\begin{aligned}\mathbf{u} \oplus \mathbf{v} &= [x_1 + y_1, x_2 + y_2] = [y_1 + x_1, y_2 + x_2] \\ &\quad \text{(commutativity of } \mathbb{R} \text{ under addition)} \\ &= [y_1, y_2] \oplus [x_1, x_2] = \mathbf{v} \oplus \mathbf{u}\end{aligned}$$

3 **Associative Property:**

$$(\mathbf{u} \oplus \mathbf{v}) \oplus \mathbf{w}$$



**Soln. of Example 3:** Let  $\mathbf{u} = [x_1, x_2]$ ,  $\mathbf{v} = [y_1, y_2]$  and  $\mathbf{w} = [z_1, z_2] \in \mathbb{R}^2$  and  $a, b \in \mathbb{R}$ .

1 **Closure Property:**

$$\mathbf{u} \oplus \mathbf{v} = [x_1, x_2] \oplus [y_1, y_2] = [x_1 + y_1, x_2 + y_2] \in \mathbb{R}^2.$$

2 **Commutative Property:**

$$\begin{aligned}\mathbf{u} \oplus \mathbf{v} &= [x_1 + y_1, x_2 + y_2] = [y_1 + x_1, y_2 + x_2] \\ &\quad \text{(commutativity of } \mathbb{R} \text{ under addition)} \\ &= [y_1, y_2] \oplus [x_1, x_2] = \mathbf{v} \oplus \mathbf{u}\end{aligned}$$

3 **Associative Property:**

$$(\mathbf{u} \oplus \mathbf{v}) \oplus \mathbf{w} = [(x_1 + y_1) + z_1, (x_2 + y_2) + z_2]$$



**Soln. of Example 3:** Let  $\mathbf{u} = [x_1, x_2]$ ,  $\mathbf{v} = [y_1, y_2]$  and  $\mathbf{w} = [z_1, z_2] \in \mathbb{R}^2$  and  $a, b \in \mathbb{R}$ .

1 **Closure Property:**

$$\mathbf{u} \oplus \mathbf{v} = [x_1, x_2] \oplus [y_1, y_2] = [x_1 + y_1, x_2 + y_2] \in \mathbb{R}^2.$$

2 **Commutative Property:**

$$\begin{aligned}\mathbf{u} \oplus \mathbf{v} &= [x_1 + y_1, x_2 + y_2] = [y_1 + x_1, y_2 + x_2] \\ &\quad \text{(commutativity of } \mathbb{R} \text{ under addition)} \\ &= [y_1, y_2] \oplus [x_1, x_2] = \mathbf{v} \oplus \mathbf{u}\end{aligned}$$

3 **Associative Property:**

$$\begin{aligned}(\mathbf{u} \oplus \mathbf{v}) \oplus \mathbf{w} &= [(x_1 + y_1) + z_1, (x_2 + y_2) + z_2] \\ &= [x_1 + (y_1 + z_1), x_2 + (y_2 + z_2)]\end{aligned}$$



**Soln. of Example 3:** Let  $\mathbf{u} = [x_1, x_2]$ ,  $\mathbf{v} = [y_1, y_2]$  and  $\mathbf{w} = [z_1, z_2] \in \mathbb{R}^2$  and  $a, b \in \mathbb{R}$ .

1 **Closure Property:**

$$\mathbf{u} \oplus \mathbf{v} = [x_1, x_2] \oplus [y_1, y_2] = [x_1 + y_1, x_2 + y_2] \in \mathbb{R}^2.$$

2 **Commutative Property:**

$$\begin{aligned}\mathbf{u} \oplus \mathbf{v} &= [x_1 + y_1, x_2 + y_2] = [y_1 + x_1, y_2 + x_2] \\ &\quad \text{(commutativity of } \mathbb{R} \text{ under addition)} \\ &= [y_1, y_2] \oplus [x_1, x_2] = \mathbf{v} \oplus \mathbf{u}\end{aligned}$$

3 **Associative Property:**

$$\begin{aligned}(\mathbf{u} \oplus \mathbf{v}) \oplus \mathbf{w} &= [(x_1 + y_1) + z_1, (x_2 + y_2) + z_2] \\ &= [x_1 + (y_1 + z_1), x_2 + (y_2 + z_2)] \\ &\quad \text{(associativity of } \mathbb{R} \text{ under addition)}\end{aligned}$$



$$= [x_1, x_2] \oplus [y_1 + z_1, y_2 + z_2]$$





$$\begin{aligned}
 &= [x_1, x_2] \oplus [y_1 + z_1, y_2 + z_2] \\
 &= [x_1, x_2] \oplus ([y_1, y_2] \oplus [z_1, z_2])
 \end{aligned}$$



$$\begin{aligned}
 &= [x_1, x_2] \oplus [y_1 + z_1, y_2 + z_2] \\
 &= [x_1, x_2] \oplus ([y_1, y_2] \oplus [z_1, z_2]) \\
 &= \mathbf{u} \oplus (\mathbf{v} \oplus \mathbf{w})
 \end{aligned}$$



$$\begin{aligned}
 &= [x_1, x_2] \oplus [y_1 + z_1, y_2 + z_2] \\
 &= [x_1, x_2] \oplus ([y_1, y_2] \oplus [z_1, z_2]) \\
 &= \mathbf{u} \oplus (\mathbf{v} \oplus \mathbf{w})
 \end{aligned}$$

## 4 Existence of additive identity (zero vector):



$$\begin{aligned}
 &= [x_1, x_2] \oplus [y_1 + z_1, y_2 + z_2] \\
 &= [x_1, x_2] \oplus ([y_1, y_2] \oplus [z_1, z_2]) \\
 &= \mathbf{u} \oplus (\mathbf{v} \oplus \mathbf{w})
 \end{aligned}$$

- 4 **Existence of additive identity (zero vector):** For any  $\mathbf{u} = [x_1, x_2] \in \mathbb{R}^2$  there exists  $\mathbf{0} = [0, 0] \in \mathbb{R}^2$  such that



$$\begin{aligned}
 &= [x_1, x_2] \oplus [y_1 + z_1, y_2 + z_2] \\
 &= [x_1, x_2] \oplus ([y_1, y_2] \oplus [z_1, z_2]) \\
 &= \mathbf{u} \oplus (\mathbf{v} \oplus \mathbf{w})
 \end{aligned}$$

- 4 **Existence of additive identity (zero vector):** For any  $\mathbf{u} = [x_1, x_2] \in \mathbb{R}^2$  there exists  $\mathbf{0} = [0, 0] \in \mathbb{R}^2$  such that
- $$\mathbf{u} \oplus \mathbf{0} =$$



$$\begin{aligned}
 &= [x_1, x_2] \oplus [y_1 + z_1, y_2 + z_2] \\
 &= [x_1, x_2] \oplus ([y_1, y_2] \oplus [z_1, z_2]) \\
 &= \mathbf{u} \oplus (\mathbf{v} \oplus \mathbf{w})
 \end{aligned}$$

4 **Existence of additive identity (zero vector):** For any  $\mathbf{u} = [x_1, x_2] \in \mathbb{R}^2$  there exists  $\mathbf{0} = [0, 0] \in \mathbb{R}^2$  such that

$$\mathbf{u} \oplus \mathbf{0} = [x_1, x_2] \oplus [0, 0]$$



$$\begin{aligned}
 &= [x_1, x_2] \oplus [y_1 + z_1, y_2 + z_2] \\
 &= [x_1, x_2] \oplus ([y_1, y_2] \oplus [z_1, z_2]) \\
 &= \mathbf{u} \oplus (\mathbf{v} \oplus \mathbf{w})
 \end{aligned}$$

4 **Existence of additive identity (zero vector):** For any  $\mathbf{u} = [x_1, x_2] \in \mathbb{R}^2$  there exists  $\mathbf{0} = [0, 0] \in \mathbb{R}^2$  such that

$$\mathbf{u} \oplus \mathbf{0} = [x_1, x_2] \oplus [0, 0] = [x_1 + 0, x_2 + 0]$$



$$\begin{aligned}
 &= [x_1, x_2] \oplus [y_1 + z_1, y_2 + z_2] \\
 &= [x_1, x_2] \oplus ([y_1, y_2] \oplus [z_1, z_2]) \\
 &= \mathbf{u} \oplus (\mathbf{v} \oplus \mathbf{w})
 \end{aligned}$$

4 **Existence of additive identity (zero vector):** For any  $\mathbf{u} = [x_1, x_2] \in \mathbb{R}^2$  there exists  $\mathbf{0} = [0, 0] \in \mathbb{R}^2$  such that

$$\begin{aligned}
 \mathbf{u} \oplus \mathbf{0} &= [x_1, x_2] \oplus [0, 0] = [x_1 + 0, x_2 + 0] \\
 &= [x_1, x_2]
 \end{aligned}$$





$$\begin{aligned}
 &= [x_1, x_2] \oplus [y_1 + z_1, y_2 + z_2] \\
 &= [x_1, x_2] \oplus ([y_1, y_2] \oplus [z_1, z_2]) \\
 &= \mathbf{u} \oplus (\mathbf{v} \oplus \mathbf{w})
 \end{aligned}$$

4 **Existence of additive identity (zero vector):** For any  $\mathbf{u} = [x_1, x_2] \in \mathbb{R}^2$  there exists  $\mathbf{0} = [0, 0] \in \mathbb{R}^2$  such that

$$\begin{aligned}
 \mathbf{u} \oplus \mathbf{0} &= [x_1, x_2] \oplus [0, 0] = [x_1 + 0, x_2 + 0] \\
 &= [x_1, x_2] \\
 &= \mathbf{u}
 \end{aligned}$$



$$\begin{aligned}
 &= [x_1, x_2] \oplus [y_1 + z_1, y_2 + z_2] \\
 &= [x_1, x_2] \oplus ([y_1, y_2] \oplus [z_1, z_2]) \\
 &= \mathbf{u} \oplus (\mathbf{v} \oplus \mathbf{w})
 \end{aligned}$$

- 4 **Existence of additive identity (zero vector):** For any  $\mathbf{u} = [x_1, x_2] \in \mathbb{R}^2$  there exists  $\mathbf{0} = [0, 0] \in \mathbb{R}^2$  such that

$$\begin{aligned}
 \mathbf{u} \oplus \mathbf{0} &= [x_1, x_2] \oplus [0, 0] = [x_1 + 0, x_2 + 0] \\
 &= [x_1, x_2] \\
 &= \mathbf{u}
 \end{aligned}$$

- 5 **Existence of additive inverse:**



$$\begin{aligned}
 &= [x_1, x_2] \oplus [y_1 + z_1, y_2 + z_2] \\
 &= [x_1, x_2] \oplus ([y_1, y_2] \oplus [z_1, z_2]) \\
 &= \mathbf{u} \oplus (\mathbf{v} \oplus \mathbf{w})
 \end{aligned}$$

- 4 **Existence of additive identity (zero vector):** For any  $\mathbf{u} = [x_1, x_2] \in \mathbb{R}^2$  there exists  $\mathbf{0} = [0, 0] \in \mathbb{R}^2$  such that

$$\begin{aligned}
 \mathbf{u} \oplus \mathbf{0} &= [x_1, x_2] \oplus [0, 0] = [x_1 + 0, x_2 + 0] \\
 &= [x_1, x_2] \\
 &= \mathbf{u}
 \end{aligned}$$

- 5 **Existence of additive inverse:** For each  $\mathbf{u} = [x_1, x_2] \in \mathbb{R}^2$  there exists  $-\mathbf{u} = [-x_1, -x_2]$  in  $\mathbb{R}^2$  such that



$$\begin{aligned}
 &= [x_1, x_2] \oplus [y_1 + z_1, y_2 + z_2] \\
 &= [x_1, x_2] \oplus ([y_1, y_2] \oplus [z_1, z_2]) \\
 &= \mathbf{u} \oplus (\mathbf{v} \oplus \mathbf{w})
 \end{aligned}$$

- 4 **Existence of additive identity (zero vector):** For any  $\mathbf{u} = [x_1, x_2] \in \mathbb{R}^2$  there exists  $\mathbf{0} = [0, 0] \in \mathbb{R}^2$  such that

$$\begin{aligned}
 \mathbf{u} \oplus \mathbf{0} &= [x_1, x_2] \oplus [0, 0] = [x_1 + 0, x_2 + 0] \\
 &= [x_1, x_2] \\
 &= \mathbf{u}
 \end{aligned}$$

- 5 **Existence of additive inverse:** For each  $\mathbf{u} = [x_1, x_2] \in \mathbb{R}^2$  there exists  $-\mathbf{u} = [-x_1, -x_2]$  in  $\mathbb{R}^2$  such that

$$\mathbf{u} \oplus (-\mathbf{u}) = [x_1, x_2] \oplus [-x_1, -x_2]$$



$$\begin{aligned}
 &= [x_1, x_2] \oplus [y_1 + z_1, y_2 + z_2] \\
 &= [x_1, x_2] \oplus ([y_1, y_2] \oplus [z_1, z_2]) \\
 &= \mathbf{u} \oplus (\mathbf{v} \oplus \mathbf{w})
 \end{aligned}$$

- 4 **Existence of additive identity (zero vector):** For any  $\mathbf{u} = [x_1, x_2] \in \mathbb{R}^2$  there exists  $\mathbf{0} = [0, 0] \in \mathbb{R}^2$  such that

$$\begin{aligned}
 \mathbf{u} \oplus \mathbf{0} &= [x_1, x_2] \oplus [0, 0] = [x_1 + 0, x_2 + 0] \\
 &= [x_1, x_2] \\
 &= \mathbf{u}
 \end{aligned}$$

- 5 **Existence of additive inverse:** For each  $\mathbf{u} = [x_1, x_2] \in \mathbb{R}^2$  there exists  $-\mathbf{u} = [-x_1, -x_2]$  in  $\mathbb{R}^2$  such that

$$\begin{aligned}
 \mathbf{u} \oplus (-\mathbf{u}) &= [x_1, x_2] \oplus [-x_1, -x_2] \\
 &= [x_1 + (-x_1), x_2 + (-x_2)]
 \end{aligned}$$



$$\begin{aligned}
 &= [x_1, x_2] \oplus [y_1 + z_1, y_2 + z_2] \\
 &= [x_1, x_2] \oplus ([y_1, y_2] \oplus [z_1, z_2]) \\
 &= \mathbf{u} \oplus (\mathbf{v} \oplus \mathbf{w})
 \end{aligned}$$

- 4 **Existence of additive identity (zero vector):** For any  $\mathbf{u} = [x_1, x_2] \in \mathbb{R}^2$  there exists  $\mathbf{0} = [0, 0] \in \mathbb{R}^2$  such that

$$\begin{aligned}
 \mathbf{u} \oplus \mathbf{0} &= [x_1, x_2] \oplus [0, 0] = [x_1 + 0, x_2 + 0] \\
 &= [x_1, x_2] \\
 &= \mathbf{u}
 \end{aligned}$$

- 5 **Existence of additive inverse:** For each  $\mathbf{u} = [x_1, x_2] \in \mathbb{R}^2$  there exists  $-\mathbf{u} = [-x_1, -x_2]$  in  $\mathbb{R}^2$  such that

$$\begin{aligned}
 \mathbf{u} \oplus (-\mathbf{u}) &= [x_1, x_2] \oplus [-x_1, -x_2] \\
 &= [x_1 + (-x_1), x_2 + (-x_2)] = [0, 0]
 \end{aligned}$$



$$\begin{aligned}
 &= [x_1, x_2] \oplus [y_1 + z_1, y_2 + z_2] \\
 &= [x_1, x_2] \oplus ([y_1, y_2] \oplus [z_1, z_2]) \\
 &= \mathbf{u} \oplus (\mathbf{v} \oplus \mathbf{w})
 \end{aligned}$$

- 4 **Existence of additive identity (zero vector):** For any  $\mathbf{u} = [x_1, x_2] \in \mathbb{R}^2$  there exists  $\mathbf{0} = [0, 0] \in \mathbb{R}^2$  such that

$$\begin{aligned}
 \mathbf{u} \oplus \mathbf{0} &= [x_1, x_2] \oplus [0, 0] = [x_1 + 0, x_2 + 0] \\
 &= [x_1, x_2] \\
 &= \mathbf{u}
 \end{aligned}$$

- 5 **Existence of additive inverse:** For each  $\mathbf{u} = [x_1, x_2] \in \mathbb{R}^2$  there exists  $-\mathbf{u} = [-x_1, -x_2]$  in  $\mathbb{R}^2$  such that

$$\begin{aligned}
 \mathbf{u} \oplus (-\mathbf{u}) &= [x_1, x_2] \oplus [-x_1, -x_2] \\
 &= [x_1 + (-x_1), x_2 + (-x_2)] = [0, 0] = \mathbf{0}
 \end{aligned}$$



6

## Closure Property of scalar multiplication:





6

## Closure Property of scalar multiplication:

$$a \odot \mathbf{u}$$



6

## Closure Property of scalar multiplication:

$$a \odot \mathbf{u} = a \odot [x_1, x_2]$$



6

## Closure Property of scalar multiplication:

$$a \odot \mathbf{u} = a \odot [x_1, x_2] = [ax_1, ax_2]$$



6 Closure Property of scalar multiplication:

$$a \odot \mathbf{u} = a \odot [x_1, x_2] = [ax_1, ax_2] \in \mathbb{R}^2.$$



6 Closure Property of scalar multiplication:

$a \odot \mathbf{u} = a \odot [x_1, x_2] = [ax_1, ax_2] \in \mathbb{R}^2$ . Thus,  $\mathbb{R}^2$  is closed under scalar multiplication.



- 6 Closure Property of scalar multiplication:  
 $a \odot \mathbf{u} = a \odot [x_1, x_2] = [ax_1, ax_2] \in \mathbb{R}^2$ . Thus,  $\mathbb{R}^2$  is closed under scalar multiplication.
- 7 Distributivity over vector addition:



- 6 Closure Property of scalar multiplication:  
 $a \odot \mathbf{u} = a \odot [x_1, x_2] = [ax_1, ax_2] \in \mathbb{R}^2$ . Thus,  $\mathbb{R}^2$  is closed under scalar multiplication.
- 7 Distributivity over vector addition:  
 $a \odot (\mathbf{u} \oplus \mathbf{v})$



- 6 Closure Property of scalar multiplication:  
 $a \odot \mathbf{u} = a \odot [x_1, x_2] = [ax_1, ax_2] \in \mathbb{R}^2$ . Thus,  $\mathbb{R}^2$  is closed under scalar multiplication.
- 7 Distributivity over vector addition:  
 $a \odot (\mathbf{u} \oplus \mathbf{v}) = a \odot ([x_1, x_2] \oplus [y_1, y_2])$





- 6 Closure Property of scalar multiplication:  
 $a \odot \mathbf{u} = a \odot [x_1, x_2] = [ax_1, ax_2] \in \mathbb{R}^2$ . Thus,  $\mathbb{R}^2$  is closed under scalar multiplication.
- 7 Distributivity over vector addition:  
$$\begin{aligned} a \odot (\mathbf{u} \oplus \mathbf{v}) &= a \odot ([x_1, x_2] \oplus [y_1, y_2]) \\ &= a \odot [x_1 + y_1, x_2 + y_2] \end{aligned}$$



- 6 Closure Property of scalar multiplication:  
 $a \odot \mathbf{u} = a \odot [x_1, x_2] = [ax_1, ax_2] \in \mathbb{R}^2$ . Thus,  $\mathbb{R}^2$  is closed under scalar multiplication.
- 7 Distributivity over vector addition:  
$$\begin{aligned} a \odot (\mathbf{u} \oplus \mathbf{v}) &= a \odot ([x_1, x_2] \oplus [y_1, y_2]) \\ &= a \odot [x_1 + y_1, x_2 + y_2] \\ &= [a(x_1 + y_1), a(x_2 + y_2)] \end{aligned}$$



6 Closure Property of scalar multiplication:  
 $a \odot \mathbf{u} = a \odot [x_1, x_2] = [ax_1, ax_2] \in \mathbb{R}^2$ . Thus,  $\mathbb{R}^2$  is closed under scalar multiplication.

7 Distributivity over vector addition:  
$$\begin{aligned} a \odot (\mathbf{u} \oplus \mathbf{v}) &= a \odot ([x_1, x_2] \oplus [y_1, y_2]) \\ &= a \odot [x_1 + y_1, x_2 + y_2] \\ &= [a(x_1 + y_1), a(x_2 + y_2)] \\ &= [ax_1 + ay_1, ax_2 + ay_2] \text{ (distributivity in } \mathbb{R}) \end{aligned}$$



6 Closure Property of scalar multiplication:  
 $a \odot \mathbf{u} = a \odot [x_1, x_2] = [ax_1, ax_2] \in \mathbb{R}^2$ . Thus,  $\mathbb{R}^2$  is closed under scalar multiplication.

7 Distributivity over vector addition:  
$$\begin{aligned} a \odot (\mathbf{u} \oplus \mathbf{v}) &= a \odot ([x_1, x_2] \oplus [y_1, y_2]) \\ &= a \odot [x_1 + y_1, x_2 + y_2] \\ &= [a(x_1 + y_1), a(x_2 + y_2)] \\ &= [ax_1 + ay_1, ax_2 + ay_2] \text{ (distributivity in } \mathbb{R}) \\ &= [ax_1, ax_2] \oplus [ay_1, ay_2] \end{aligned}$$



6 Closure Property of scalar multiplication:  
 $a \odot \mathbf{u} = a \odot [x_1, x_2] = [ax_1, ax_2] \in \mathbb{R}^2$ . Thus,  $\mathbb{R}^2$  is closed under scalar multiplication.

7 Distributivity over vector addition:  
$$\begin{aligned} a \odot (\mathbf{u} \oplus \mathbf{v}) &= a \odot ([x_1, x_2] \oplus [y_1, y_2]) \\ &= a \odot [x_1 + y_1, x_2 + y_2] \\ &= [a(x_1 + y_1), a(x_2 + y_2)] \\ &= [ax_1 + ay_1, ax_2 + ay_2] \text{ (distributivity in } \mathbb{R}) \\ &= [ax_1, ax_2] \oplus [ay_1, ay_2] \\ &= (a \odot [x_1, x_2]) \oplus (a \odot [y_1, y_2]) \end{aligned}$$



6 Closure Property of scalar multiplication:  
 $a \odot \mathbf{u} = a \odot [x_1, x_2] = [ax_1, ax_2] \in \mathbb{R}^2$ . Thus,  $\mathbb{R}^2$  is closed under scalar multiplication.

7 Distributivity over vector addition:  
$$\begin{aligned} a \odot (\mathbf{u} \oplus \mathbf{v}) &= a \odot ([x_1, x_2] \oplus [y_1, y_2]) \\ &= a \odot [x_1 + y_1, x_2 + y_2] \\ &= [a(x_1 + y_1), a(x_2 + y_2)] \\ &= [ax_1 + ay_1, ax_2 + ay_2] \text{ (distributivity in } \mathbb{R}) \\ &= [ax_1, ax_2] \oplus [ay_1, ay_2] \\ &= (a \odot [x_1, x_2]) \oplus (a \odot [y_1, y_2]) \\ &= (a \odot \mathbf{u}) \oplus (a \odot \mathbf{v}) \end{aligned}$$



## 8 Distributivity over scalar addition:



## 8 Distributivity over scalar addition:

$$(a + b) \odot \mathbf{u} = (a + b) \odot [x_1, x_2]$$





8

## Distributivity over scalar addition:

$$\begin{aligned}(a + b) \odot \mathbf{u} &= (a + b) \odot [x_1, x_2] \\ &= [(a + b)x_1, (a + b)x_2]\end{aligned}$$



8

## Distributivity over scalar addition:

$$\begin{aligned}(a + b) \odot \mathbf{u} &= (a + b) \odot [x_1, x_2] \\ &= [(a + b)x_1, (a + b)x_2] \\ &= [ax_1 + bx_1, ax_2 + bx_2] \text{ (distributivity in } \mathbb{R})\end{aligned}$$



8

## Distributivity over scalar addition:

$$\begin{aligned}(a + b) \odot \mathbf{u} &= (a + b) \odot [x_1, x_2] \\ &= [(a + b)x_1, (a + b)x_2] \\ &= [ax_1 + bx_1, ax_2 + bx_2] \text{ (distributivity in } \mathbb{R}) \\ &= [ax_1, ax_2] \oplus [bx_1, bx_2]\end{aligned}$$



8

## Distributivity over scalar addition:

$$\begin{aligned}(a + b) \odot \mathbf{u} &= (a + b) \odot [x_1, x_2] \\&= [(a + b)x_1, (a + b)x_2] \\&= [ax_1 + bx_1, ax_2 + bx_2] \text{ (distributivity in } \mathbb{R}) \\&= [ax_1, ax_2] \oplus [bx_1, bx_2] \\&= (a \odot [x_1, x_2]) \oplus (b \odot [x_1, x_2])\end{aligned}$$



## 8 Distributivity over scalar addition:

$$\begin{aligned}(a + b) \odot \mathbf{u} &= (a + b) \odot [x_1, x_2] \\&= [(a + b)x_1, (a + b)x_2] \\&= [ax_1 + bx_1, ax_2 + bx_2] \text{ (distributivity in } \mathbb{R}) \\&= [ax_1, ax_2] \oplus [bx_1, bx_2] \\&= (a \odot [x_1, x_2]) \oplus (b \odot [x_1, x_2]) \\&= (a \odot \mathbf{u}) \oplus (b \odot \mathbf{u})\end{aligned}$$



## 8 Distributivity over scalar addition:

$$\begin{aligned}(a + b) \odot \mathbf{u} &= (a + b) \odot [x_1, x_2] \\&= [(a + b)x_1, (a + b)x_2] \\&= [ax_1 + bx_1, ax_2 + bx_2] \text{ (distributivity in } \mathbb{R}) \\&= [ax_1, ax_2] \oplus [bx_1, bx_2] \\&= (a \odot [x_1, x_2]) \oplus (b \odot [x_1, x_2]) \\&= (a \odot \mathbf{u}) \oplus (b \odot \mathbf{u})\end{aligned}$$

## 9 $(ab) \odot \mathbf{u}$



8 Distributivity over scalar addition:

$$\begin{aligned}(a + b) \odot \mathbf{u} &= (a + b) \odot [x_1, x_2] \\&= [(a + b)x_1, (a + b)x_2] \\&= [ax_1 + bx_1, ax_2 + bx_2] \text{ (distributivity in } \mathbb{R}) \\&= [ax_1, ax_2] \oplus [bx_1, bx_2] \\&= (a \odot [x_1, x_2]) \oplus (b \odot [x_1, x_2]) \\&= (a \odot \mathbf{u}) \oplus (b \odot \mathbf{u})\end{aligned}$$

9  $(ab) \odot \mathbf{u} = (ab) \odot [x_1, x_2]$



## 8 Distributivity over scalar addition:

$$\begin{aligned}(a + b) \odot \mathbf{u} &= (a + b) \odot [x_1, x_2] \\&= [(a + b)x_1, (a + b)x_2] \\&= [ax_1 + bx_1, ax_2 + bx_2] \text{ (distributivity in } \mathbb{R}) \\&= [ax_1, ax_2] \oplus [bx_1, bx_2] \\&= (a \odot [x_1, x_2]) \oplus (b \odot [x_1, x_2]) \\&= (a \odot \mathbf{u}) \oplus (b \odot \mathbf{u})\end{aligned}$$

## 9

$$\begin{aligned}(ab) \odot \mathbf{u} &= (ab) \odot [x_1, x_2] \\&= [(ab)x_1, (ab)x_2]\end{aligned}$$





## 8 Distributivity over scalar addition:

$$\begin{aligned}(a + b) \odot \mathbf{u} &= (a + b) \odot [x_1, x_2] \\&= [(a + b)x_1, (a + b)x_2] \\&= [ax_1 + bx_1, ax_2 + bx_2] \text{ (distributivity in } \mathbb{R}) \\&= [ax_1, ax_2] \oplus [bx_1, bx_2] \\&= (a \odot [x_1, x_2]) \oplus (b \odot [x_1, x_2]) \\&= (a \odot \mathbf{u}) \oplus (b \odot \mathbf{u})\end{aligned}$$

## 9

$$\begin{aligned}(ab) \odot \mathbf{u} &= (ab) \odot [x_1, x_2] \\&= [(ab)x_1, (ab)x_2] \\&= [a(bx_1), a(bx_2)]\end{aligned}$$



8 Distributivity over scalar addition:

$$\begin{aligned}(a + b) \odot \mathbf{u} &= (a + b) \odot [x_1, x_2] \\&= [(a + b)x_1, (a + b)x_2] \\&= [ax_1 + bx_1, ax_2 + bx_2] \text{ (distributivity in } \mathbb{R}) \\&= [ax_1, ax_2] \oplus [bx_1, bx_2] \\&= (a \odot [x_1, x_2]) \oplus (b \odot [x_1, x_2]) \\&= (a \odot \mathbf{u}) \oplus (b \odot \mathbf{u})\end{aligned}$$

9

$$\begin{aligned}(ab) \odot \mathbf{u} &= (ab) \odot [x_1, x_2] \\&= [(ab)x_1, (ab)x_2] \\&= [a(bx_1), a(bx_2)] \\&\text{(associativity of } \mathbb{R} \text{ under multiplication)}\end{aligned}$$



## 8 Distributivity over scalar addition:

$$\begin{aligned}(a + b) \odot \mathbf{u} &= (a + b) \odot [x_1, x_2] \\&= [(a + b)x_1, (a + b)x_2] \\&= [ax_1 + bx_1, ax_2 + bx_2] \text{ (distributivity in } \mathbb{R}) \\&= [ax_1, ax_2] \oplus [bx_1, bx_2] \\&= (a \odot [x_1, x_2]) \oplus (b \odot [x_1, x_2]) \\&= (a \odot \mathbf{u}) \oplus (b \odot \mathbf{u})\end{aligned}$$

## 9

$$\begin{aligned}(ab) \odot \mathbf{u} &= (ab) \odot [x_1, x_2] \\&= [(ab)x_1, (ab)x_2] \\&= [a(bx_1), a(bx_2)] \\&\text{(associativity of } \mathbb{R} \text{ under multiplication)} \\&= a \odot [bx_1, bx_2]\end{aligned}$$



8 Distributivity over scalar addition:

$$\begin{aligned}(a + b) \odot \mathbf{u} &= (a + b) \odot [x_1, x_2] \\&= [(a + b)x_1, (a + b)x_2] \\&= [ax_1 + bx_1, ax_2 + bx_2] \text{ (distributivity in } \mathbb{R}) \\&= [ax_1, ax_2] \oplus [bx_1, bx_2] \\&= (a \odot [x_1, x_2]) \oplus (b \odot [x_1, x_2]) \\&= (a \odot \mathbf{u}) \oplus (b \odot \mathbf{u})\end{aligned}$$

9

$$\begin{aligned}(ab) \odot \mathbf{u} &= (ab) \odot [x_1, x_2] \\&= [(ab)x_1, (ab)x_2] \\&= [a(bx_1), a(bx_2)] \\&\text{ (associativity of } \mathbb{R} \text{ under multiplication)} \\&= a \odot [bx_1, bx_2] \\&= a \odot (b \odot [x_1, x_2])\end{aligned}$$



## 8 Distributivity over scalar addition:

$$\begin{aligned}(a + b) \odot \mathbf{u} &= (a + b) \odot [x_1, x_2] \\&= [(a + b)x_1, (a + b)x_2] \\&= [ax_1 + bx_1, ax_2 + bx_2] \text{ (distributivity in } \mathbb{R}) \\&= [ax_1, ax_2] \oplus [bx_1, bx_2] \\&= (a \odot [x_1, x_2]) \oplus (b \odot [x_1, x_2]) \\&= (a \odot \mathbf{u}) \oplus (b \odot \mathbf{u})\end{aligned}$$

## 9

$$\begin{aligned}(ab) \odot \mathbf{u} &= (ab) \odot [x_1, x_2] \\&= [(ab)x_1, (ab)x_2] \\&= [a(bx_1), a(bx_2)] \\&\text{ (associativity of } \mathbb{R} \text{ under multiplication)} \\&= a \odot [bx_1, bx_2] \\&= a \odot (b \odot [x_1, x_2]) \\&= a \odot (b \odot \mathbf{u})\end{aligned}$$



$$10 \quad 1 \odot \mathbf{u} =$$



$$10 \quad 1 \odot \mathbf{u} = 1 \odot [x_1, x_2] =$$



$$10 \quad 1 \odot \mathbf{u} = 1 \odot [x_1, x_2] = [1x_1, 1x_2] =$$





$$10 \quad 1 \odot \mathbf{u} = 1 \odot [x_1, x_2] = [1x_1, 1x_2] = [x_1, x_2] =$$



$$10 \quad 1 \odot \mathbf{u} = 1 \odot [x_1, x_2] = [1x_1, 1x_2] = [x_1, x_2] = \mathbf{u}$$



10  $1 \odot \mathbf{u} = 1 \odot [x_1, x_2] = [1x_1, 1x_2] = [x_1, x_2] = \mathbf{u}.$

Thus  $\mathbb{R}^2$  is vector space under usual vector addition and scalar multiplication.



**Example 4:** The set  $\mathbb{R}^n = \{[x_1, x_2, \dots, x_n] \mid x_i \in \mathbb{R}\}$  is a **vector space** with respect to the following operations:

- $[x_1, x_2, \dots, x_n] \oplus [y_1, y_2, \dots, y_n]$   
 $= [x_1 + y_1, x_2 + y_2, \dots, x_n + y_n]$  (**vector addition**)



**Example 4:** The set  $\mathbb{R}^n = \{[x_1, x_2, \dots, x_n] \mid x_i \in \mathbb{R}\}$  is a **vector space** with respect to the following operations:

- $[x_1, x_2, \dots, x_n] \oplus [y_1, y_2, \dots, y_n]$   
 $= [x_1 + y_1, x_2 + y_2, \dots, x_n + y_n]$  (**vector addition**)
- $a \odot [x_1, x_2, \dots, x_n] = [ax_1, ax_2, \dots, ax_n]$  (**scalar multiplication**)

for all  $a \in \mathbb{R}$  and  $[x_1, x_2, \dots, x_n], [y_1, y_2, \dots, y_n] \in \mathbb{R}^n$ .



## Example 5: The set

$$\mathcal{M}_{mn} = \{[a_{ij}]_{m \times n} \mid a_{ij} \in \mathbb{R}\}$$

of all  $m \times n$  matrices with real entries is a **vector space** with respect to the following operations:

- $[a_{ij}]_{m \times n} \oplus [b_{ij}]_{m \times n} = [a_{ij} + b_{ij}]_{m \times n}$  (**vector addition**)
- $a \odot [a_{ij}]_{m \times n} = [aa_{ij}]_{m \times n}$  (**scalar multiplication**)

for all  $a \in \mathbb{R}$  and  $[a_{ij}]_{m \times n}, [b_{ij}]_{m \times n} \in \mathcal{M}_{mn}$ .



## Example 6: Let

$$\Phi = \{f \mid f : \mathbb{R} \rightarrow \mathbb{R}\}$$

be the set of real-valued functions defined on  $\mathbb{R}$ .  
Define

$$f \oplus g = f + g \text{ (vector addition),}$$

and  $a \odot f = af \text{ (scalar multiplication),}$



## Example 6: Let

$$\Phi = \{f \mid f : \mathbb{R} \rightarrow \mathbb{R}\}$$

be the set of real-valued functions defined on  $\mathbb{R}$ .  
Define

$$f \oplus g = f + g \quad (\text{vector addition}),$$

where  $(f + g)(x) = f(x) + g(x) \quad \forall x \in \mathbb{R}$ .

$$\text{and} \quad a \odot f = af \quad (\text{scalar multiplication}),$$





## Example 6: Let

$$\Phi = \{f \mid f : \mathbb{R} \rightarrow \mathbb{R}\}$$

be the set of real-valued functions defined on  $\mathbb{R}$ .  
Define

$$f \oplus g = f + g \quad (\text{vector addition}),$$

where  $(f + g)(x) = f(x) + g(x) \quad \forall x \in \mathbb{R}$ .

$$\text{and} \quad a \odot f = af \quad (\text{scalar multiplication}),$$

where  $(af)(x) = af(x) \quad \forall x \in \mathbb{R}$ .



## Example 6: Let

$$\Phi = \{f \mid f : \mathbb{R} \rightarrow \mathbb{R}\}$$

be the set of real-valued functions defined on  $\mathbb{R}$ .  
Define

$$f \oplus g = f + g \quad (\text{vector addition}),$$

where  $(f + g)(x) = f(x) + g(x) \quad \forall x \in \mathbb{R}$ .

$$\text{and} \quad a \odot f = af \quad (\text{scalar multiplication}),$$

where  $(af)(x) = af(x) \quad \forall x \in \mathbb{R}$ .

Then  $\Phi$  is a **vector space** with respect to above defined vector addition and scalar multiplication.



## Example 7: Let

$$\mathcal{P}_2 = \{a_2x^2 + a_1x + a_0 \mid a_2, a_1, a_0 \in \mathbb{R}\}$$

be the set of all polynomials of degree  $\leq 2$  with real coefficients. Define addition and scalar multiplication in usual way i.e. if

$$p(x) = a_2x^2 + a_1x + a_0 \text{ and } q(x) = b_2x^2 + b_1x + b_0$$

are in  $\mathcal{P}_2$ , then

$$p(x) \oplus q(x) = (a_2 + b_2)x^2 + (a_1 + b_1)x + (a_0 + b_0)$$

$$c \odot p(x) = ca_2x^2 + ca_1x + ca_0.$$

Show that  $\mathcal{P}_2$  is a **vector space**.



**In general**, for any fixed natural number  $n$ , the set

$$\mathcal{P}_n = \{a_n x^n + \cdots + a_1 x + a_0 \mid a_n, \dots, a_1, a_0 \in \mathbb{R}\}$$

of all polynomials of degree less than or equal to  $n$  is a **vector space**. under the usual addition (term by term) and scalar multiplication of polynomials.



**In general**, for any fixed natural number  $n$ , the set

$$\mathcal{P}_n = \{a_n x^n + \cdots + a_1 x + a_0 \mid a_n, \dots, a_1, a_0 \in \mathbb{R}\}$$

of all polynomials of degree less than or equal to  $n$  is a **vector space**. under the usual addition (term by term) and scalar multiplication of polynomials.

**Question:** Does the set of all polynomials of degree 7 form a vector space under the usual operation of addition and scalar multiplication?



**Example 8:** The set  $\mathcal{P}$  of all polynomials with real coefficients is a **vector space** under the usual operation of polynomial (term by term) addition and scalar multiplication.



**Theorem:** Let  $\mathcal{V}$  be a vector space. Then for every  $\mathbf{v} \in \mathcal{V}$  and  $\alpha \in \mathbb{R}$ , we have

- $\alpha \mathbf{0} = \mathbf{0}$
- $0\mathbf{v} = \mathbf{0}$
- $(-1)\mathbf{v} = -\mathbf{v}$
- If  $\alpha\mathbf{v} = \mathbf{0}$ , then  $\alpha = 0$  or  $\mathbf{v} = \mathbf{0}$ .



# Section 4.2 (Subspaces)





# Section 4.2 (Subspaces)

**Subspace:** A nonempty subset  $W$  of a vector space  $\mathcal{V}$  is said to be a **subspace** of  $\mathcal{V}$  if  $W$  is itself a vector space with respect to the same operations (vector addition and scalar multiplication) of  $\mathcal{V}$ .



## Section 4.2 (Subspaces)

**Subspace:** A nonempty subset  $W$  of a vector space  $\mathcal{V}$  is said to be a **subspace** of  $\mathcal{V}$  if  $W$  is itself a vector space with respect to the same operations (vector addition and scalar multiplication) of  $\mathcal{V}$ .

**Note that** every vector space  $\mathcal{V}$  has at least two subspaces:  $\{0\}$  and  $\mathcal{V}$  itself.



## Section 4.2 (Subspaces)

**Subspace:** A nonempty subset  $W$  of a vector space  $\mathcal{V}$  is said to be a **subspace** of  $\mathcal{V}$  if  $W$  is itself a vector space with respect to the same operations (vector addition and scalar multiplication) of  $\mathcal{V}$ .

**Note that** every vector space  $\mathcal{V}$  has at least two subspaces:  $\{0\}$  and  $\mathcal{V}$  itself. The subspace  $\{0\}$  is known as **trivial subspace**.



**Example:** The set

$$W = \{[x, y] \in \mathbb{R}^2 \mid y = 0\}$$

forms a vector space with respect to usual vector addition and scalar multiplication in  $\mathbb{R}^2$ .



**Example:** The set

$$W = \{[x, y] \in \mathbb{R}^2 \mid y = 0\}$$

forms a vector space with respect to usual vector addition and scalar multiplication in  $\mathbb{R}^2$ . Thus,  $W$  is a subspace of  $\mathbb{R}^2$ .



**Example:** The set

$$W = \{[x, y] \in \mathbb{R}^2 \mid y = 0\}$$

forms a vector space with respect to usual vector addition and scalar multiplication in  $\mathbb{R}^2$ . Thus,  $W$  is a subspace of  $\mathbb{R}^2$ .

**Question:** Does the set

$$W = \{[x, y] \in \mathbb{R}^2 \mid x \neq y\}$$

form a subspace of  $\mathbb{R}^2$ ?



**Theorem:** A **nonempty** subset  $W$  of a vector space  $\mathcal{V}$  is a subspace of  $\mathcal{V}$  if and only if the following conditions hold:



**Theorem:** A **nonempty** subset  $W$  of a vector space  $\mathcal{V}$  is a subspace of  $\mathcal{V}$  if and only if the following conditions hold:

- $\mathbf{u} + \mathbf{v} \in W$  for all  $\mathbf{u}, \mathbf{v} \in W$ .
- $a\mathbf{u} \in W$  for all  $a \in \mathbb{R}, \mathbf{u} \in W$ .





**Theorem:** A **nonempty** subset  $W$  of a vector space  $\mathcal{V}$  is a subspace of  $\mathcal{V}$  if and only if the following conditions hold:

- $\mathbf{u} + \mathbf{v} \in W$  for all  $\mathbf{u}, \mathbf{v} \in W$ .
- $a\mathbf{u} \in W$  for all  $a \in \mathbb{R}, \mathbf{u} \in W$ .

**Remark:** If  $W$  is a subspace of a vector space  $\mathcal{V}$ , then  $0 \in W$ .



**Exercise:** Examine whether the following sets are subspaces of the vector space  $\mathbb{R}^3$ .

- $W_1 = \{[x, y, z] \in \mathbb{R}^3 \mid x \geq 0\}.$



**Exercise:** Examine whether the following sets are subspaces of the vector space  $\mathbb{R}^3$ .

- $W_1 = \{[x, y, z] \in \mathbb{R}^3 \mid x \geq 0\}.$
- $W_2 = \{[x, y, z] \in \mathbb{R}^3 \mid x + y + z = 0\}.$



**Exercise:** Examine whether the following sets are subspaces of the vector space  $\mathbb{R}^3$ .

- $W_1 = \{[x, y, z] \in \mathbb{R}^3 \mid x \geq 0\}.$
- $W_2 = \{[x, y, z] \in \mathbb{R}^3 \mid x + y + z = 0\}.$
- $W_3 = \{[x, y, z] \in \mathbb{R}^3 \mid x = y^2\}.$



**Exercise:** Examine whether the following sets are subspaces of the vector space  $\mathbb{R}^3$ .

- $W_1 = \{[x, y, z] \in \mathbb{R}^3 \mid x \geq 0\}.$
- $W_2 = \{[x, y, z] \in \mathbb{R}^3 \mid x + y + z = 0\}.$
- $W_3 = \{[x, y, z] \in \mathbb{R}^3 \mid x = y^2\}.$
- $W_4 = \{[x, y, z] \in \mathbb{R}^3 \mid x + y + z = 2\}.$



**Exercise:** Examine whether the following sets are subspaces of the vector space  $\mathbb{R}^3$ .

- $W_1 = \{[x, y, z] \in \mathbb{R}^3 \mid x \geq 0\}$ .
- $W_2 = \{[x, y, z] \in \mathbb{R}^3 \mid x + y + z = 0\}$ .
- $W_3 = \{[x, y, z] \in \mathbb{R}^3 \mid x = y^2\}$ .
- $W_4 = \{[x, y, z] \in \mathbb{R}^3 \mid x + y + z = 2\}$ .
- $W_5 = \{[x, y, z] \in \mathbb{R}^3 \mid x^2 + y^2 + z^2 = 1\}$ .



**Exercise:** Examine whether the following sets are subspaces of the vector space  $\mathcal{M}_{22}$  (see Example 5).

- $W_1 = \{A \in \mathcal{M}_{22} \mid A \text{ is singular}\}.$



**Exercise:** Examine whether the following sets are subspaces of the vector space  $\mathcal{M}_{22}$  (see Example 5).

- $W_1 = \{A \in \mathcal{M}_{22} \mid A \text{ is singular}\}.$
- $W_2 = \{A \in \mathcal{M}_{22} \mid A \text{ is nonsingular}\}.$





**Exercise:** Examine whether the following sets are subspaces of the vector space  $\mathcal{M}_{22}$  (see Example 5).

- $W_1 = \{A \in \mathcal{M}_{22} \mid A \text{ is singular}\}.$
- $W_2 = \{A \in \mathcal{M}_{22} \mid A \text{ is nonsingular}\}.$
- $W_3 = \{A \in \mathcal{M}_{22} \mid A \text{ is in RREF}\}.$



**Exercise:** Examine whether the following sets are subspaces of the vector space  $\mathcal{M}_{22}$  (see Example 5).

- $W_1 = \{A \in \mathcal{M}_{22} \mid A \text{ is singular}\}.$
- $W_2 = \{A \in \mathcal{M}_{22} \mid A \text{ is nonsingular}\}.$
- $W_3 = \{A \in \mathcal{M}_{22} \mid A \text{ is in RREF}\}.$
- $W_4 = \{A \in \mathcal{M}_{22} \mid A \text{ is symmetric}\}.$



**Exercise:** Examine whether the following sets are subspaces of the vector space  $\mathcal{M}_{22}$  (see Example 5).

- $W_1 = \{A \in \mathcal{M}_{22} \mid A \text{ is singular}\}.$
- $W_2 = \{A \in \mathcal{M}_{22} \mid A \text{ is nonsingular}\}.$
- $W_3 = \{A \in \mathcal{M}_{22} \mid A \text{ is in RREF}\}.$
- $W_4 = \{A \in \mathcal{M}_{22} \mid A \text{ is symmetric}\}.$
- $W_5 = \{A \in \mathcal{M}_{22} \mid A^2 = A\}.$



**Exercise:** Examine whether the following sets are subspaces of the vector space  $\Phi$  (see Example 6).



**Exercise:** Examine whether the following sets are subspaces of the vector space  $\Phi$  (see Example 6).

- $W_1 = \{f \in \Phi \mid f(-x) = f(x) \text{ for all } x \in \mathbb{R}\}.$



**Exercise:** Examine whether the following sets are subspaces of the vector space  $\Phi$  (see Example 6).

- $W_1 = \{f \in \Phi \mid f(-x) = f(x) \text{ for all } x \in \mathbb{R}\}.$
- $W_2 = \{f \in \Phi \mid f(-x) = -f(x) \text{ for all } x \in \mathbb{R}\}.$



**Exercise:** Examine whether the following sets are subspaces of the vector space  $\Phi$  (see Example 6).

- $W_1 = \{f \in \Phi \mid f(-x) = f(x) \text{ for all } x \in \mathbb{R}\}.$
- $W_2 = \{f \in \Phi \mid f(-x) = -f(x) \text{ for all } x \in \mathbb{R}\}.$
- $W_3 = \{f \in \Phi \mid f(1) = 0\}.$



**Exercise:** Examine whether the following sets are subspaces of the vector space  $\Phi$  (see Example 6).

- $W_1 = \{f \in \Phi \mid f(-x) = f(x) \text{ for all } x \in \mathbb{R}\}.$
- $W_2 = \{f \in \Phi \mid f(-x) = -f(x) \text{ for all } x \in \mathbb{R}\}.$
- $W_3 = \{f \in \Phi \mid f(1) = 0\}.$
- $W_4 = \{f \in \Phi \mid f\left(\frac{1}{2}\right) = f(1)\}.$





**Exercise:** Examine whether the following sets are subspaces of the vector space  $\Phi$  (see Example 6).

- $W_1 = \{f \in \Phi \mid f(-x) = f(x) \text{ for all } x \in \mathbb{R}\}.$
- $W_2 = \{f \in \Phi \mid f(-x) = -f(x) \text{ for all } x \in \mathbb{R}\}.$
- $W_3 = \{f \in \Phi \mid f(1) = 0\}.$
- $W_4 = \{f \in \Phi \mid f\left(\frac{1}{2}\right) = f(1)\}.$
- $W_5 = \{f \in \Phi \mid f(1) = \frac{1}{2}\}.$



**Result:** Let  $W_1$  and  $W_2$  be two subspaces of vector space  $\mathcal{V}$ . Then



**Result:** Let  $W_1$  and  $W_2$  be two subspaces of vector space  $\mathcal{V}$ . Then

- their intersection i.e.  $W_1 \cap W_2$  is a subspace of  $\mathcal{V}$ .



**Result:** Let  $W_1$  and  $W_2$  be two subspaces of vector space  $\mathcal{V}$ . Then

- their intersection i.e.  $W_1 \cap W_2$  is a subspace of  $\mathcal{V}$ .

- their sum, defined as

$$W_1 + W_2 = \{w_1 + w_2 \mid w_1 \in W_1, w_2 \in W_2\},$$

is a subspace of  $\mathcal{V}$ .



**Result:** Let  $W_1$  and  $W_2$  be two subspaces of vector space  $\mathcal{V}$ . Then

- their intersection i.e.  $W_1 \cap W_2$  is a subspace of  $\mathcal{V}$ .
- their union  $W_1 \cup W_2$  **need not** be a subspace of  $\mathcal{V}$ .
- their sum, defined as

$$W_1 + W_2 = \{w_1 + w_2 \mid w_1 \in W_1, w_2 \in W_2\},$$

is a subspace of  $\mathcal{V}$ .



**Result:** Let  $W_1$  and  $W_2$  be two subspaces of vector space  $\mathcal{V}$ . Then

- their intersection i.e.  $W_1 \cap W_2$  is a subspace of  $\mathcal{V}$ .
- their union  $W_1 \cup W_2$  **need not** be a subspace of  $\mathcal{V}$ .
- $W_1 \cup W_2$  is subspace of  $\mathcal{V}$  if and only if either  $W_1 \subset W_2$  or  $W_2 \subset W_1$ .
- their sum, defined as

$$W_1 + W_2 = \{w_1 + w_2 \mid w_1 \in W_1, w_2 \in W_2\},$$

is a subspace of  $\mathcal{V}$ .



## Section 4.3 (Span)

**Question:** Given a subset  $S$  of a vector space  $\mathcal{V}$ , how to construct a subspace containing  $S$ ?



**Linear combination:** Let  $\mathcal{V}$  be a vector space and  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k \in \mathcal{V}$ . Then a vector  $\mathbf{v} \in \mathcal{V}$  is a linear combination of  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$  if





**Linear combination:** Let  $\mathcal{V}$  be a vector space and  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k \in \mathcal{V}$ . Then a vector  $\mathbf{v} \in \mathcal{V}$  is a linear combination of  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$  if

$$\mathbf{v} = a_1 \mathbf{v}_1 + a_2 \mathbf{v}_2 + \dots + a_k \mathbf{v}_k; \quad a_i (1 \leq i \leq k) \in \mathbb{R}$$



**Linear combination:** Let  $\mathcal{V}$  be a vector space and  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k \in \mathcal{V}$ . Then a vector  $\mathbf{v} \in \mathcal{V}$  is a linear combination of  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$  if

$$\mathbf{v} = a_1 \mathbf{v}_1 + a_2 \mathbf{v}_2 + \dots + a_k \mathbf{v}_k; \quad a_i (1 \leq i \leq k) \in \mathbb{R}$$

**Example:** The vector  $[3, 4]$  is a linear combination of  $[1, 0]$  and  $[0, 1]$  in  $\mathbb{R}^2$ .



**Linear combination:** Let  $\mathcal{V}$  be a vector space and  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k \in \mathcal{V}$ . Then a vector  $\mathbf{v} \in \mathcal{V}$  is a linear combination of  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$  if

$$\mathbf{v} = a_1 \mathbf{v}_1 + a_2 \mathbf{v}_2 + \cdots + a_k \mathbf{v}_k; \quad a_i (1 \leq i \leq k) \in \mathbb{R}$$

**Example:** The vector  $[3, 4]$  is a linear combination of  $[1, 0]$  and  $[0, 1]$  in  $\mathbb{R}^2$ .

Note that

$$[3, 4] = 2[1, 1] + [1, 2].$$



**Linear combination:** Let  $\mathcal{V}$  be a vector space and  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k \in \mathcal{V}$ . Then a vector  $\mathbf{v} \in \mathcal{V}$  is a linear combination of  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$  if

$$\mathbf{v} = a_1\mathbf{v}_1 + a_2\mathbf{v}_2 + \cdots + a_k\mathbf{v}_k; \quad a_i (1 \leq i \leq k) \in \mathbb{R}$$

**Example:** The vector  $[3, 4]$  is a linear combination of  $[1, 0]$  and  $[0, 1]$  in  $\mathbb{R}^2$ .

Note that

$$[3, 4] = 2[1, 1] + [1, 2].$$

Thus,  $[3, 4]$  is a linear combination of  $[1, 1]$  and  $[1, 2]$  also.



**Span of a set:** Let  $S$  be a nonempty subset of a vector space  $\mathcal{V}$ . Then the **span** of  $S$  is the set of all possible (finite) linear combinations of the vectors in  $S$  and it is denoted by  **$\text{span}(S)$**



**Span of a set:** Let  $S$  be a nonempty subset of a vector space  $\mathcal{V}$ . Then the **span** of  $S$  is the set of all possible (finite) linear combinations of the vectors in  $S$  and it is denoted by  **$\text{span}(S)$**  i.e.

$$\text{span}(S) = \{a_1 \mathbf{v}_1 + \cdots + a_k \mathbf{v}_k \mid \mathbf{v}_i \in S, a_i \in \mathbb{R}, 1 \leq i \leq k\}$$



**Span of a set:** Let  $S$  be a nonempty subset of a vector space  $\mathcal{V}$ . Then the **span** of  $S$  is the set of all possible (finite) linear combinations of the vectors in  $S$  and it is denoted by  **$\text{span}(S)$**  i.e.

$$\text{span}(S) = \{a_1 \mathbf{v}_1 + \cdots + a_k \mathbf{v}_k \mid \mathbf{v}_i \in S, a_i \in \mathbb{R}, 1 \leq i \leq k\}$$

- For a subset  $S = \{[1, 0], [0, 1]\}$  of  $\mathbb{R}^2$ , we have  $\text{span}(S)$



**Span of a set:** Let  $S$  be a nonempty subset of a vector space  $\mathcal{V}$ . Then the **span** of  $S$  is the set of all possible (finite) linear combinations of the vectors in  $S$  and it is denoted by  $\text{span}(S)$  i.e.

$$\text{span}(S) = \{a_1 \mathbf{v}_1 + \cdots + a_k \mathbf{v}_k \mid \mathbf{v}_i \in S, a_i \in \mathbb{R}, 1 \leq i \leq k\}$$

- For a subset  $S = \{[1, 0], [0, 1]\}$  of  $\mathbb{R}^2$ , we have  $\text{span}(S) = \mathbb{R}^2$ .





**Span of a set:** Let  $S$  be a nonempty subset of a vector space  $\mathcal{V}$ . Then the **span** of  $S$  is the set of all possible (finite) linear combinations of the vectors in  $S$  and it is denoted by **span**( $S$ ) i.e.

$$\text{span}(S) = \{a_1 \mathbf{v}_1 + \cdots + a_k \mathbf{v}_k \mid \mathbf{v}_i \in S, a_i \in \mathbb{R}, 1 \leq i \leq k\}$$

- For a subset  $S = \{[1, 0], [0, 1]\}$  of  $\mathbb{R}^2$ , we have  $\text{span}(S) = \mathbb{R}^2$ .
- For a subset  $S = \{[1, 0, 0], [0, 1, 0], [0, 0, 1]\}$  of  $\mathbb{R}^3$  we have  $\text{span}(S)$



**Span of a set:** Let  $S$  be a nonempty subset of a vector space  $\mathcal{V}$ . Then the **span** of  $S$  is the set of all possible (finite) linear combinations of the vectors in  $S$  and it is denoted by **span**( $S$ ) i.e.

$$\text{span}(S) = \{a_1 \mathbf{v}_1 + \cdots + a_k \mathbf{v}_k \mid \mathbf{v}_i \in S, a_i \in \mathbb{R}, 1 \leq i \leq k\}$$

- For a subset  $S = \{[1, 0], [0, 1]\}$  of  $\mathbb{R}^2$ , we have  $\text{span}(S) = \mathbb{R}^2$ .
- For a subset  $S = \{[1, 0, 0], [0, 1, 0], [0, 0, 1]\}$  of  $\mathbb{R}^3$  we have  $\text{span}(S) = \mathbb{R}^3$ .



**Exercise:** Let  $\mathcal{V} = \mathbb{R}^3$  and  $S = \{[1, 0, 0], [0, 1, 0]\}$ .

- Find  $\text{span}(S)$ .
- Do  $[3, 2, 0]$  and  $[2, 5, 1]$  belong to  $\text{span}(S)$ ?



**Exercise:** Let  $\mathcal{V} = \mathbb{R}^3$  and  $S = \{[1, 0, 0], [0, 1, 0]\}$ .

- Find  $\text{span}(S)$ .
- Do  $[3, 2, 0]$  and  $[2, 5, 1]$  belong to  $\text{span}(S)$ ?

**Solution:**

$$\text{span}(S) = \{a[1, 0, 0] + b[0, 1, 0] \mid a, b \in \mathbb{R}\}$$



**Exercise:** Let  $\mathcal{V} = \mathbb{R}^3$  and  $S = \{[1, 0, 0], [0, 1, 0]\}$ .

- Find  $\text{span}(S)$ .
- Do  $[3, 2, 0]$  and  $[2, 5, 1]$  belong to  $\text{span}(S)$ ?

**Solution:**

$$\begin{aligned}\text{span}(S) &= \{a[1, 0, 0] + b[0, 1, 0] \mid a, b \in \mathbb{R}\} \\ &= \{[a, b, 0] \mid a, b \in \mathbb{R}\}\end{aligned}$$



**Exercise:** Let  $\mathcal{V} = \mathbb{R}^3$  and  $S = \{[1, 0, 0], [0, 1, 0]\}$ .

- Find  $\text{span}(S)$ .
- Do  $[3, 2, 0]$  and  $[2, 5, 1]$  belong to  $\text{span}(S)$ ?

**Solution:**

$$\begin{aligned}\text{span}(S) &= \{a[1, 0, 0] + b[0, 1, 0] \mid a, b \in \mathbb{R}\} \\ &= \{[a, b, 0] \mid a, b \in \mathbb{R}\}\end{aligned}$$

Clearly,  $[3, 2, 0] \in \text{span}(S)$  but  $[2, 5, 1] \notin \text{span}(S)$ .



**Exercise:** Let  $\mathcal{V} = \mathbb{R}^3$  and  $S = \{[1, 0, 0], [0, 1, 0]\}$ .

- Find  $\text{span}(S)$ .
- Do  $[3, 2, 0]$  and  $[2, 5, 1]$  belong to  $\text{span}(S)$ ?

**Solution:**

$$\begin{aligned}\text{span}(S) &= \{a[1, 0, 0] + b[0, 1, 0] \mid a, b \in \mathbb{R}\} \\ &= \{[a, b, 0] \mid a, b \in \mathbb{R}\}\end{aligned}$$

Clearly,  $[3, 2, 0] \in \text{span}(S)$  but  $[2, 5, 1] \notin \text{span}(S)$ .

In this exercise **note that**  $\text{span}(S)$  is a subspace of  $\mathbb{R}^3$ .



**Exercise:** Let  $\mathbf{v}_1, \mathbf{v}_2$  be in a vector space  $\mathcal{V}$ . Then show that  $W = \text{span}\{\mathbf{v}_1, \mathbf{v}_2\}$  is a subspace of  $\mathcal{V}$ .





**Exercise:** Let  $\mathbf{v}_1, \mathbf{v}_2$  be in a vector space  $\mathcal{V}$ . Then show that  $W = \text{span}\{\mathbf{v}_1, \mathbf{v}_2\}$  is a subspace of  $\mathcal{V}$ .

**Theorem:** Let  $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$  be a nonempty subset of a vector space  $\mathcal{V}$ . Then

- $\text{span}(S)$  is a subspace of  $\mathcal{V}$ .
- $\text{span}(S)$  is the smallest subspace of  $\mathcal{V}$  containing  $S$ .



**Exercise:** Let  $\mathbf{v}_1, \mathbf{v}_2$  be in a vector space  $\mathcal{V}$ . Then show that  $W = \text{span}\{\mathbf{v}_1, \mathbf{v}_2\}$  is a subspace of  $\mathcal{V}$ .

**Theorem:** Let  $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$  be a nonempty subset of a vector space  $\mathcal{V}$ . Then

- $\text{span}(S)$  is a subspace of  $\mathcal{V}$ .
- $\text{span}(S)$  is the smallest subspace of  $\mathcal{V}$  containing  $S$ .

**Convention:**  $\text{span}(\emptyset) = \{0\}$ .



**Row space of a matrix:** Let  $A$  be an  $m \times n$  matrix. The **row space of  $A$** , denoted by  $\text{row}(A)$ , is the subspace of  $\mathbb{R}^n$  spanned by the rows of  $A$ .



**Row space of a matrix:** Let  $A$  be an  $m \times n$  matrix. The **row space of  $A$** , denoted by  $\text{row}(A)$ , is the subspace of  $\mathbb{R}^n$  spanned by the rows of  $A$ .

**Theorem:** Let  $B$  be any matrix that is row equivalent to a matrix  $A$ . Then  $\text{row}(B) = \text{row}(A)$ .



**Row space of a matrix:** Let  $A$  be an  $m \times n$  matrix. The **row space of  $A$** , denoted by  $\text{row}(A)$ , is the subspace of  $\mathbb{R}^n$  spanned by the rows of  $A$ .

**Theorem:** Let  $B$  be any matrix that is row equivalent to a matrix  $A$ . Then  $\text{row}(B) = \text{row}(A)$ .

**Corollary:** For any matrix  $A$ , we have

$$\text{row}(A) = \text{row}(\text{RREF}(A)).$$



**Exercise:** Let  $\mathcal{V} = \mathbb{R}^3$  and

$$S = \{[1, 3, -1], [2, 7, -3], [4, 8, -7]\}.$$

Then find  $\text{span}(S)$  in simplified form.



**Exercise:** Let  $\mathcal{V} = \mathbb{R}^3$  and

$$S = \{[1, 3, -1], [2, 7, -3], [4, 8, -7]\}.$$

Then find  $\text{span}(S)$  in simplified form.

**Solution:** To determine  $\text{span}(S)$  in simplified form consider

$$A = \begin{bmatrix} 1 & 3 & -1 \\ 2 & 7 & -3 \\ 4 & 8 & -7 \end{bmatrix}$$



**Exercise:** Let  $\mathcal{V} = \mathbb{R}^3$  and

$$S = \{[1, 3, -1], [2, 7, -3], [4, 8, -7]\}.$$

Then find  $\text{span}(S)$  in simplified form.

**Solution:** To determine  $\text{span}(S)$  in simplified form consider

$$A = \begin{bmatrix} 1 & 3 & -1 \\ 2 & 7 & -3 \\ 4 & 8 & -7 \end{bmatrix}$$

Note that  $\text{span}(S) = \text{row}(A) =$





**Exercise:** Let  $\mathcal{V} = \mathbb{R}^3$  and

$$S = \{[1, 3, -1], [2, 7, -3], [4, 8, -7]\}.$$

Then find  $\text{span}(S)$  in simplified form.

**Solution:** To determine  $\text{span}(S)$  in simplified form consider

$$A = \begin{bmatrix} 1 & 3 & -1 \\ 2 & 7 & -3 \\ 4 & 8 & -7 \end{bmatrix}$$

Note that  $\text{span}(S) = \text{row}(A) = \text{row}(\text{RREF}(A))$ .



Note that

$$\text{RREF}(A) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$



Note that

$$\text{RREF}(A) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$\text{row}(\text{RREF}(A))$

$$= \{a[1, 0, 0] + b[0, 1, 0] + c[0, 0, 1] \mid a, b, c \in \mathbb{R}\}$$



Note that

$$\text{RREF}(A) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\text{row}(\text{RREF}(A))$$

$$= \{a[1, 0, 0] + b[0, 1, 0] + c[0, 0, 1] \mid a, b, c \in \mathbb{R}\}$$

$$\text{row}(\text{RREF}(A)) = \text{span}(S) = \{[a, b, c] \mid a, b, c \in \mathbb{R}\}$$

$$\text{span}(S) = \mathbb{R}^3$$



**Simplified Span Method:** Let  $S$  be a finite subset of  $\mathbb{R}^n$  containing  $k$  vectors, with  $k \geq 2$ .

**Step 1:** Construct a matrix  $A$  of order  $k \times n$  by using the vectors in  $S$  as **the rows of  $A$** .



**Simplified Span Method:** Let  $S$  be a finite subset of  $\mathbb{R}^n$  containing  $k$  vectors, with  $k \geq 2$ .

**Step 1:** Construct a matrix  $A$  of order  $k \times n$  by using the vectors in  $S$  as **the rows of**  $A$ . Then  $\text{span}(S) = \text{row}(A)$ .



**Simplified Span Method:** Let  $S$  be a finite subset of  $\mathbb{R}^n$  containing  $k$  vectors, with  $k \geq 2$ .

**Step 1:** Construct a matrix  $A$  of order  $k \times n$  by using the vectors in  $S$  as **the rows of**  $A$ . Then  $\text{span}(S) = \text{row}(A)$ .

**Step 2:** Find  $\text{RREF}(A)$ .



**Simplified Span Method:** Let  $S$  be a finite subset of  $\mathbb{R}^n$  containing  $k$  vectors, with  $k \geq 2$ .

**Step 1:** Construct a matrix  $A$  of order  $k \times n$  by using the vectors in  $S$  as **the rows of**  $A$ . Then  $\text{span}(S) = \text{row}(A)$ .

**Step 2:** Find  $\text{RREF}(A)$ .

**Step 3:** Then, the set of all linear combinations of the **nonzero rows** of  $\text{RREF}(A)$  gives a simplified form for  $\text{span}(S)$ .





**Exercise:** For a given vector space  $\mathcal{V}$  and a subset  $S$  of  $\mathcal{V}$ , find a simplified general form of  $\text{span}(S)$  using Simplified Span Method:



**Exercise:** For a given vector space  $\mathcal{V}$  and a subset  $S$  of  $\mathcal{V}$ , find a simplified general form of  $\text{span}(S)$  using Simplified Span Method:

1  $\mathcal{V} = \mathbb{R}^3, S = \{[1, 1, 1], [2, 1, 1], [1, 1, 2]\}.$



**Exercise:** For a given vector space  $\mathcal{V}$  and a subset  $S$  of  $\mathcal{V}$ , find a simplified general form of  $\text{span}(S)$  using Simplified Span Method:

1  $\mathcal{V} = \mathbb{R}^3, S = \{[1, 1, 1], [2, 1, 1], [1, 1, 2]\}.$

2  $\mathcal{V} = \mathcal{P}_2, S = \{x^2 + x + 1, x + 1, 1\}.$



**Exercise:** For a given vector space  $\mathcal{V}$  and a subset  $S$  of  $\mathcal{V}$ , find a simplified general form of  $\text{span}(S)$  using Simplified Span Method:

1  $\mathcal{V} = \mathbb{R}^3, S = \{[1, 1, 1], [2, 1, 1], [1, 1, 2]\}.$

2  $\mathcal{V} = \mathcal{P}_2, S = \{x^2 + x + 1, x + 1, 1\}.$

3  $\mathcal{V} = \mathcal{P}_2, S = \{x^2 + 4x - 3, 2x^2 + x + 5, 7x - 11\}.$



**Exercise:** For a given vector space  $\mathcal{V}$  and a subset  $S$  of  $\mathcal{V}$ , find a simplified general form of  $\text{span}(S)$  using Simplified Span Method:

1  $\mathcal{V} = \mathbb{R}^3, S = \{[1, 1, 1], [2, 1, 1], [1, 1, 2]\}.$

2  $\mathcal{V} = \mathcal{P}_2, S = \{x^2 + x + 1, x + 1, 1\}.$

3  $\mathcal{V} = \mathcal{P}_2, S = \{x^2 + 4x - 3, 2x^2 + x + 5, 7x - 11\}.$

4  $\mathcal{V} = \mathcal{M}_{22}, S = \left\{ \begin{bmatrix} 1 & 3 \\ -2 & 1 \end{bmatrix}, \begin{bmatrix} -2 & -5 \\ 3 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ -3 & 4 \end{bmatrix} \right\}.$



# Section 4.4 (Linear Independence)



# Section 4.4 (Linear Independence)

**Definition:** A subset  $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$  of a vector space  $\mathcal{V}$  is said to be **linearly dependent** (LD) if there exist real numbers  $a_1, a_2, \dots, a_n$  not all zero such that

$$a_1\mathbf{v}_1 + a_2\mathbf{v}_2 + \dots + a_n\mathbf{v}_n = \mathbf{0}.$$



# Section 4.4 (Linear Independence)

**Definition:** A subset  $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$  of a vector space  $\mathcal{V}$  is said to be **linearly dependent** (LD) if there exist real numbers  $a_1, a_2, \dots, a_n$  not all zero such that

$$a_1\mathbf{v}_1 + a_2\mathbf{v}_2 + \dots + a_n\mathbf{v}_n = \mathbf{0}.$$

$S$  is **linearly independent** (LI) if it is not linearly dependent





## Section 4.4 (Linear Independence)

**Definition:** A subset  $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$  of a vector space  $\mathcal{V}$  is said to be **linearly dependent** (LD) if there exist real numbers  $a_1, a_2, \dots, a_n$  not all zero such that

$$a_1\mathbf{v}_1 + a_2\mathbf{v}_2 + \cdots + a_n\mathbf{v}_n = \mathbf{0}.$$

$S$  is **linearly independent** (LI) if it is not linearly dependent i.e. if

$$a_1\mathbf{v}_1 + a_2\mathbf{v}_2 + \cdots + a_n\mathbf{v}_n = \mathbf{0}$$

Then

$$a_1 = a_2 = \cdots = a_n = 0.$$



## Examples

- The subset  $S = \{[1, 0], [0, 1]\}$  of  $\mathbb{R}^2$  is linearly independent.



## Examples

- The subset  $S = \{[1, 0], [0, 1]\}$  of  $\mathbb{R}^2$  is linearly independent.
- The subset  $S = \{[1, 2], [5, 10]\}$  of  $\mathbb{R}^2$  is linearly dependent.



## Examples

- The subset  $S = \{[1, 0], [0, 1]\}$  of  $\mathbb{R}^2$  is linearly independent.
- The subset  $S = \{[1, 2], [5, 10]\}$  of  $\mathbb{R}^2$  is linearly dependent.
- The subset  $S = \{[1, 0, 0], [0, 1, 0], [0, 0, 1]\}$  of  $\mathbb{R}^3$  is linearly independent.



- The singleton set containing  $0 \in \mathcal{V}$  i.e.  $\{0\}$



- The singleton set containing  $0 \in \mathcal{V}$  i.e.  $\{0\}$  is LD.



- The singleton set containing  $0 \in \mathcal{V}$  i.e.  $\{0\}$  is LD.
- For  $\mathbf{v} \neq 0$  of  $\mathcal{V}$ , the set  $\{\mathbf{v}\}$



- The singleton set containing  $0 \in \mathcal{V}$  i.e.  $\{0\}$  is LD.
- For  $\mathbf{v} \neq 0$  of  $\mathcal{V}$ , the set  $\{\mathbf{v}\}$  is LI.





- The singleton set containing  $0 \in \mathcal{V}$  i.e.  $\{0\}$  is LD.
- For  $\mathbf{v} \neq 0$  of  $\mathcal{V}$ , the set  $\{\mathbf{v}\}$  is LI.
- Any set containing zero vector is



- The singleton set containing  $0 \in \mathcal{V}$  i.e.  $\{0\}$  is LD.
- For  $\mathbf{v} \neq 0$  of  $\mathcal{V}$ , the set  $\{\mathbf{v}\}$  is LI.
- Any set containing zero vector is LD.



- The singleton set containing  $0 \in \mathcal{V}$  i.e.  $\{0\}$  is LD.
- For  $\mathbf{v} \neq 0$  of  $\mathcal{V}$ , the set  $\{\mathbf{v}\}$  is LI.
- Any set containing zero vector is LD.
- Let  $S = \{\mathbf{v}_1, \mathbf{v}_2\}$  be a set of nonzero vectors of  $\mathcal{V}$ . Then  $S$  is linearly dependent iff one vector is a scalar multiple of the other.



- The singleton set containing  $0 \in \mathcal{V}$  i.e.  $\{0\}$  is LD.
- For  $\mathbf{v} \neq 0$  of  $\mathcal{V}$ , the set  $\{\mathbf{v}\}$  is LI.
- Any set containing zero vector is LD.
- Let  $S = \{\mathbf{v}_1, \mathbf{v}_2\}$  be a set of nonzero vectors of  $\mathcal{V}$ . Then  $S$  is linearly dependent iff one vector is a scalar multiple of the other.
- Let  $S$  be a finite set of nonzero vectors having at least two elements. Then  $S$  is LD if and only if some vector in  $S$  can be expressed as a linear combination of the other vectors in  $S$ .



**Exercise:** For a given vector space  $\mathcal{V}$  and a given subset  $S$  of  $\mathcal{V}$ , check the linear independence of  $S$  in the following:

1  $\mathcal{V} = \mathcal{P}_2, S = \{1 + x + x^2, 1 - x + 3x^2, 1 + 3x - x^2\}.$



**Exercise:** For a given vector space  $\mathcal{V}$  and a given subset  $S$  of  $\mathcal{V}$ , check the linear independence of  $S$  in the following:

- 1  $\mathcal{V} = \mathcal{P}_2, S = \{1 + x + x^2, 1 - x + 3x^2, 1 + 3x - x^2\}.$
- 2  $\mathcal{V} = \mathcal{P}_2, S = \{1 + x, x + x^2, 1 + x^2\}.$



**Exercise:** For a given vector space  $\mathcal{V}$  and a given subset  $S$  of  $\mathcal{V}$ , check the linear independence of  $S$  in the following:

- 1  $\mathcal{V} = \mathcal{P}_2, S = \{1 + x + x^2, 1 - x + 3x^2, 1 + 3x - x^2\}.$
- 2  $\mathcal{V} = \mathcal{P}_2, S = \{1 + x, x + x^2, 1 + x^2\}.$
- 3  $\mathcal{V} = \mathcal{P}_n, S = \{1, x, x^2, \dots, x^n\}.$



**Exercise:** For a given vector space  $\mathcal{V}$  and a given subset  $S$  of  $\mathcal{V}$ , check the linear independence of  $S$  in the following:

- 1  $\mathcal{V} = \mathcal{P}_2, S = \{1 + x + x^2, 1 - x + 3x^2, 1 + 3x - x^2\}.$
- 2  $\mathcal{V} = \mathcal{P}_2, S = \{1 + x, x + x^2, 1 + x^2\}.$
- 3  $\mathcal{V} = \mathcal{P}_n, S = \{1, x, x^2, \dots, x^n\}.$
- 4  $\mathcal{V} = \Phi, S = \{\sin^2 x, \cos^2 x, \cos 2x\}.$





**Exercise:** For a given vector space  $\mathcal{V}$  and a given subset  $S$  of  $\mathcal{V}$ , check the linear independence of  $S$  in the following:

- 1  $\mathcal{V} = \mathcal{P}_2, S = \{1 + x + x^2, 1 - x + 3x^2, 1 + 3x - x^2\}.$
- 2  $\mathcal{V} = \mathcal{P}_2, S = \{1 + x, x + x^2, 1 + x^2\}.$
- 3  $\mathcal{V} = \mathcal{P}_n, S = \{1, x, x^2, \dots, x^n\}.$
- 4  $\mathcal{V} = \Phi, S = \{\sin^2 x, \cos^2 x, \cos 2x\}.$
- 5  $\mathcal{V} = \Phi, S = \{\sin x, \cos x\}.$



**Exercise:** For a given vector space  $\mathcal{V}$  and a given subset  $S$  of  $\mathcal{V}$ , check the linear independence of  $S$  in the following:

1  $\mathcal{V} = \mathcal{P}_2, S = \{1 + x + x^2, 1 - x + 3x^2, 1 + 3x - x^2\}.$

2  $\mathcal{V} = \mathcal{P}_2, S = \{1 + x, x + x^2, 1 + x^2\}.$

3  $\mathcal{V} = \mathcal{P}_n, S = \{1, x, x^2, \dots, x^n\}.$

4  $\mathcal{V} = \Phi, S = \{\sin^2 x, \cos^2 x, \cos 2x\}.$

5  $\mathcal{V} = \Phi, S = \{\sin x, \cos x\}.$

6  $\mathcal{V} = \mathcal{M}_{22}, S = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}.$



**Exercise:** For a given vector space  $\mathcal{V}$  and a given subset  $S$  of  $\mathcal{V}$ , check the linear independence of  $S$  in the following:

1  $\mathcal{V} = \mathcal{P}_2, S = \{1 + x + x^2, 1 - x + 3x^2, 1 + 3x - x^2\}.$

2  $\mathcal{V} = \mathcal{P}_2, S = \{1 + x, x + x^2, 1 + x^2\}.$

3  $\mathcal{V} = \mathcal{P}_n, S = \{1, x, x^2, \dots, x^n\}.$

4  $\mathcal{V} = \Phi, S = \{\sin^2 x, \cos^2 x, \cos 2x\}.$

5  $\mathcal{V} = \Phi, S = \{\sin x, \cos x\}.$

6  $\mathcal{V} = \mathcal{M}_{22}, S = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}.$

7  $\mathcal{V} = \mathcal{M}_{22}, S = \left\{ \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & -1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 2 & 0 \\ 1 & 1 \end{bmatrix} \right\}.$



**Exercise:** Show that

$$S = \{[3, 1, -1], [-5, -2, 2], [2, 2, -1]\}$$

is linearly independent subset of  $\mathbb{R}^3$ .



**Exercise:** Show that

$$S = \{[3, 1, -1], [-5, -2, 2], [2, 2, -1]\}$$

is linearly independent subset of  $\mathbb{R}^3$ .

**Solution:** Let  $a, b, c \in \mathbb{R}$  such that

$$a[3, 1, -1] + b[-5, -2, 2] + c[2, 2, -1] = 0$$



**Exercise:** Show that

$$S = \{[3, 1, -1], [-5, -2, 2], [2, 2, -1]\}$$

is linearly independent subset of  $\mathbb{R}^3$ .

**Solution:** Let  $a, b, c \in \mathbb{R}$  such that

$$a[3, 1, -1] + b[-5, -2, 2] + c[2, 2, -1] = \mathbf{0}$$

$$[3a, a, -a] + [-5b, -2b, 2b] + [2c, 2c, -c] = [0, 0, 0]$$



**Exercise:** Show that

$$S = \{[3, 1, -1], [-5, -2, 2], [2, 2, -1]\}$$

is linearly independent subset of  $\mathbb{R}^3$ .

**Solution:** Let  $a, b, c \in \mathbb{R}$  such that

$$a[3, 1, -1] + b[-5, -2, 2] + c[2, 2, -1] = \mathbf{0}$$

$$[3a, a, -a] + [-5b, -2b, 2b] + [2c, 2c, -c] = [0, 0, 0]$$

$$[3a - 5b + 2c, a - 2b + 2c, -a + 2b - c] = [0, 0, 0]$$



To find  $a, b, c \in \mathbb{R}$ , we need to solve the following homogenous system:





To find  $a, b, c \in \mathbb{R}$ , we need to solve the following homogenous system:

$$3a - 5b + 2c = 0$$

$$a - 2b + 2c = 0$$

$$-a + 2b - c = 0$$



To find  $a, b, c \in \mathbb{R}$ , we need to solve the following homogenous system:

$$3a - 5b + 2c = 0$$

$$a - 2b + 2c = 0$$

$$-a + 2b - c = 0$$

To solve above homogenous system, write augmented matrix

$$[A|0] = \left[ \begin{array}{ccc|c} 3 & -5 & 2 & 0 \\ 1 & -2 & 2 & 0 \\ -1 & 2 & -1 & 0 \end{array} \right]$$



reduced row echelon form of  $[A|0]$  is

$$\left[ \begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right]$$

Thus, we have  $a = 0, b = 0, c = 0$ .



reduced row echelon form of  $[A|0]$  is

$$\left[ \begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right]$$

Thus, we have  $a = 0, b = 0, c = 0$ . Hence,  $S$  is linearly independent subset of  $\mathbb{R}^3$ .



**Independence Test Method:** Let  $S$  be a finite set of vectors in  $\mathbb{R}^n$ . To check whether  $S$  is LI, perform the following steps:



**Independence Test Method:** Let  $S$  be a finite set of vectors in  $\mathbb{R}^n$ . To check whether  $S$  is LI, perform the following steps:

**Step 1:** Form a matrix  $A$  whose **columns** are the vectors in  $S$ .



**Independence Test Method:** Let  $S$  be a finite set of vectors in  $\mathbb{R}^n$ . To check whether  $S$  is LI, perform the following steps:

**Step 1:** Form a matrix  $A$  whose **columns** are the vectors in  $S$ .

**Step 2:** Find  $\text{RREF}(A)$ .



**Independence Test Method:** Let  $S$  be a finite set of vectors in  $\mathbb{R}^n$ . To check whether  $S$  is LI, perform the following steps:

**Step 1:** Form a matrix  $A$  whose **columns** are the vectors in  $S$ .

**Step 2:** Find  $\text{RREF}(A)$ .

**Step 3:** If there is a pivot in every column of  $\text{RREF}(A)$ , then  $S$  is LI.





**Independence Test Method:** Let  $S$  be a finite set of vectors in  $\mathbb{R}^n$ . To check whether  $S$  is LI, perform the following steps:

**Step 1:** Form a matrix  $A$  whose **columns** are the vectors in  $S$ .

**Step 2:** Find  $\text{RREF}(A)$ .

**Step 3:** If there is a pivot in every column of  $\text{RREF}(A)$ , then  $S$  is LI. Otherwise  $S$  is LD.



**Theorem:** If  $S$  is any subset of  $\mathbb{R}^n$  containing  $k$  distinct vectors, where  $k > n$ , then  $S$  is linearly dependent.



**Theorem:** If  $S$  is any subset of  $\mathbb{R}^n$  containing  $k$  distinct vectors, where  $k > n$ , then  $S$  is linearly dependent.

**Exercise:** Examine the linear independence of a subset  $S = \{[2, -5, 1], [1, 1, -1], [0, 2, -3], [2, 2, 6]\}$  of  $\mathbb{R}^3$ .



**Theorem:** If  $S$  is any subset of  $\mathbb{R}^n$  containing  $k$  distinct vectors, where  $k > n$ , then  $S$  is linearly dependent.

**Exercise:** Examine the linear independence of a subset  $S = \{[2, -5, 1], [1, 1, -1], [0, 2, -3], [2, 2, 6]\}$  of  $\mathbb{R}^3$ .

**Result:** Let  $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$  be a linearly independent subset of a vector space  $\mathcal{V}$ . If  $\mathbf{v} \in \mathcal{V}$  and  $\mathbf{v} \notin \text{span}(S)$ , then  $S_1 = \{\mathbf{v}, \mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$  is a linearly independent set.



**Theorem:** A nonempty finite subset  $S$  of a vector space  $\mathcal{V}$  is LI iff every vector  $\mathbf{v} \in \text{span}(S)$  can be expressed **uniquely** as a linear combination of the elements of  $S$ .



**Definition:** An **infinite** subset  $S$  of a vector space  $\mathcal{V}$  is linearly dependent if there is some finite subset  $T$  of  $S$  such that  $T$  is linearly dependent.



**Definition:** An **infinite** subset  $S$  of a vector space  $\mathcal{V}$  is linearly dependent if there is some finite subset  $T$  of  $S$  such that  $T$  is linearly dependent.

**Example:** The subset

$$S = \{A \in \mathcal{M}_{22} \mid A \text{ is nonsingular}\}$$

of vector space  $\mathcal{M}_{22}$  is linearly dependent.



**Definition:** An **infinite** subset  $S$  of a vector space  $\mathcal{V}$  is linearly dependent if there is some finite subset  $T$  of  $S$  such that  $T$  is linearly dependent.

**Example:** The subset

$$S = \{A \in \mathcal{M}_{22} \mid A \text{ is nonsingular}\}$$

of vector space  $\mathcal{M}_{22}$  is linearly dependent.

**Solution:** Note that the finite subset  $T = \{I_2, 2I_2\}$  of  $S$  is linearly dependent





**Definition:** An **infinite** subset  $S$  of a vector space  $\mathcal{V}$  is linearly dependent if there is some finite subset  $T$  of  $S$  such that  $T$  is linearly dependent.

**Example:** The subset

$$S = \{A \in \mathcal{M}_{22} \mid A \text{ is nonsingular}\}$$

of vector space  $\mathcal{M}_{22}$  is linearly dependent.

**Solution:** Note that the finite subset  $T = \{I_2, 2I_2\}$  of  $S$  is linearly dependent as  $2I_2$  is scalar multiple of  $I_2$ .



**Definition:** An **infinite** subset  $S$  of a vector space  $\mathcal{V}$  is linearly dependent if there is some finite subset  $T$  of  $S$  such that  $T$  is linearly dependent.

**Example:** The subset

$$S = \{A \in \mathcal{M}_{22} \mid A \text{ is nonsingular}\}$$

of vector space  $\mathcal{M}_{22}$  is linearly dependent.

**Solution:** Note that the finite subset  $T = \{I_2, 2I_2\}$  of  $S$  is linearly dependent as  $2I_2$  is scalar multiple of  $I_2$ . Hence,  $S$  is linearly dependent.



**Definition:** An **infinite** subset  $S$  of a vector space  $\mathcal{V}$  is linearly independent if every finite subset of  $S$  is linearly independent.



**Definition:** An **infinite** subset  $S$  of a vector space  $\mathcal{V}$  is linearly independent if every finite subset of  $S$  is linearly independent.

**Result:** An infinite subset  $S$  of a vector space  $\mathcal{V}$  is linearly independent if and only if no vector in  $S$  is a finite linear combination of other vector in  $S$ .



**Definition:** An **infinite** subset  $S$  of a vector space  $\mathcal{V}$  is linearly independent if every finite subset of  $S$  is linearly independent.

**Result:** An infinite subset  $S$  of a vector space  $\mathcal{V}$  is linearly independent if and only if no vector in  $S$  is a finite linear combination of other vector in  $S$ .

**Example:** The subset  $S = \{1, x, x^2, x^3, x^4, \dots\}$  of vector space  $\mathcal{P}$  is



**Definition:** An **infinite** subset  $S$  of a vector space  $\mathcal{V}$  is linearly independent if every finite subset of  $S$  is linearly independent.

**Result:** An infinite subset  $S$  of a vector space  $\mathcal{V}$  is linearly independent if and only if no vector in  $S$  is a finite linear combination of other vector in  $S$ .

**Example:** The subset  $S = \{1, x, x^2, x^3, x^4, \dots\}$  of vector space  $\mathcal{P}$  is linearly independent.



# Section 4.5, Basis and Dimension



# Section 4.5, Basis and Dimension

**Basis:** A subset  $B$  of a vector space  $\mathcal{V}$  is said to be a **basis** of  $\mathcal{V}$  if

- 1  $B$  is LI, and
- 2  $\text{span}(B) = \mathcal{V}$ .





## Examples

- The subset  $B = \{[1, 0], [0, 1]\} = \{e_1, e_2\}$  is a basis of  $\mathbb{R}^2$  as  $B$  is LI and  $\text{span}(B) = \mathbb{R}^2$ .



## Examples

- The subset  $B = \{[1, 0], [0, 1]\} = \{e_1, e_2\}$  is a basis of  $\mathbb{R}^2$  as  $B$  is LI and  $\text{span}(B) = \mathbb{R}^2$ . The subset  $B$  is called the **standard basis** of  $\mathbb{R}^2$ .



## Examples

- The subset  $B = \{[1, 0], [0, 1]\} = \{e_1, e_2\}$  is a basis of  $\mathbb{R}^2$  as  $B$  is LI and  $\text{span}(B) = \mathbb{R}^2$ . The subset  $B$  is called the **standard basis** of  $\mathbb{R}^2$ .
- The subset  $B = \{[1, 2], [3, 4]\}$  is a basis of  $\mathbb{R}^2$  as  $B$  is LI (verify!) and  $\text{span}(B) = \mathbb{R}^2$  (verify!).



## Examples

- The subset  $B = \{[1, 0], [0, 1]\} = \{e_1, e_2\}$  is a basis of  $\mathbb{R}^2$  as  $B$  is LI and  $\text{span}(B) = \mathbb{R}^2$ . The subset  $B$  is called the **standard basis** of  $\mathbb{R}^2$ .
- The subset  $B = \{[1, 2], [3, 4]\}$  is a basis of  $\mathbb{R}^2$  as  $B$  is LI (verify!) and  $\text{span}(B) = \mathbb{R}^2$  (verify!).
- The subset  $B = \{[1, 0, 0], [0, 1, 0], [0, 0, 1]\}$ , also denoted by  $\{e_1, e_2, e_3\}$ , is a basis of  $\mathbb{R}^3$  as it is LI and  $\text{span}(B) = \mathbb{R}^3$ .



## Examples

- The subset  $B = \{[1, 0], [0, 1]\} = \{e_1, e_2\}$  is a basis of  $\mathbb{R}^2$  as  $B$  is LI and  $\text{span}(B) = \mathbb{R}^2$ . The subset  $B$  is called the **standard basis** of  $\mathbb{R}^2$ .
- The subset  $B = \{[1, 2], [3, 4]\}$  is a basis of  $\mathbb{R}^2$  as  $B$  is LI (verify!) and  $\text{span}(B) = \mathbb{R}^2$  (verify!).
- The subset  $B = \{[1, 0, 0], [0, 1, 0], [0, 0, 1]\}$ , also denoted by  $\{e_1, e_2, e_3\}$ , is a basis of  $\mathbb{R}^3$  as it is LI and  $\text{span}(B) = \mathbb{R}^3$ . The subset  $B$  is called the **standard basis** of  $\mathbb{R}^3$ .



## Examples

- The subset  $B = \{[1, 0], [0, 1]\} = \{e_1, e_2\}$  is a basis of  $\mathbb{R}^2$  as  $B$  is LI and  $\text{span}(B) = \mathbb{R}^2$ . The subset  $B$  is called the **standard basis** of  $\mathbb{R}^2$ .
- The subset  $B = \{[1, 2], [3, 4]\}$  is a basis of  $\mathbb{R}^2$  as  $B$  is LI (verify!) and  $\text{span}(B) = \mathbb{R}^2$  (verify!).
- The subset  $B = \{[1, 0, 0], [0, 1, 0], [0, 0, 1]\}$ , also denoted by  $\{e_1, e_2, e_3\}$ , is a basis of  $\mathbb{R}^3$  as it is LI and  $\text{span}(B) = \mathbb{R}^3$ . The subset  $B$  is called the **standard basis** of  $\mathbb{R}^3$ .

Think about some more basis of  $\mathbb{R}^2$  and  $\mathbb{R}^3$ .



- The subset  $B = \{1, x, x^2, \dots, x^n\}$  is a basis of  $\mathcal{P}_n$  as  $B$  is LI (verify!) and  $\text{span}(B) = \mathcal{P}_n$  (verify!).



- The subset  $B = \{1, x, x^2, \dots, x^n\}$  is a basis of  $\mathcal{P}_n$  as  $B$  is LI (verify!) and  $\text{span}(B) = \mathcal{P}_n$  (verify!).
- The subset

$$B = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}$$

is a basis of  $\mathcal{M}_{22}$ .





- The subset  $B = \{1, x, x^2, \dots, x^n\}$  is a basis of  $\mathcal{P}_n$  as  $B$  is LI (verify!) and  $\text{span}(B) = \mathcal{P}_n$  (verify!).
- The subset

$$B = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}$$

is a basis of  $\mathcal{M}_{22}$ .

Verify that  $B$  is LI and  $\text{span}(B) = \mathcal{M}_{22}$ .



**Theorem:** Every vector space has a basis.



**Theorem:** Every vector space has a basis.

**Theorem:** If a vector space  $V$  has a finite basis, then all bases for  $V$  are finite and have the same number of elements.



**Definition:** The **dimension** of a vector space  $\mathcal{V}$  is the number of elements in a basis of  $\mathcal{V}$



**Definition:** The **dimension** of a vector space  $\mathcal{V}$  is the number of elements in a basis of  $\mathcal{V}$  and it is denoted by  $\dim(\mathcal{V})$ .



**Definition:** The **dimension** of a vector space  $\mathcal{V}$  is the number of elements in a basis of  $\mathcal{V}$  and it is denoted by  $\dim(\mathcal{V})$ .

The dimension of the trivial vector space  $\{0\}$  is defined to be zero.



**Definition:** The **dimension** of a vector space  $\mathcal{V}$  is the number of elements in a basis of  $\mathcal{V}$  and it is denoted by  **$\dim(\mathcal{V})$** .

The dimension of the trivial vector space  $\{0\}$  is defined to be zero.

**Definition:** A vector space  $\mathcal{V}$  is said to be **finite dimensional** if it has a basis containing finite number of elements.



**Definition:** The **dimension** of a vector space  $\mathcal{V}$  is the number of elements in a basis of  $\mathcal{V}$  and it is denoted by  **$\dim(\mathcal{V})$** .

The dimension of the trivial vector space  $\{0\}$  is defined to be zero.

**Definition:** A vector space  $\mathcal{V}$  is said to be **finite dimensional** if it has a basis containing finite number of elements. If a vector space  $\mathcal{V}$  has no finite basis then  $\mathcal{V}$  is called **infinite dimensional**.





## Examples

- $\dim(\mathbb{R}^2) = 2.$



## Examples

- $\dim(\mathbb{R}^2) = 2.$
- $\dim(\mathbb{R}^3) = 3.$



## Examples

- $\dim(\mathbb{R}^2) = 2.$
- $\dim(\mathbb{R}^3) = 3.$
- $\dim(\mathbb{R}^n) = n.$



## Examples

- $\dim(\mathbb{R}^2) = 2.$
- $\dim(\mathbb{R}^3) = 3.$
- $\dim(\mathbb{R}^n) = n.$
- $\dim(\mathcal{P}_n) = n + 1.$



## Examples

- $\dim(\mathbb{R}^2) = 2.$
- $\dim(\mathbb{R}^3) = 3.$
- $\dim(\mathbb{R}^n) = n.$
- $\dim(\mathcal{P}_n) = n + 1.$
- $\dim(\mathcal{M}_{mn}) = mn.$



## Examples

- $\dim(\mathbb{R}^2) = 2.$
- $\dim(\mathbb{R}^3) = 3.$
- $\dim(\mathbb{R}^n) = n.$
- $\dim(\mathcal{P}_n) = n + 1.$
- $\dim(\mathcal{M}_{mn}) = mn.$
- Since  $\{1, x, x^2, x^3, \dots\}$  is a basis of  $\mathcal{P}$  (the vector space of all polynomials with real coefficients), thus the vector space  $\mathcal{P}$  is infinite dimensional.



**Exercise:** Find a basis and the dimension of a subspace  $W$  of  $\mathbb{R}^3$ , where

$$W = \{[x, y, z] \in \mathbb{R}^3 \mid x + 2z = 0\}.$$



**Exercise:** Find a basis and the dimension of a subspace  $W$  of  $\mathbb{R}^3$ , where

$$W = \{[x, y, z] \in \mathbb{R}^3 \mid x + 2z = 0\}.$$

**Solution:** The general solution of the equation  $x + 2z = 0$  is given by  $\{[-2s, t, s] \mid t, s \in \mathbb{R}\}$ . Thus

$$W = \{[-2s, t, s] \mid t, s \in \mathbb{R}\}$$





**Exercise:** Find a basis and the dimension of a subspace  $W$  of  $\mathbb{R}^3$ , where

$$W = \{[x, y, z] \in \mathbb{R}^3 \mid x + 2z = 0\}.$$

**Solution:** The general solution of the equation  $x + 2z = 0$  is given by  $\{[-2s, t, s] \mid t, s \in \mathbb{R}\}$ . Thus

$$W = \{[-2s, t, s] \mid t, s \in \mathbb{R}\}$$

$$W = \{s[-2, 0, 1] + t[0, 1, 0] \mid t, s \in \mathbb{R}\}$$



**Exercise:** Find a basis and the dimension of a subspace  $W$  of  $\mathbb{R}^3$ , where

$$W = \{[x, y, z] \in \mathbb{R}^3 \mid x + 2z = 0\}.$$

**Solution:** The general solution of the equation  $x + 2z = 0$  is given by  $\{[-2s, t, s] \mid t, s \in \mathbb{R}\}$ . Thus

$$W = \{[-2s, t, s] \mid t, s \in \mathbb{R}\}$$

$$W = \{s[-2, 0, 1] + t[0, 1, 0] \mid t, s \in \mathbb{R}\}$$

$$W = \text{span}(\{[-2, 0, 1], [0, 1, 0]\}).$$



**Exercise:** Find a basis and the dimension of a subspace  $W$  of  $\mathbb{R}^3$ , where

$$W = \{[x, y, z] \in \mathbb{R}^3 \mid x + 2z = 0\}.$$

**Solution:** The general solution of the equation  $x + 2z = 0$  is given by  $\{[-2s, t, s] \mid t, s \in \mathbb{R}\}$ . Thus

$$W = \{[-2s, t, s] \mid t, s \in \mathbb{R}\}$$

$$W = \{s[-2, 0, 1] + t[0, 1, 0] \mid t, s \in \mathbb{R}\}$$

$$W = \text{span}(\{[-2, 0, 1], [0, 1, 0]\}).$$

Note that the set  $\{[-2, 0, 1], [0, 1, 0]\}$  is linearly independent ([show it](#)).



Hence, the subset  $\{[-2, 0, 1], [0, 1, 0]\}$  is a basis of  $W$  and  $\dim(W) = 2$ .



Hence, the subset  $\{[-2, 0, 1], [0, 1, 0]\}$  is a basis of  $W$  and  $\dim(W) = 2$ .

**Exercise:** Find a basis and the dimension of a subspace  $W$  of  $\mathcal{P}_3$ , where

$$W = \{\mathbf{p} \in \mathcal{P}_3 \mid \mathbf{p}(2) = 0\}.$$



**Theorem:** Let  $W$  be a subspace of a finite dimensional vector space  $\mathcal{V}$ . Then

- $W$  is also finite dimensional and  $\dim W \leq \dim \mathcal{V}$ .
- $\dim W = \dim \mathcal{V}$  if and only if  $W = \mathcal{V}$ .



**Notation:**  $|A|$  - The number of elements in  $A$ .



**Notation:**  $|A|$ -The number of elements in  $A$ .

**Theorem:** Let  $\mathcal{V}$  be a finite dimensional vector space such that  $\dim(\mathcal{V}) = n$ .





**Notation:**  $|A|$ -The number of elements in  $A$ .

**Theorem:** Let  $\mathcal{V}$  be a finite dimensional vector space such that  $\dim(\mathcal{V}) = n$ .

- Suppose  $S$  is a finite subset of  $\mathcal{V}$  that spans  $\mathcal{V}$ .  
Then  $|S| \geq n$ .



**Notation:**  $|A|$ -The number of elements in  $A$ .

**Theorem:** Let  $\mathcal{V}$  be a finite dimensional vector space such that  $\dim(\mathcal{V}) = n$ .

- Suppose  $S$  is a finite subset of  $\mathcal{V}$  that spans  $\mathcal{V}$ . Then  $|S| \geq n$ . Moreover,  $|S| = n$  if and only if  $S$  is a basis of  $\mathcal{V}$ .



**Notation:**  $|A|$ -The number of elements in  $A$ .

**Theorem:** Let  $\mathcal{V}$  be a finite dimensional vector space such that  $\dim(\mathcal{V}) = n$ .

- Suppose  $S$  is a finite subset of  $\mathcal{V}$  that spans  $\mathcal{V}$ . Then  $|S| \geq n$ . Moreover,  $|S| = n$  if and only if  $S$  is a basis of  $\mathcal{V}$ .
- Suppose  $T$  is a linearly independent subset of  $\mathcal{V}$ . Then  $T$  is finite and  $|T| \leq n$ .



**Notation:**  $|A|$ -The number of elements in  $A$ .

**Theorem:** Let  $\mathcal{V}$  be a finite dimensional vector space such that  $\dim(\mathcal{V}) = n$ .

- Suppose  $S$  is a finite subset of  $\mathcal{V}$  that spans  $\mathcal{V}$ . Then  $|S| \geq n$ . Moreover,  $|S| = n$  if and only if  $S$  is a basis of  $\mathcal{V}$ .
- Suppose  $T$  is a linearly independent subset of  $\mathcal{V}$ . Then  $T$  is finite and  $|T| \leq n$ . Moreover,  $|T| = n$  if and only if  $T$  is a basis for  $\mathcal{V}$ .



**Exercise:** For a given vector space  $\mathcal{V}$  and a given subset  $B$  of  $\mathcal{V}$ , determine which of following  $B$  form a basis of the respective vector space  $\mathcal{V}$ :



**Exercise:** For a given vector space  $\mathcal{V}$  and a given subset  $B$  of  $\mathcal{V}$ , determine which of following  $B$  form a basis of the respective vector space  $\mathcal{V}$ :

1  $\mathcal{V} = \mathbb{R}^3$ ,  $B = \{[3, 1, -1], [-5, -2, 2], [2, 2, -1]\}$ .



**Exercise:** For a given vector space  $\mathcal{V}$  and a given subset  $B$  of  $\mathcal{V}$ , determine which of following  $B$  form a basis of the respective vector space  $\mathcal{V}$ :

- 1  $\mathcal{V} = \mathbb{R}^3$ ,  $B = \{[3, 1, -1], [-5, -2, 2], [2, 2, -1]\}$ .
- 2  $\mathcal{V} = \mathbb{R}^4$ ,  $B = \{[7, 1, 2, 0], [8, 0, 1, -1], [1, 0, 0, -2]\}$ .



**Exercise:** For a given vector space  $\mathcal{V}$  and a given subset  $B$  of  $\mathcal{V}$ , determine which of following  $B$  form a basis of the respective vector space  $\mathcal{V}$ :

- 1  $\mathcal{V} = \mathbb{R}^3$ ,  $B = \{[3, 1, -1], [-5, -2, 2], [2, 2, -1]\}$ .
- 2  $\mathcal{V} = \mathbb{R}^4$ ,  $B = \{[7, 1, 2, 0], [8, 0, 1, -1], [1, 0, 0, -2]\}$ .
- 3  $\mathcal{V} = \mathcal{P}_2$ ,  $B = \{1 + x, x + x^2, 1 + x^2\}$ .





**Exercise:** For a given vector space  $\mathcal{V}$  and a given subset  $B$  of  $\mathcal{V}$ , determine which of following  $B$  form a basis of the respective vector space  $\mathcal{V}$ :

- 1  $\mathcal{V} = \mathbb{R}^3$ ,  $B = \{[3, 1, -1], [-5, -2, 2], [2, 2, -1]\}$ .
- 2  $\mathcal{V} = \mathbb{R}^4$ ,  $B = \{[7, 1, 2, 0], [8, 0, 1, -1], [1, 0, 0, -2]\}$ .
- 3  $\mathcal{V} = \mathcal{P}_2$ ,  $B = \{1 + x, x + x^2, 1 + x^2\}$ .
- 4  $\mathcal{V} = \mathcal{P}_2$ ,  $B = \{1 - x, x - x^2, 1 - x^2\}$ .



**Exercise:** For a given vector space  $\mathcal{V}$  and a given subset  $B$  of  $\mathcal{V}$ , determine which of following  $B$  form a basis of the respective vector space  $\mathcal{V}$ :

- 1  $\mathcal{V} = \mathbb{R}^3$ ,  $B = \{[3, 1, -1], [-5, -2, 2], [2, 2, -1]\}$ .
- 2  $\mathcal{V} = \mathbb{R}^4$ ,  $B = \{[7, 1, 2, 0], [8, 0, 1, -1], [1, 0, 0, -2]\}$ .
- 3  $\mathcal{V} = \mathcal{P}_2$ ,  $B = \{1 + x, x + x^2, 1 + x^2\}$ .
- 4  $\mathcal{V} = \mathcal{P}_2$ ,  $B = \{1 - x, x - x^2, 1 - x^2\}$ .
- 5  $\mathcal{V} = \mathcal{M}_{22}$ ,

$$B = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \right\}.$$



**Exercise:** Let  $S = \{[1, 2], [3, 0], [0, 2]\} \subseteq \mathbb{R}^2$ .



**Exercise:** Let  $S = \{[1, 2], [3, 0], [0, 2]\} \subseteq \mathbb{R}^2$ . Find a basis for  $\text{span}(S)$ .



**Exercise:** Let  $S = \{[1, 2], [3, 0], [0, 2]\} \subseteq \mathbb{R}^2$ . Find a basis for  $\text{span}(S)$ .

**Solution:** In order to find a basis, we need a subset  $B$  of  $\text{span}(S)$  such that

- $\text{span}(S) = \text{span}(B)$ ;
- $B$  is LI.



**Exercise:** Let  $S = \{[1, 2], [3, 0], [0, 2]\} \subseteq \mathbb{R}^2$ . Find a basis for  $\text{span}(S)$ .

**Solution:** In order to find a basis, we need a subset  $B$  of  $\text{span}(S)$  such that

- $\text{span}(S) = \text{span}(B)$ ;
- $B$  is LI.

Since  $[1, 2] = \frac{1}{3}[3, 0] + [0, 2]$



**Exercise:** Let  $S = \{[1, 2], [3, 0], [0, 2]\} \subseteq \mathbb{R}^2$ . Find a basis for  $\text{span}(S)$ .

**Solution:** In order to find a basis, we need a subset  $B$  of  $\text{span}(S)$  such that

- $\text{span}(S) = \text{span}(B)$ ;
- $B$  is LI.

Since  $[1, 2] = \frac{1}{3}[3, 0] + [0, 2]$  implies

$$\text{span}(S) = \text{span}(B), \text{ where } B = \{[3, 0], [0, 2]\}.$$



**Exercise:** Let  $S = \{[1, 2], [3, 0], [0, 2]\} \subseteq \mathbb{R}^2$ . Find a basis for  $\text{span}(S)$ .

**Solution:** In order to find a basis, we need a subset  $B$  of  $\text{span}(S)$  such that

- $\text{span}(S) = \text{span}(B)$ ;
- $B$  is LI.

Since  $[1, 2] = \frac{1}{3}[3, 0] + [0, 2]$  implies

$$\text{span}(S) = \text{span}(B), \text{ where } B = \{[3, 0], [0, 2]\}.$$

Also,  $B$  is LI ([show it](#)).





**Exercise:** Let  $S = \{[1, 2], [3, 0], [0, 2]\} \subseteq \mathbb{R}^2$ . Find a basis for  $\text{span}(S)$ .

**Solution:** In order to find a basis, we need a subset  $B$  of  $\text{span}(S)$  such that

- $\text{span}(S) = \text{span}(B)$ ;
- $B$  is LI.

Since  $[1, 2] = \frac{1}{3}[3, 0] + [0, 2]$  implies

$$\text{span}(S) = \text{span}(B), \text{ where } B = \{[3, 0], [0, 2]\}.$$

Also,  $B$  is LI ([show it](#)). Hence,  $B$  is a basis of  $\text{span}(S)$ .



**Exercise:** Let  $W = \{X \in \mathbb{R}^5 : AX = 0\}$ ,



**Exercise:** Let  $W = \{X \in \mathbb{R}^5 : AX = 0\}$ , where

$$A = \begin{bmatrix} 1 & 2 & 1 & 0 & -1 \\ 2 & -1 & 0 & 1 & 3 \\ 1 & -3 & -1 & 1 & 4 \\ 2 & 9 & 4 & -1 & -7 \end{bmatrix}.$$

Find a basis for  $W$ .



**Exercise:** Let  $W = \{X \in \mathbb{R}^5 : AX = 0\}$ , where

$$A = \begin{bmatrix} 1 & 2 & 1 & 0 & -1 \\ 2 & -1 & 0 & 1 & 3 \\ 1 & -3 & -1 & 1 & 4 \\ 2 & 9 & 4 & -1 & -7 \end{bmatrix}.$$

Find a basis for  $W$ .

**Answer:**

$$B = \left\{ \begin{bmatrix} -\frac{1}{5} \\ -\frac{2}{5} \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -\frac{2}{5} \\ \frac{1}{5} \\ 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \\ 0 \\ 0 \\ 1 \end{bmatrix} \right\}$$



**Exercise:** Let  $S = \{[4, 2, 1], [2, 6, -5], [1, -2, 3]\}$  be a subset of vector space  $\mathbb{R}^3$ .

- Examine the linear independence of  $S$ .



**Exercise:** Let  $S = \{[4, 2, 1], [2, 6, -5], [1, -2, 3]\}$  be a subset of vector space  $\mathbb{R}^3$ .

- Examine the linear independence of  $S$ .
- Find  $\dim(\text{span}(S))$ .



**Exercise:** Let  $S = \{[4, 2, 1], [2, 6, -5], [1, -2, 3]\}$  be a subset of vector space  $\mathbb{R}^3$ .

- Examine the linear independence of  $S$ .
- Find  $\dim(\text{span}(S))$ .

**Hint:**

- Let

$$a_1[4, 2, 1] + a_2[2, 6, -5] + a_3[1, -2, 3] = \mathbf{0} = [0, 0, 0]$$

On solving above system of equations, we get

$$a_1 = -1, a_2 = 1, a_3 = 2$$

implies  $S$  is not LI.



- Note that

$$[2, 6, -5] = [4, 2, 1] - 2[1, -2, 3]$$

implies  $\text{span}(S) = \text{span}(B)$ , where

$$B = \{[4, 2, 1], [1, -2, 3]\}.$$

Now, note that  $B$  is LI





- Note that

$$[2, 6, -5] = [4, 2, 1] - 2[1, -2, 3]$$

implies  $\text{span}(S) = \text{span}(B)$ , where

$$B = \{[4, 2, 1], [1, -2, 3]\}.$$

Now, note that  $B$  is LI ([Show it](#)). Thus  $B$  (a set of two elements) is a basis of  $\text{span}(S)$



- Note that

$$[2, 6, -5] = [4, 2, 1] - 2[1, -2, 3]$$

implies  $\text{span}(S) = \text{span}(B)$ , where

$$B = \{[4, 2, 1], [1, -2, 3]\}.$$

Now, note that  $B$  is LI ([Show it](#)). Thus  $B$  (a set of two elements) is a basis of  $\text{span}(S)$  and

$$\dim(\text{span}(S)) = 2$$

.



# Simplified Span Method to find a Basis for $\text{span}(S)$



# Simplified Span Method to find a Basis for $\text{span}(S)$

Let  $S \subseteq \mathbb{R}^n$  containing  $k$  vectors.



## Simplified Span Method to find a Basis for $\text{span}(S)$

Let  $S \subseteq \mathbb{R}^n$  containing  $k$  vectors.

- Construct a matrix  $A$  of order  $k \times n$  by using vectors of  $S$



## Simplified Span Method to find a Basis for $\text{span}(S)$

Let  $S \subseteq \mathbb{R}^n$  containing  $k$  vectors.

- Construct a matrix  $A$  of order  $k \times n$  by using vectors of  $S$  **as rows of**  $A$ .



## Simplified Span Method to find a Basis for $\text{span}(S)$

Let  $S \subseteq \mathbb{R}^n$  containing  $k$  vectors.

- Construct a matrix  $A$  of order  $k \times n$  by using vectors of  $S$  **as rows of**  $A$ .
- Compute  $C = \text{RREF}(A)$ .



## Simplified Span Method to find a Basis for $\text{span}(S)$

Let  $S \subseteq \mathbb{R}^n$  containing  $k$  vectors.

- Construct a matrix  $A$  of order  $k \times n$  by using vectors of  $S$  **as rows of**  $A$ .
- Compute  $C = \text{RREF}(A)$ .
- **Nonzero rows** of  $C$  forms a basis for  $\text{span}(S)$ .





**Example:** Let  $S = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4\}$ , where

$$\mathbf{v}_1 = [1, 2, 3, -1, 0], \mathbf{v}_2 = [3, 6, 8, -2, 0]$$

$$\mathbf{v}_3 = [-1, -1, -3, 1, 1], \mathbf{v}_4 = [-2, -3, -5, 1, 1]$$

be a subset of  $\mathbb{R}^5$ .



**Example:** Let  $S = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4\}$ , where

$$\mathbf{v}_1 = [1, 2, 3, -1, 0], \mathbf{v}_2 = [3, 6, 8, -2, 0]$$

$$\mathbf{v}_3 = [-1, -1, -3, 1, 1], \mathbf{v}_4 = [-2, -3, -5, 1, 1]$$

be a subset of  $\mathbb{R}^5$ . Find a basis for  $\text{span}(S)$ .



**Example:** Let  $S = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4\}$ , where

$$\mathbf{v}_1 = [1, 2, 3, -1, 0], \mathbf{v}_2 = [3, 6, 8, -2, 0]$$

$$\mathbf{v}_3 = [-1, -1, -3, 1, 1], \mathbf{v}_4 = [-2, -3, -5, 1, 1]$$

be a subset of  $\mathbb{R}^5$ . Find a basis for  $\text{span}(S)$ .

**Solution:**

**Step 1:**

$$A = \begin{bmatrix} 1 & 2 & 3 & -1 & 0 \\ 3 & 6 & 8 & -2 & 0 \\ -1 & -1 & -3 & 1 & 1 \\ -2 & -3 & -5 & 1 & 1 \end{bmatrix}$$



## Step 2:

$$C = \text{RREF}(A) = \begin{bmatrix} 1 & 0 & 0 & 2 & -2 \\ 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$



## Step 2:

$$C = \text{RREF}(A) = \begin{bmatrix} 1 & 0 & 0 & 2 & -2 \\ 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

## Step 3:

$$B = \{[1, 0, 0, 2, -2], [0, 1, 0, 0, 1], [0, 0, 1, -1, 0]\}$$

is a basis for  $\text{span}(S)$ .



**Exercise:** Let

$$S = \{x^3 - 3x^2 + 2, 2x^3 - 7x^2 + x - 3, 4x^3 - 13x^2 + x + 5\}$$

be a subset of  $\mathcal{P}_3$ . Find a basis for  $\text{span}(S)$ .



**Exercise:** Let

$$S = \{x^3 - 3x^2 + 2, 2x^3 - 7x^2 + x - 3, 4x^3 - 13x^2 + x + 5\}$$

be a subset of  $\mathcal{P}_3$ . Find a basis for  $\text{span}(S)$ .

**Answer:**  $B = \{x^3 - 3x, x^2 - x, 1\}$ .



Next is to reduce a spanning set to a basis





Next is to reduce a spanning set to a basis by eliminating redundant vectors



Next is to reduce a spanning set to a basis by eliminating redundant vectors **without changing the form** of the original vectors.



Next is to reduce a spanning set to a basis by eliminating redundant vectors **without changing the form** of the original vectors.

**Theorem:** If  $S$  is a spanning set for a finite dimensional vector space  $\mathcal{V}$ , then there is a set  $B \subseteq S$  that is a basis for  $\mathcal{V}$ .



# Independence Test Method to find a Basis for $\mathcal{V} = \text{span}(S)$

Let  $S \subseteq \mathbb{R}^n$  containing  $k$  vectors.



# Independence Test Method to find a Basis for $\mathcal{V} = \text{span}(S)$

Let  $S \subseteq \mathbb{R}^n$  containing  $k$  vectors.

- Construct a matrix  $A$  of order  $n \times k$  by using vectors of  $S$



# Independence Test Method to find a Basis for $\mathcal{V} = \text{span}(S)$

Let  $S \subseteq \mathbb{R}^n$  containing  $k$  vectors.

- Construct a matrix  $A$  of order  $n \times k$  by using vectors of  $S$  **as columns of  $A$ .**



# Independence Test Method to find a Basis for $\mathcal{V} = \text{span}(S)$

Let  $S \subseteq \mathbb{R}^n$  containing  $k$  vectors.

- Construct a matrix  $A$  of order  $n \times k$  by using vectors of  $S$  **as columns of  $A$** .
- Compute  $C = \text{RREF}(A)$ .



# Independence Test Method to find a Basis for $\mathcal{V} = \text{span}(S)$

Let  $S \subseteq \mathbb{R}^n$  containing  $k$  vectors.

- Construct a matrix  $A$  of order  $n \times k$  by using vectors of  $S$  **as columns of  $A$** .
- Compute  $C = \text{RREF}(A)$ .
- Column vectors in  $A$  corresponding to **pivot columns** of  $C$  forms a basis for  $\text{span}(S)$ .





**Example:** Let  $S = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4, \mathbf{v}_5\}$ , where

$$\mathbf{v}_1 = [1, 2, -2, 1], \quad \mathbf{v}_2 = [-3, 0, -4, 3]$$

$$\mathbf{v}_3 = [2, 1, 1, -1], \quad \mathbf{v}_4 = [-3, 3, -9, 6]$$

$$\text{and } \mathbf{v}_5 = [9, 3, 7, -6]$$

be a subset of  $\mathbb{R}^4$ .



**Example:** Let  $S = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4, \mathbf{v}_5\}$ , where

$$\mathbf{v}_1 = [1, 2, -2, 1], \quad \mathbf{v}_2 = [-3, 0, -4, 3]$$

$$\mathbf{v}_3 = [2, 1, 1, -1], \quad \mathbf{v}_4 = [-3, 3, -9, 6]$$

$$\text{and } \mathbf{v}_5 = [9, 3, 7, -6]$$

be a subset of  $\mathbb{R}^4$ . Find a basis for  $\text{span}(S)$ .



**Example:** Let  $S = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4, \mathbf{v}_5\}$ , where

$$\mathbf{v}_1 = [1, 2, -2, 1], \quad \mathbf{v}_2 = [-3, 0, -4, 3]$$

$$\mathbf{v}_3 = [2, 1, 1, -1], \quad \mathbf{v}_4 = [-3, 3, -9, 6]$$

$$\text{and } \mathbf{v}_5 = [9, 3, 7, -6]$$

be a subset of  $\mathbb{R}^4$ . Find a basis for  $\text{span}(S)$ .

**Solution:**

$$A = \begin{bmatrix} 1 & -3 & 2 & -3 & 9 \\ 2 & 0 & 1 & 3 & 3 \\ -2 & -4 & 1 & -9 & 7 \\ 1 & 3 & -1 & 6 & -6 \end{bmatrix}$$



$$C = \text{RREF}(A) = \begin{bmatrix} 1 & 0 & 1/2 & 3/2 & 3/2 \\ 0 & 1 & -1/2 & 3/2 & -5/2 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

implies



$$C = \text{RREF}(A) = \begin{bmatrix} 1 & 0 & 1/2 & 3/2 & 3/2 \\ 0 & 1 & -1/2 & 3/2 & -5/2 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

implies the set of vectors corresponding to pivot columns i.e.

$$B = \{[1, 2, -2, 1], [-3, 0, -4, 3]\}$$



$$C = \text{RREF}(A) = \begin{bmatrix} \color{red}{1} & 0 & 1/2 & 3/2 & 3/2 \\ 0 & \color{red}{1} & -1/2 & 3/2 & -5/2 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

implies the set of vectors corresponding to pivot columns i.e.

$$B = \{[1, 2, -2, 1], [-3, 0, -4, 3]\}$$

forms a basis for the subspace  $\text{span}(S)$ .



**Exercise:** Let

$$S = \{x^3 - 3x^2 + 1, 2x^2 + x, 2x^3 + 3x + 2, 4x - 5\}$$

be a subset of  $\mathcal{P}_3$ . Find a basis for  $\text{span}(S)$ .



**Exercise:** Let

$$S = \{x^3 - 3x^2 + 1, 2x^2 + x, 2x^3 + 3x + 2, 4x - 5\}$$

be a subset of  $\mathcal{P}_3$ . Find a basis for  $\text{span}(S)$ .

**Answer:**  $B = \{x^3 - 3x^2 + 1, 2x^2 + x, 4x - 5\}$ .





**Example:** Let  $S = \{[1, 0, 1, 0], [-1, 1, -1, 0]\}$  be LI a subset of  $\mathbb{R}^4$ .



**Example:** Let  $S = \{[1, 0, 1, 0], [-1, 1, -1, 0]\}$  be LI a subset of  $\mathbb{R}^4$ . Extend  $S$  to form a basis for  $\mathbb{R}^4$ .



**Example:** Let  $S = \{[1, 0, 1, 0], [-1, 1, -1, 0]\}$  be LI a subset of  $\mathbb{R}^4$ . Extend  $S$  to form a basis for  $\mathbb{R}^4$ .

**Note that** the set  $S$  is **NOT** a basis for  $\mathbb{R}^4$  (**Why?**).



**Example:** Let  $S = \{[1, 0, 1, 0], [-1, 1, -1, 0]\}$  be LI a subset of  $\mathbb{R}^4$ . Extend  $S$  to form a basis for  $\mathbb{R}^4$ .

**Note that** the set  $S$  is **NOT** a basis for  $\mathbb{R}^4$  (**Why?**).

**Solution:**



**Example:** Let  $S = \{[1, 0, 1, 0], [-1, 1, -1, 0]\}$  be LI a subset of  $\mathbb{R}^4$ . Extend  $S$  to form a basis for  $\mathbb{R}^4$ .

**Note that** the set  $S$  is **NOT** a basis for  $\mathbb{R}^4$  (**Why?**).

**Solution:** We know that  $B' = \{e_1, e_2, e_3, e_4\}$  is a standard basis for  $\mathbb{R}^4$ .



**Example:** Let  $S = \{[1, 0, 1, 0], [-1, 1, -1, 0]\}$  be LI a subset of  $\mathbb{R}^4$ . Extend  $S$  to form a basis for  $\mathbb{R}^4$ .

**Note that** the set  $S$  is **NOT** a basis for  $\mathbb{R}^4$  (**Why?**).

**Solution:** We know that  $B' = \{e_1, e_2, e_3, e_4\}$  is a standard basis for  $\mathbb{R}^4$ . Consider

$$B_1 = \{[1, 0, 1, 0], [-1, 1, -1, 0], e_1, e_2, e_3, e_4\}.$$



**Example:** Let  $S = \{[1, 0, 1, 0], [-1, 1, -1, 0]\}$  be LI a subset of  $\mathbb{R}^4$ . Extend  $S$  to form a basis for  $\mathbb{R}^4$ .

**Note that** the set  $S$  is **NOT** a basis for  $\mathbb{R}^4$  (**Why?**).

**Solution:** We know that  $B' = \{e_1, e_2, e_3, e_4\}$  is a standard basis for  $\mathbb{R}^4$ . Consider

$$B_1 = \{[1, 0, 1, 0], [-1, 1, -1, 0], e_1, e_2, e_3, e_4\}.$$

Now  $B'$  spans  $\mathbb{R}^4$  implies



**Example:** Let  $S = \{[1, 0, 1, 0], [-1, 1, -1, 0]\}$  be LI a subset of  $\mathbb{R}^4$ . Extend  $S$  to form a basis for  $\mathbb{R}^4$ .

**Note that** the set  $S$  is **NOT** a basis for  $\mathbb{R}^4$  (**Why?**).

**Solution:** We know that  $B' = \{e_1, e_2, e_3, e_4\}$  is a standard basis for  $\mathbb{R}^4$ . Consider

$$B_1 = \{[1, 0, 1, 0], [-1, 1, -1, 0], e_1, e_2, e_3, e_4\}.$$

Now  $B'$  spans  $\mathbb{R}^4$  implies  $\text{span}(B_1) = \mathbb{R}^4$ .





**Example:** Let  $S = \{[1, 0, 1, 0], [-1, 1, -1, 0]\}$  be LI a subset of  $\mathbb{R}^4$ . Extend  $S$  to form a basis for  $\mathbb{R}^4$ .

**Note that** the set  $S$  is **NOT** a basis for  $\mathbb{R}^4$  (**Why?**).

**Solution:** We know that  $B' = \{e_1, e_2, e_3, e_4\}$  is a standard basis for  $\mathbb{R}^4$ . Consider

$$B_1 = \{[1, 0, 1, 0], [-1, 1, -1, 0], e_1, e_2, e_3, e_4\}.$$

Now  $B'$  spans  $\mathbb{R}^4$  implies  $\text{span}(B_1) = \mathbb{R}^4$ . Consider

$$A = \begin{bmatrix} 1 & -1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 1 & -1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$



$$C = \text{RREF}(A) = \begin{bmatrix} 1 & 0 & 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$



$$C = \text{RREF}(A) = \begin{bmatrix} 1 & 0 & 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Using Independence Test Method, we have



$$C = \text{RREF}(A) = \begin{bmatrix} 1 & 0 & 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Using Independence Test Method, we have

$$B = \{[1, 0, 1, 0], [-1, 1, -1, 0], e_1, e_4\}$$

is a basis of  $\mathbb{R}^4$  containing  $S$ .



**Theorem:** Every LI subset of a finite dimensional vector space  $\mathcal{V}$  can be extended to form a basis of  $\mathcal{V}$ .



# Method for finding a Basis by Enlarging a LI Subset



## Method for finding a Basis by Enlarging a LI Subset

Let  $T = \{t_1, \dots, t_k\}$  be a LI subset of a finite dimensional vector space  $\mathcal{V}$ .



## Method for finding a Basis by Enlarging a LI Subset

Let  $T = \{t_1, \dots, t_k\}$  be a LI subset of a finite dimensional vector space  $\mathcal{V}$ .

- Find a spanning set  $A = \{a_1, \dots, a_n\}$  for  $\mathcal{V}$ .





## Method for finding a Basis by Enlarging a LI Subset

Let  $T = \{t_1, \dots, t_k\}$  be a LI subset of a finite dimensional vector space  $\mathcal{V}$ .

- Find a spanning set  $A = \{a_1, \dots, a_n\}$  for  $\mathcal{V}$ .
- Form the ordered spanning set

$$S = \{t_1, \dots, t_k, a_1, \dots, a_n\}$$

for  $\mathcal{V}$ .



## Method for finding a Basis by Enlarging a LI Subset

Let  $T = \{t_1, \dots, t_k\}$  be a LI subset of a finite dimensional vector space  $\mathcal{V}$ .

- Find a spanning set  $A = \{a_1, \dots, a_n\}$  for  $\mathcal{V}$ .
- Form the ordered spanning set

$$S = \{t_1, \dots, t_k, a_1, \dots, a_n\}$$

for  $\mathcal{V}$ .

- Use Independence Test Method to produce a subset  $B$  of  $S$ .



## Method for finding a Basis by Enlarging a LI Subset

Let  $T = \{t_1, \dots, t_k\}$  be a LI subset of a finite dimensional vector space  $\mathcal{V}$ .

- Find a spanning set  $A = \{a_1, \dots, a_n\}$  for  $\mathcal{V}$ .
- Form the ordered spanning set

$$S = \{t_1, \dots, t_k, a_1, \dots, a_n\}$$

for  $\mathcal{V}$ .

- Use Independence Test Method to produce a subset  $B$  of  $S$ . Then  $B$  is a basis for  $\mathcal{V}$  containing  $T$ .



**Example:** Let

$$T = \{x^3 - x^2, x^4 - 3x^3 + 5x^2 - x\}$$

be a LI subset of  $\mathcal{P}_4$ .



**Example:** Let

$$T = \{x^3 - x^2, x^4 - 3x^3 + 5x^2 - x\}$$

be a LI subset of  $\mathcal{P}_4$ . Extend  $T$  to form a basis for  $\mathcal{P}_4$ .



**Example:** Let

$$T = \{x^3 - x^2, x^4 - 3x^3 + 5x^2 - x\}$$

be a LI subset of  $\mathcal{P}_4$ . Extend  $T$  to form a basis for  $\mathcal{P}_4$ .

**Solution:** By considering the polynomials in  $T$  as vectors in  $\mathbb{R}^5$ , note that



**Example:** Let

$$T = \{x^3 - x^2, x^4 - 3x^3 + 5x^2 - x\}$$

be a LI subset of  $\mathcal{P}_4$ . Extend  $T$  to form a basis for  $\mathcal{P}_4$ .

**Solution:** By considering the polynomials in  $T$  as vectors in  $\mathbb{R}^5$ , note that

$$T = \{\mathbf{v}_1, \mathbf{v}_2\},$$



**Example:** Let

$$T = \{x^3 - x^2, x^4 - 3x^3 + 5x^2 - x\}$$

be a LI subset of  $\mathcal{P}_4$ . Extend  $T$  to form a basis for  $\mathcal{P}_4$ .

**Solution:** By considering the polynomials in  $T$  as vectors in  $\mathbb{R}^5$ , note that

$$T = \{\mathbf{v}_1, \mathbf{v}_2\},$$

where  $\mathbf{v}_1 = [0, 1, -1, 0, 0]$  and  $\mathbf{v}_2 = [1, -3, 5, -1, 0]$ .





**Step 1:** We know that  $A = \{e_1, e_2, e_3, e_4, e_5\}$  is a standard basis for  $\mathbb{R}^5$ .



**Step 1:** We know that  $A = \{e_1, e_2, e_3, e_4, e_5\}$  is a standard basis for  $\mathbb{R}^5$ .

**Step 2:** Now we form the spanning set

$$S = \{\mathbf{v}_1, \mathbf{v}_2, e_1, e_2, e_3, e_4, e_5\}.$$

of  $\mathbb{R}^5$ .



**Step 1:** We know that  $A = \{e_1, e_2, e_3, e_4, e_5\}$  is a standard basis for  $\mathbb{R}^5$ .

**Step 2:** Now we form the spanning set

$$S = \{\mathbf{v}_1, \mathbf{v}_2, e_1, e_2, e_3, e_4, e_5\}.$$

of  $\mathbb{R}^5$ .

**Step 3:**

$$C = \begin{bmatrix} 0 & 1 & 1 & 0 & 0 & 0 & 0 \\ 1 & -3 & 0 & 1 & 0 & 0 & 0 \\ -1 & 5 & 0 & 0 & 1 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$



$$\text{RREF}(C) = \begin{bmatrix} 1 & 0 & 0 & 0 & -1 & -5 & 0 \\ 0 & 1 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$



$$\text{RREF}(C) = \begin{bmatrix} 1 & 0 & 0 & 0 & -1 & -5 & 0 \\ 0 & 1 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Since columns I, II, III, IV and VII are pivot columns, so by independent test method, the set

$$B = \{\mathbf{v}_1, \mathbf{v}_2, e_1, e_2, e_5\}$$

is a basis of  $\text{span}(S)$ ,



$$\text{RREF}(C) = \begin{bmatrix} 1 & 0 & 0 & 0 & -1 & -5 & 0 \\ 0 & 1 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Since columns I, II, III, IV and VII are pivot columns, so by independent test method, the set

$$B = \{\mathbf{v}_1, \mathbf{v}_2, e_1, e_2, e_5\}$$

is a basis of  $\text{span}(S)$ , i.e.

$$B = \{x^3 - x^2, x^4 - 3x^3 + 5x^2 - x, x^4, x^3, 1\}$$

is a basis of  $\mathcal{P}_4$  containing  $T$ .



**Exercise:** Let

$$T = \left\{ \begin{bmatrix} 1 & -1 \\ -1 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & -1 \\ -1 & 1 \end{bmatrix} \right\}$$

be a LI subset of  $\mathcal{M}_{32}$ .



**Exercise:** Let

$$T = \left\{ \begin{bmatrix} 1 & -1 \\ -1 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & -1 \\ -1 & 1 \end{bmatrix} \right\}$$

be a LI subset of  $\mathcal{M}_{32}$ . Extend  $T$  to form a basis for  $\mathcal{M}_{32}$ .





**Exercise:** Let

$$T = \left\{ \begin{bmatrix} 1 & -1 \\ -1 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & -1 \\ -1 & 1 \end{bmatrix} \right\}$$

be a LI subset of  $\mathcal{M}_{32}$ . Extend  $T$  to form a basis for  $\mathcal{M}_{32}$ .

**Answer:**

$$B = \left\{ \begin{bmatrix} 1 & -1 \\ -1 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & -1 \\ -1 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \\ 1 & 0 \end{bmatrix} \right\}$$



# ***Thank You***

