

# MATHEMATICS-II (MATH F112)

**Dr. Krishnendra Shekhawat**

**BITS PILANI**  
**Department of Mathematics**



# Section 5.5

## *Isomorphism*



## Isomorphism

A LT  $L : V \rightarrow W$  that is both one-to-one and onto is called as **isomorphism** from  $V$  to  $W$ .



## Example 1

**Q:.** Check if  $L : P_n \rightarrow P_n$  given by  $L(p) = p + p'$  is an isomorphism.



## Example 1

**Q:.** Check if  $L : P_n \rightarrow P_n$  given by  $L(p) = p + p'$  is an isomorphism.

**Sol.** First we need to show that



## Example 1

**Q:.** Check if  $L : P_n \rightarrow P_n$  given by  $L(p) = p + p'$  is an isomorphism.

**Sol.** First we need to show that  $L$  is a linear operator.



## Example 1

**Q:.** Check if  $L : P_n \rightarrow P_n$  given by  $L(p) = p + p'$  is an isomorphism.

**Sol.** First we need to show that  $L$  is a linear operator.

$$L(p + q) = (p + q) + (p + q)' = p + p' + q + q' = L(p) + L(q)$$

for all  $p, q \in P_n$ .



## Example 1

**Q:.** Check if  $L : P_n \rightarrow P_n$  given by  $L(p) = p + p'$  is an isomorphism.

**Sol.** First we need to show that  $L$  is a linear operator.

$$L(p + q) = (p + q) + (p + q)' = p + p' + q + q' = L(p) + L(q)$$

for all  $p, q \in P_n$ .

Similarly,  $L(cp) = cL(p)$  for all real  $c$  and  $p \in P_n$ .





## Example 1

**Q:.** Check if  $L : P_n \rightarrow P_n$  given by  $L(p) = p + p'$  is an isomorphism.

**Sol.** First we need to show that  $L$  is a linear operator.

$$L(p + q) = (p + q) + (p + q)' = p + p' + q + q' = L(p) + L(q)$$

for all  $p, q \in P_n$ .

Similarly,  $L(cp) = cL(p)$  for all real  $c$  and  $p \in P_n$ . Hence,  $L$  is a linear operator.



Now,  $\ker(L) = \{p | L(p) = 0\}$ .



Now,  $\ker(L) = \{p | L(p) = 0\}$ .

$$L(p) = 0 \implies p + p' = 0 \implies p = -p'.$$



Now,  $\ker(L) = \{p | L(p) = 0\}$ .

$$L(p) = 0 \implies p + p' = 0 \implies p = -p'.$$

If  $p$  is not a constant polynomial,  $p$  and  $p'$  must have different degrees, i.e.,



Now,  $\ker(L) = \{p | L(p) = 0\}$ .

$$L(p) = 0 \implies p + p' = 0 \implies p = -p'.$$

If  $p$  is not a constant polynomial,  $p$  and  $p'$  must have different degrees, i.e.,  $p = -p'$  is a contradiction



Now,  $\ker(L) = \{p | L(p) = 0\}$ .

$$L(p) = 0 \implies p + p' = 0 \implies p = -p'.$$

If  $p$  is not a constant polynomial,  $p$  and  $p'$  must have different degrees, i.e.,  $p = -p'$  is a contradiction  
 $\implies p = 0$  and  $p' = 0$



Now,  $\ker(L) = \{p | L(p) = 0\}$ .

$$L(p) = 0 \implies p + p' = 0 \implies p = -p'.$$

If  $p$  is not a constant polynomial,  $p$  and  $p'$  must have different degrees, i.e.,  $p = -p'$  is a contradiction  
 $\implies p = 0$  and  $p' = 0 \implies \ker(L) = \{0\}$



Now,  $\ker(L) = \{p | L(p) = 0\}$ .

$$L(p) = 0 \implies p + p' = 0 \implies p = -p'.$$

If  $p$  is not a constant polynomial,  $p$  and  $p'$  must have different degrees, i.e.,  $p = -p'$  is a contradiction  
 $\implies p = 0$  and  $p' = 0 \implies \ker(L) = \{0\} \implies L$  is one-to-one.





Now,  $\ker(L) = \{p | L(p) = 0\}$ .

$$L(p) = 0 \implies p + p' = 0 \implies p = -p'.$$

If  $p$  is not a constant polynomial,  $p$  and  $p'$  must have different degrees, i.e.,  $p = -p'$  is a contradiction  
 $\implies p = 0$  and  $p' = 0 \implies \ker(L) = \{0\} \implies L$  is one-to-one.

Also,  $L$  is onto



Now,  $\ker(L) = \{p | L(p) = 0\}$ .

$$L(p) = 0 \implies p + p' = 0 \implies p = -p'.$$

If  $p$  is not a constant polynomial,  $p$  and  $p'$  must have different degrees, i.e.,  $p = -p'$  is a contradiction  
 $\implies p = 0$  and  $p' = 0 \implies \ker(L) = \{0\} \implies L$  is one-to-one.

Also,  $L$  is onto because  $\dim \text{range}(L) = \dim(P_n) = n + 1$ .



Now,  $\ker(L) = \{p | L(p) = 0\}$ .

$$L(p) = 0 \implies p + p' = 0 \implies p = -p'.$$

If  $p$  is not a constant polynomial,  $p$  and  $p'$  must have different degrees, i.e.,  $p = -p'$  is a contradiction  
 $\implies p = 0$  and  $p' = 0 \implies \ker(L) = \{0\} \implies L$  is one-to-one.

Also,  $L$  is onto because  $\dim \text{range}(L) = \dim(P_n) = n + 1$ .

Hence,  $L$  is an isomorphism.



## Example 2

**Q:.** Check if the linear operator  $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  given by  $L(e_1) = e_1 + e_2, L(e_2) = e_2 + e_3, L(e_3) = e_1 + e_2 + e_3$  is an isomorphism.



## Example 2

**Q:.** Check if the linear operator  $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  given by  $L(e_1) = e_1 + e_2, L(e_2) = e_2 + e_3, L(e_3) = e_1 + e_2 + e_3$  is an isomorphism.

**Sol.** For  $u = [x, y, z] \in \mathbb{R}^3$ ,



## Example 2

**Q:.** Check if the linear operator  $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  given by  $L(e_1) = e_1 + e_2, L(e_2) = e_2 + e_3, L(e_3) = e_1 + e_2 + e_3$  is an isomorphism.

**Sol.** For  $u = [x, y, z] \in \mathbb{R}^3$ ,  $L(u) = L([x, y, z]) =$



## Example 2

**Q:.** Check if the linear operator  $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  given by  $L(e_1) = e_1 + e_2, L(e_2) = e_2 + e_3, L(e_3) = e_1 + e_2 + e_3$  is an isomorphism.

**Sol.** For  $u = [x, y, z] \in \mathbb{R}^3$ ,  $L(u) = L([x, y, z]) = L([xe_1 + ye_2 + ze_3]) = xL(e_1) + yL(e_2) + zL(e_3) = [x + z, x + y + z, y + z]$ .



## Example 2

**Q.:** Check if the linear operator  $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  given by  $L(e_1) = e_1 + e_2, L(e_2) = e_2 + e_3, L(e_3) = e_1 + e_2 + e_3$  is an isomorphism.

**Sol.** For  $u = [x, y, z] \in \mathbb{R}^3$ ,  $L(u) = L([x, y, z]) = L([xe_1 + ye_2 + ze_3]) = xL(e_1) + yL(e_2) + zL(e_3) = [x + z, x + y + z, y + z]$ .

Now  $L([x, y, z]) = [0, 0, 0] \implies$





## Example 2

**Q.:** Check if the linear operator  $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  given by  $L(e_1) = e_1 + e_2, L(e_2) = e_2 + e_3, L(e_3) = e_1 + e_2 + e_3$  is an isomorphism.

**Sol.** For  $u = [x, y, z] \in \mathbb{R}^3$ ,  $L(u) = L([x, y, z]) = L([xe_1 + ye_2 + ze_3]) = xL(e_1) + yL(e_2) + zL(e_3) = [x + z, x + y + z, y + z]$ .

Now  $L([x, y, z]) = [0, 0, 0] \implies$

$[x + z, x + y + z, y + z] = [0, 0, 0] \implies x = y = z = 0.$



## Example 2

**Q.:** Check if the linear operator  $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  given by  $L(e_1) = e_1 + e_2, L(e_2) = e_2 + e_3, L(e_3) = e_1 + e_2 + e_3$  is an isomorphism.

**Sol.** For  $u = [x, y, z] \in \mathbb{R}^3$ ,  $L(u) = L([x, y, z]) = L([xe_1 + ye_2 + ze_3]) = xL(e_1) + yL(e_2) + zL(e_3) = [x + z, x + y + z, y + z]$ .

Now  $L([x, y, z]) = [0, 0, 0] \implies$

$[x + z, x + y + z, y + z] = [0, 0, 0] \implies x = y = z = 0$ . Since  $\ker(L) = \{0_{\mathbb{R}^3}\}$ ,  $L$  is one-to-one.



## Example 2

**Q.:** Check if the linear operator  $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  given by  $L(e_1) = e_1 + e_2, L(e_2) = e_2 + e_3, L(e_3) = e_1 + e_2 + e_3$  is an isomorphism.

**Sol.** For  $u = [x, y, z] \in \mathbb{R}^3$ ,  $L(u) = L([x, y, z]) = L([xe_1 + ye_2 + ze_3]) = xL(e_1) + yL(e_2) + zL(e_3) = [x + z, x + y + z, y + z]$ .

Now  $L([x, y, z]) = [0, 0, 0] \implies$

$[x + z, x + y + z, y + z] = [0, 0, 0] \implies x = y = z = 0$ . Since  $\ker(L) = \{0_{\mathbb{R}^3}\}$ ,  $L$  is one-to-one.  $L$  is also onto since  $\dim(\text{range}(L)) = 3 - 0 = 3 = \dim(\mathbb{R}^3)$ , from Dimension Theorem.



## Example 2

**Q.:** Check if the linear operator  $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  given by  $L(e_1) = e_1 + e_2, L(e_2) = e_2 + e_3, L(e_3) = e_1 + e_2 + e_3$  is an isomorphism.

**Sol.** For  $u = [x, y, z] \in \mathbb{R}^3$ ,  $L(u) = L([x, y, z]) = L([xe_1 + ye_2 + ze_3]) = xL(e_1) + yL(e_2) + zL(e_3) = [x + z, x + y + z, y + z]$ .

Now  $L([x, y, z]) = [0, 0, 0] \implies$

$[x + z, x + y + z, y + z] = [0, 0, 0] \implies x = y = z = 0$ . Since  $\ker(L) = \{0_{\mathbb{R}^3}\}$ ,  $L$  is one-to-one.  $L$  is also onto since  $\dim(\text{range}(L)) = 3 - 0 = 3 = \dim(\mathbb{R}^3)$ , from Dimension Theorem. Hence,  $L$  is an isomorphism.



### Example 3

**Q:.** Check if the LT  $L : P_4 \rightarrow \mathbb{R}^3$  given by  $L(p) = [p(-1), p(0), p(1)]$  is an isomorphism.



### Example 3

**Q:.** Check if the LT  $L : P_4 \rightarrow \mathbb{R}^3$  given by  $L(p) = [p(-1), p(0), p(1)]$  is an isomorphism.

**Sol.** Basis for  $\ker(L)$  is  $\{x(x^2 - 1), x^2(x^2 - 1)\}$ .



### Example 3

**Q:.** Check if the LT  $L : P_4 \rightarrow \mathbb{R}^3$  given by  $L(p) = [p(-1), p(0), p(1)]$  is an isomorphism.

**Sol.** Basis for  $\ker(L)$  is  $\{x(x^2 - 1), x^2(x^2 - 1)\}$ . Hence,  $\dim(\ker(L)) = 2$ .



### Example 3

**Q.:** Check if the LT  $L : P_4 \rightarrow \mathbb{R}^3$  given by  $L(p) = [p(-1), p(0), p(1)]$  is an isomorphism.

**Sol.** Basis for  $\ker(L)$  is  $\{x(x^2 - 1), x^2(x^2 - 1)\}$ . Hence,  $\dim(\ker(L)) = 2$ .

Hence,  $L$  is not one-to-one, i.e., it is not an isomorphism.





## Exercise

**Q:.** Show that the linear operator  $L : P_2 \rightarrow P_2$  given by  $L(a + bt + ct^2) = (b + c) + (a + c)t + (a + b)t^2$  is an isomorphism.



## Exercise

**Q:.** Show that the linear operator  $L : P_2 \rightarrow P_2$  given by  $L(a + bt + ct^2) = (b + c) + (a + c)t + (a + b)t^2$  is an isomorphism.

**Q:.** Show that the linear operator  $L : M_{mn} \rightarrow M_{nm}$  given by  $L(A) = A^T$  is an isomorphism.



## Invertible LT

Let  $L : V \rightarrow W$  be a LT. Then  $L$  is an **invertible LT** if and only if



## Invertible LT

Let  $L : V \rightarrow W$  be a LT. Then  $L$  is an **invertible LT** if and only if there is a function  $M : W \rightarrow V$  such that  $(M \circ L)(v) = v$ , for all  $v \in V$ , and  $(L \circ M)(w) = w$ , for all  $w \in W$ .



## Invertible LT

Let  $L : V \rightarrow W$  be a LT. Then  $L$  is an **invertible LT** if and only if there is a function  $M : W \rightarrow V$  such that  $(M \circ L)(v) = v$ , for all  $v \in V$ , and  $(L \circ M)(w) = w$ , for all  $w \in W$ .

Such a function  $M$ , **denoted by  $L^{-1}$** , is called an **inverse** of  $L$ .



**Theorem:** A LT  $L : V \rightarrow W$  is an isomorphism if and only if  $L$  is an invertible LT.



**Theorem:** A LT  $L : V \rightarrow W$  is an isomorphism if and only if  $L$  is an invertible LT. Moreover, if  $L$  is invertible, then  $L^{-1}$  is also a LT.



## Example 4

**Q:.** Let  $L : \mathbb{R}^3 \rightarrow P_2$  be a LT given by  
$$L([x, y, z]) = x + (x + y - z)t + (x + y + z)t^2.$$





## Example 4

**Q:.** Let  $L : \mathbb{R}^3 \rightarrow P_2$  be a LT given by  
$$L([x, y, z]) = x + (x + y - z)t + (x + y + z)t^2.$$

Is  $L$  invertible.



## Example 4

**Q:.** Let  $L : \mathbb{R}^3 \rightarrow P_2$  be a LT given by  
$$L([x, y, z]) = x + (x + y - z)t + (x + y + z)t^2.$$

Is  $L$  invertible. If yes, find  $L^{-1}$ .



## Example 4

**Q:.** Let  $L : \mathbb{R}^3 \rightarrow P_2$  be a LT given by  
$$L([x, y, z]) = x + (x + y - z)t + (x + y + z)t^2.$$

Is  $L$  invertible. If yes, find  $L^{-1}$ .

**Sol.** It can be easily verified that  $L$  is both one-to-one and onto. Hence, invertible.



Let  $L^{-1} : P_2 \rightarrow \mathbb{R}^3$  be defined by



Let  $L^{-1} : P_2 \rightarrow \mathbb{R}^3$  be defined by  
 $L^{-1}(a + bt + ct^2) = [x, y, z]$



Let  $L^{-1} : P_2 \rightarrow \mathbb{R}^3$  be defined by

$$L^{-1}(a + bt + ct^2) = [x, y, z]$$

$$\implies L([x, y, z]) = a + bt + ct^2$$



Let  $L^{-1} : P_2 \rightarrow \mathbb{R}^3$  be defined by

$$L^{-1}(a + bt + ct^2) = [x, y, z]$$

$$\implies L([x, y, z]) = a + bt + ct^2$$

$$\implies x + (x + y - z)t + (x + y + z)t^2 = a + bt + ct^2$$



Let  $L^{-1} : P_2 \rightarrow \mathbb{R}^3$  be defined by

$$L^{-1}(a + bt + ct^2) = [x, y, z]$$

$$\implies L([x, y, z]) = a + bt + ct^2$$

$$\implies x + (x + y - z)t + (x + y + z)t^2 = a + bt + ct^2$$

$$\implies x = a, x + y - z = b, x + y + z = c$$





Let  $L^{-1} : P_2 \rightarrow \mathbb{R}^3$  be defined by

$$L^{-1}(a + bt + ct^2) = [x, y, z]$$

$$\implies L([x, y, z]) = a + bt + ct^2$$

$$\implies x + (x + y - z)t + (x + y + z)t^2 = a + bt + ct^2$$

$$\implies x = a, x + y - z = b, x + y + z = c$$

$$\implies x = a, y = \frac{b+c-2a}{2}, z = \frac{c-b}{2}.$$



Let  $L^{-1} : P_2 \rightarrow \mathbb{R}^3$  be defined by

$$L^{-1}(a + bt + ct^2) = [x, y, z]$$

$$\implies L([x, y, z]) = a + bt + ct^2$$

$$\implies x + (x + y - z)t + (x + y + z)t^2 = a + bt + ct^2$$

$$\implies x = a, x + y - z = b, x + y + z = c$$

$$\implies x = a, y = \frac{b+c-2a}{2}, z = \frac{c-b}{2}.$$

$$\text{Hence, } L^{-1}(a + bt + ct^2) = [a, \frac{b+c-2a}{2}, \frac{c-b}{2}].$$



## Example 5

**Q:.** Let  $L : P_2 \rightarrow P_2$  be a LT given by  
$$L(a + bt + ct^2) = (b + c) + (a + c)t + (a + b)t^2.$$



## Example 5

**Q:.** Let  $L : P_2 \rightarrow P_2$  be a LT given by  
$$L(a + bt + ct^2) = (b + c) + (a + c)t + (a + b)t^2.$$

Is  $L$  invertible.



## Example 5

**Q:.** Let  $L : P_2 \rightarrow P_2$  be a LT given by  
$$L(a + bt + ct^2) = (b + c) + (a + c)t + (a + b)t^2.$$

Is  $L$  invertible. If yes, find  $L^{-1}$ .



## Example 5

**Q:.** Let  $L : P_2 \rightarrow P_2$  be a LT given by  
$$L(a + bt + ct^2) = (b + c) + (a + c)t + (a + b)t^2.$$

Is  $L$  invertible. If yes, find  $L^{-1}$ .

**Sol.** It can be easily verified that  $L$  is both one-to-one and onto. Hence, invertible.



Let  $L^{-1} : P_2 \rightarrow P_2$  be defined by



Let  $L^{-1} : P_2 \rightarrow P_2$  be defined by  
$$L^{-1}(a + bt + ct^2) = a_1 + b_1t + c_1t^2$$





Let  $L^{-1} : P_2 \rightarrow P_2$  be defined by

$$L^{-1}(a + bt + ct^2) = a_1 + b_1t + c_1t^2$$
$$\implies L(a_1 + b_1t + c_1t^2) = a + bt + ct^2$$


Let  $L^{-1} : P_2 \rightarrow P_2$  be defined by

$$L^{-1}(a + bt + ct^2) = a_1 + b_1t + c_1t^2$$

$$\implies L(a_1 + b_1t + c_1t^2) = a + bt + ct^2$$

$$\implies (b_1 + c_1) + (a_1 + c_1)t + (a_1 + b_1)t^2 = a + bt + ct^2.$$



Let  $L^{-1} : P_2 \rightarrow P_2$  be defined by

$$L^{-1}(a + bt + ct^2) = a_1 + b_1t + c_1t^2$$

$$\implies L(a_1 + b_1t + c_1t^2) = a + bt + ct^2$$

$$\implies (b_1 + c_1) + (a_1 + c_1)t + (a_1 + b_1)t^2 = a + bt + ct^2.$$

On equating the coefficient of  $t$  on both sides, we get



Let  $L^{-1} : P_2 \rightarrow P_2$  be defined by

$$L^{-1}(a + bt + ct^2) = a_1 + b_1t + c_1t^2$$

$$\implies L(a_1 + b_1t + c_1t^2) = a + bt + ct^2$$

$$\implies (b_1 + c_1) + (a_1 + c_1)t + (a_1 + b_1)t^2 = a + bt + ct^2.$$

On equating the coefficient of  $t$  on both sides, we get

$$a = b_1 + c_1, b = a_1 + c_1, c = a_1 + b_1.$$



Let  $L^{-1} : P_2 \rightarrow P_2$  be defined by

$$L^{-1}(a + bt + ct^2) = a_1 + b_1t + c_1t^2$$

$$\implies L(a_1 + b_1t + c_1t^2) = a + bt + ct^2$$

$$\implies (b_1 + c_1) + (a_1 + c_1)t + (a_1 + b_1)t^2 = a + bt + ct^2.$$

On equating the coefficient of  $t$  on both sides, we get

$$a = b_1 + c_1, b = a_1 + c_1, c = a_1 + b_1.$$

On solving, we get

$$a_1 = \frac{1}{2}(b + c - a), b_1 = \frac{1}{2}(a + c - b), c_1 = \frac{1}{2}(a + b - c).$$



Let  $L^{-1} : P_2 \rightarrow P_2$  be defined by

$$L^{-1}(a + bt + ct^2) = a_1 + b_1t + c_1t^2$$

$$\implies L(a_1 + b_1t + c_1t^2) = a + bt + ct^2$$

$$\implies (b_1 + c_1) + (a_1 + c_1)t + (a_1 + b_1)t^2 = a + bt + ct^2.$$

On equating the coefficient of  $t$  on both sides, we get

$$a = b_1 + c_1, b = a_1 + c_1, c = a_1 + b_1.$$

On solving, we get

$$a_1 = \frac{1}{2}(b + c - a), b_1 = \frac{1}{2}(a + c - b), c_1 = \frac{1}{2}(a + b - c).$$

Hence,

$$L^{-1}(a + bt + ct^2) = \frac{1}{2}(b + c - a) + \frac{1}{2}(a + c - b)t + \frac{1}{2}(a + b - c)t^2$$



## Example 6

**Q:.** Let  $L : P_2 \rightarrow \mathbb{R}^2$  be a LT given by  $L(p) = [p(1), p'(1)]$ .



## Example 6

**Q:.** Let  $L : P_2 \rightarrow \mathbb{R}^2$  be a LT given by  $L(p) = [p(1), p'(1)]$ .

Is  $L$  invertible.





## Example 6

**Q:.** Let  $L : P_2 \rightarrow \mathbb{R}^2$  be a LT given by  $L(p) = [p(1), p'(1)]$ .

Is  $L$  invertible. If yes, find  $L^{-1}$ .



## Example 6

**Q:.** Let  $L : P_2 \rightarrow \mathbb{R}^2$  be a LT given by  $L(p) = [p(1), p'(1)]$ .

Is  $L$  invertible. If yes, find  $L^{-1}$ . If no, find  $\ker(L)$ .



## Example 6

**Q:.** Let  $L : P_2 \rightarrow \mathbb{R}^2$  be a LT given by  $L(p) = [p(1), p'(1)]$ .

Is  $L$  invertible. If yes, find  $L^{-1}$ . If no, find  $\ker(L)$ .

**Sol.**  $\ker(L) = \{p \in P_2 | L(p) = 0_{\mathbb{R}^2}\} \implies$



## Example 6

**Q.:** Let  $L : P_2 \rightarrow \mathbb{R}^2$  be a LT given by  $L(p) = [p(1), p'(1)]$ .

Is  $L$  invertible. If yes, find  $L^{-1}$ . If no, find  $\ker(L)$ .

**Sol.**  $\ker(L) = \{p \in P_2 | L(p) = 0_{\mathbb{R}^2}\} \implies$

$\ker(L) = \{p \in P_2 | [p(1), p'(1)] = [0, 0]\} \implies$



## Example 6

**Q:.** Let  $L : P_2 \rightarrow \mathbb{R}^2$  be a LT given by  $L(p) = [p(1), p'(1)]$ .

Is  $L$  invertible. If yes, find  $L^{-1}$ . If no, find  $\ker(L)$ .

**Sol.**  $\ker(L) = \{p \in P_2 | L(p) = 0_{\mathbb{R}^2}\} \implies$

$\ker(L) = \{p \in P_2 | [p(1), p'(1)] = [0, 0]\} \implies$

$\ker(L) = \{p \in P_2 | p(1) = 0, p'(1) = 0\}.$



Now,  $p(1) = 0 \implies$



Now,  $p(1) = 0 \implies p = (x - 1)(ax + b)$ .



Now,  $p(1) = 0 \implies p = (x - 1)(ax + b)$ . Also,  
 $p'(1) = 0 \implies$





Now,  $p(1) = 0 \implies p = (x - 1)(ax + b)$ . Also,  
 $p'(1) = 0 \implies a = -b$ .



Now,  $p(1) = 0 \implies p = (x-1)(ax+b)$ . Also,  
 $p'(1) = 0 \implies a = -b$ . Hence,  
 $p = (x-1)(ax-a) \implies p = a(x^2 - 2x + 1)$ .



Now,  $p(1) = 0 \implies p = (x-1)(ax+b)$ . Also,  
 $p'(1) = 0 \implies a = -b$ . Hence,  
 $p = (x-1)(ax-a) \implies p = a(x^2 - 2x + 1)$ . Thus  
 $\ker(L) = \{a(x^2 - 2x + 1) | a \in \mathbb{R}\}$ .



## Exercise

**Q.:** Let  $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  be a LT given by  
 $L(e_1) = e_1 + e_2, L(e_2) = e_2 + e_3, L(e_3) = e_1 + e_2 + e_3.$



## Exercise

**Q.:** Let  $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  be a LT given by  
 $L(e_1) = e_1 + e_2, L(e_2) = e_2 + e_3, L(e_3) = e_1 + e_2 + e_3.$

Is  $L$  invertible.



## Exercise

**Q.:** Let  $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  be a LT given by  
 $L(e_1) = e_1 + e_2, L(e_2) = e_2 + e_3, L(e_3) = e_1 + e_2 + e_3$ .  
Is  $L$  invertible. If yes, find  $L^{-1}$ .



## Exercise

**Q.:** Let  $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  be a LT given by  
 $L(e_1) = e_1 + e_2, L(e_2) = e_2 + e_3, L(e_3) = e_1 + e_2 + e_3$ .

Is  $L$  invertible. If yes, find  $L^{-1}$ .

**Sol.**  $L^{-1}([x, y, z]) = [y - z, y - x, x - y + z]$ .



## Isomorphism

Let  $V$  and  $W$  be vector spaces. Then  $V$  is isomorphic to  $W$ , denoted by  $V \cong W$ , if and only if there exists an isomorphism  $L : V \rightarrow W$ .





**Theorem:** Suppose  $V \cong W$  and  $V$  and  $W$  are finite dimensional. Then  $V$  is isomorphic to  $W$  if and only if  $\dim(V) = \dim(W)$ .



**Theorem:** Suppose  $V \cong W$  and  $V$  and  $W$  are finite dimensional. Then  $V$  is isomorphic to  $W$  if and only if  $\dim(V) = \dim(W)$ .

**Q:.** Show that  $\mathbb{R}^n$  and  $P_n$  are not isomorphic.



**Theorem:** Suppose  $V \cong W$  and  $V$  and  $W$  are finite dimensional. Then  $V$  is isomorphic to  $W$  if and only if  $\dim(V) = \dim(W)$ .

**Q:.** Show that  $\mathbb{R}^n$  and  $P_n$  are not isomorphic.

**Sol.** Since,  $\dim(\mathbb{R}^n) = n \neq n + 1 = \dim(P_n)$ ,



**Theorem:** Suppose  $V \cong W$  and  $V$  and  $W$  are finite dimensional. Then  $V$  is isomorphic to  $W$  if and only if  $\dim(V) = \dim(W)$ .

**Q:.** Show that  $\mathbb{R}^n$  and  $P_n$  are not isomorphic.

**Sol.** Since,  $\dim(\mathbb{R}^n) = n \neq n+1 = \dim(P_n)$ ,  $\mathbb{R}^n$  and  $P_n$  are not isomorphic.



## Exercise

**Q:.** Let  $W$  be the vector space of all symmetric  $2 \times 2$  matrices. Show that  $W$  is isomorphic to  $\mathbb{R}^3$ .



## Exercise

**Q:.** Let  $W$  be the vector space of all symmetric  $2 \times 2$  matrices. Show that  $W$  is isomorphic to  $\mathbb{R}^3$ .

**Q:.** Check if  $P_{4n+3} \cong M_{4,n+1}$ .

