

# MATHEMATICS-II (MATH F112)

**Dr. Krishnendra Shekhawat**

**BITS PILANI**  
**Department of Mathematics**



# Section 4.7

## *Coordinatization*



# Ordered Basis



## Ordered Basis

An **ordered basis** for vector space  $V$  is an ordered  $n$ -tuple of vectors  $(v_1, v_2, \dots, v_n)$  such that the set  $\{v_1, v_2, \dots, v_n\}$  is a basis for  $V$ .



## Ordered Basis

An **ordered basis** for vector space  $V$  is an ordered  $n$ -tuple of vectors  $(v_1, v_2, \dots, v_n)$  such that the set  $\{v_1, v_2, \dots, v_n\}$  is a basis for  $V$ .

For example,



## Ordered Basis

An **ordered basis** for vector space  $V$  is an ordered  $n$ -tuple of vectors  $(v_1, v_2, \dots, v_n)$  such that the set  $\{v_1, v_2, \dots, v_n\}$  is a basis for  $V$ .

For example,  $(e_1, e_2)$  and  $(e_2, e_1)$  are two ordered bases for  $\mathbb{R}^2$ .



# Coordinationization



## Coordinatization

Let  $B = (v_1, v_2, \dots, v_n)$  be an ordered basis for a vector space  $V$ .





## Coordinationization

Let  $B = (v_1, v_2, \dots, v_n)$  be an ordered basis for a vector space  $V$ . Suppose that  $w = a_1v_1 + \dots + a_nv_n \in V$



## Coordinationization

Let  $B = (v_1, v_2, \dots, v_n)$  be an ordered basis for a vector space  $V$ . Suppose that  $w = a_1v_1 + \dots + a_nv_n \in V$  (from Theorem 4.9, if  $B = (v_1, v_2, \dots, v_n)$  is an ordered basis for  $V$ , then for every  $w \in V$ , there are unique scalars  $a_1, a_2, \dots, a_n$  such that  $w = a_1v_1 + \dots + a_nv_n$ ).



## Coordinatization

Let  $B = (v_1, v_2, \dots, v_n)$  be an ordered basis for a vector space  $V$ . Suppose that  $w = a_1v_1 + \dots + a_nv_n \in V$  (from Theorem 4.9, if  $B = (v_1, v_2, \dots, v_n)$  is an ordered basis for  $V$ , then for every  $w \in V$ , there are unique scalars  $a_1, a_2, \dots, a_n$  such that  $w = a_1v_1 + \dots + a_nv_n$ ). Then  $[w]_B$ , the **coordinatization or coordinates of  $w$  with respect to  $B$**  is the  $n$ -vector  $[a_1, a_2, \dots, a_n]$ .



## Example 1

Let  $B = ([4, 2], [1, 3])$  be an ordered basis for  $\mathbb{R}^2$ .



## Example 1

Let  $B = ([4, 2], [1, 3])$  be an ordered basis for  $\mathbb{R}^2$ . Now,  $[4, 2] = 1[4, 2] + 0[1, 3]$ .



## Example 1

Let  $B = ([4, 2], [1, 3])$  be an ordered basis for  $\mathbb{R}^2$ . Now,  $[4, 2] = 1[4, 2] + 0[1, 3]$ . Hence,  $[4, 2]_B = [1, 0]$ .



## Example 1

Let  $B = ([4, 2], [1, 3])$  be an ordered basis for  $\mathbb{R}^2$ . Now,  $[4, 2] = 1[4, 2] + 0[1, 3]$ . Hence,  $[4, 2]_B = [1, 0]$ .

Similarly,  $[11, 13] = 2[4, 2] + 3[1, 3]$ . Hence,  $[11, 13]_B = [2, 3]$ .



## Example 2

Q: Let

$B = ([-4, 5, -1, 0, -1], [1, -3, 2, 2, 5], [1, -2, 1, 1, 3])$  be an ordered basis of the subspace  $V$  of  $\mathbb{R}^5$ .





## Example 2

**Q:.** Let

$B = ([-4, 5, -1, 0, -1], [1, -3, 2, 2, 5], [1, -2, 1, 1, 3])$  be an ordered basis of the subspace  $V$  of  $\mathbb{R}^5$ . Compute  $[-23, 30, -7, -1, -7]_B, [1, 2, 3, 4, 5]_B$ .



## Example 2

**Q:.** Let

$B = ([-4, 5, -1, 0, -1], [1, -3, 2, 2, 5], [1, -2, 1, 1, 3])$  be an ordered basis of the subspace  $V$  of  $\mathbb{R}^5$ . Compute  $[-23, 30, -7, -1, -7]_B, [1, 2, 3, 4, 5]_B$ .

**Sol.** To find  $[-23, 30, -7, -1, -7]_B$ ,



## Example 2

**Q:.** Let

$B = ([-4, 5, -1, 0, -1], [1, -3, 2, 2, 5], [1, -2, 1, 1, 3])$  be an ordered basis of the subspace  $V$  of  $\mathbb{R}^5$ . Compute  $[-23, 30, -7, -1, -7]_B, [1, 2, 3, 4, 5]_B$ .

**Sol.** To find  $[-23, 30, -7, -1, -7]_B$ , we solve the following equation

$$[-23, 30, -7, -1, -7] = a[-4, 5, -1, 0, -1] + b[1, -3, 2, 2, 5] + c[1, -2, 1, 1, 3]$$



The equivalent system is



The equivalent system is

$$-4a + b + c = -23$$

$$5a - 3b - 2c = 30$$

$$-a + 2b + c = -7$$

$$2b + c = -1$$

$$-a + 5b + 3c = -7$$



The equivalent system is

$$-4a + b + c = -23$$

$$5a - 3b - 2c = 30$$

$$-a + 2b + c = -7$$

$$2b + c = -1$$

$$-a + 5b + 3c = -7$$

To solve this system, we row reduce



$$\left[ \begin{array}{ccc|c} -4 & 1 & 1 & -23 \\ 5 & -3 & -2 & 30 \\ -1 & 2 & 1 & -7 \\ 0 & 2 & 1 & -1 \\ -1 & 5 & 3 & -7 \end{array} \right]$$



$$\left[ \begin{array}{ccc|c} -4 & 1 & 1 & -23 \\ 5 & -3 & -2 & 30 \\ -1 & 2 & 1 & -7 \\ 0 & 2 & 1 & -1 \\ -1 & 5 & 3 & -7 \end{array} \right] \text{ to obtain}$$





$$\left[ \begin{array}{ccc|c} -4 & 1 & 1 & -23 \\ 5 & -3 & -2 & 30 \\ -1 & 2 & 1 & -7 \\ 0 & 2 & 1 & -1 \\ -1 & 5 & 3 & -7 \end{array} \right] \text{ to obtain } \left[ \begin{array}{ccc|c} 1 & 0 & 0 & 6 \\ 0 & 1 & 0 & -2 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$



$$\left[ \begin{array}{ccc|c} -4 & 1 & 1 & -23 \\ 5 & -3 & -2 & 30 \\ -1 & 2 & 1 & -7 \\ 0 & 2 & 1 & -1 \\ -1 & 5 & 3 & -7 \end{array} \right] \text{ to obtain } \left[ \begin{array}{ccc|c} 1 & 0 & 0 & 6 \\ 0 & 1 & 0 & -2 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Hence, the unique solution for the system is



$$\left[ \begin{array}{ccc|c} -4 & 1 & 1 & -23 \\ 5 & -3 & -2 & 30 \\ -1 & 2 & 1 & -7 \\ 0 & 2 & 1 & -1 \\ -1 & 5 & 3 & -7 \end{array} \right] \text{ to obtain } \left[ \begin{array}{ccc|c} 1 & 0 & 0 & 6 \\ 0 & 1 & 0 & -2 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Hence, the unique solution for the system is  
 $a = 6, b = -2, c = 3 \implies$



$$\left[ \begin{array}{ccc|c} -4 & 1 & 1 & -23 \\ 5 & -3 & -2 & 30 \\ -1 & 2 & 1 & -7 \\ 0 & 2 & 1 & -1 \\ -1 & 5 & 3 & -7 \end{array} \right] \text{ to obtain } \left[ \begin{array}{ccc|c} 1 & 0 & 0 & 6 \\ 0 & 1 & 0 & -2 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Hence, the unique solution for the system is

$$a = 6, b = -2, c = 3 \implies$$

$$[-23, 30, -7, -1, -7]_B = [6, -2, 3].$$



To find  $[1,2,3,4,5]_B$ , we need solve the following system



To find  $[1,2,3,4,5]_B$ , we need solve the following system

$$-4a + b + c = 1$$

$$5a - 3b - 2c = 2$$

$$-a + 2b + c = 3$$

$$2b + c = 4$$

$$-a + 5b + 3c = 5$$



To find  $[1,2,3,4,5]_B$ , we need solve the following system

$$-4a + b + c = 1$$

$$5a - 3b - 2c = 2$$

$$-a + 2b + c = 3$$

$$2b + c = 4$$

$$-a + 5b + 3c = 5$$

To solve this system, we row reduce



$$\left[ \begin{array}{ccc|c} -4 & 1 & 1 & 1 \\ 5 & -3 & -2 & 2 \\ -1 & 2 & 1 & 3 \\ 0 & 2 & 1 & 4 \\ -1 & 5 & 3 & 5 \end{array} \right]$$





$$\left[ \begin{array}{ccc|c} -4 & 1 & 1 & 1 \\ 5 & -3 & -2 & 2 \\ -1 & 2 & 1 & 3 \\ 0 & 2 & 1 & 4 \\ -1 & 5 & 3 & 5 \end{array} \right] \text{ to obtain}$$



$$\left[ \begin{array}{ccc|c} -4 & 1 & 1 & 1 \\ 5 & -3 & -2 & 2 \\ -1 & 2 & 1 & 3 \\ 0 & 2 & 1 & 4 \\ -1 & 5 & 3 & 5 \end{array} \right] \text{ to obtain } \left[ \begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{array} \right]$$



$$\left[ \begin{array}{ccc|c} -4 & 1 & 1 & 1 \\ 5 & -3 & -2 & 2 \\ -1 & 2 & 1 & 3 \\ 0 & 2 & 1 & 4 \\ -1 & 5 & 3 & 5 \end{array} \right] \text{ to obtain } \left[ \begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

This system has no solutions,



$$\left[ \begin{array}{ccc|c} -4 & 1 & 1 & 1 \\ 5 & -3 & -2 & 2 \\ -1 & 2 & 1 & 3 \\ 0 & 2 & 1 & 4 \\ -1 & 5 & 3 & 5 \end{array} \right] \text{ to obtain } \left[ \begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

This system has no solutions, implying the vector  $[1, 2, 3, 4, 5]$  is not in  $\text{span}(\mathbf{B}) = V$ .



# Coordinatization Method



## Coordination Method

Let  $V$  be a nontrivial subspace of  $\mathbb{R}^n$ , let

$B = (v_1, v_2, \dots, v_k)$  be an ordered basis for  $V$ , and let  $v \in \mathbb{R}^n$ .



## Coordinatization Method

Let  $V$  be a nontrivial subspace of  $\mathbb{R}^n$ , let

$B = (v_1, v_2, \dots, v_k)$  be an ordered basis for  $V$ , and let  $v \in \mathbb{R}^n$ . To compute  $[v]_B$ , we perform the following steps:



## Coordinatization Method

Let  $V$  be a nontrivial subspace of  $\mathbb{R}^n$ , let

$B = (v_1, v_2, \dots, v_k)$  be an ordered basis for  $V$ , and let  $v \in \mathbb{R}^n$ . To compute  $[v]_B$ , we perform the following steps:

- Form an augmented matrix  $[A|v]$





## Coordination Method

Let  $V$  be a nontrivial subspace of  $\mathbb{R}^n$ , let

$B = (v_1, v_2, \dots, v_k)$  be an ordered basis for  $V$ , and let  $v \in \mathbb{R}^n$ . To compute  $[v]_B$ , we perform the following steps:

- Form an augmented matrix  $[A|v]$  by using the vectors in  $B$  as the columns of  $A$ , in order,



## Coordination Method

Let  $V$  be a nontrivial subspace of  $\mathbb{R}^n$ , let

$B = (v_1, v_2, \dots, v_k)$  be an ordered basis for  $V$ , and let  $v \in \mathbb{R}^n$ . To compute  $[v]_B$ , we perform the following steps:

- Form an augmented matrix  $[A|v]$  by using the vectors in  $B$  as the columns of  $A$ , in order, and using  $v$  as a column on the right.



## Coordination Method

Let  $V$  be a nontrivial subspace of  $\mathbb{R}^n$ , let

$B = (v_1, v_2, \dots, v_k)$  be an ordered basis for  $V$ , and let  $v \in \mathbb{R}^n$ . To compute  $[v]_B$ , we perform the following steps:

- Form an augmented matrix  $[A|v]$  by using the vectors in  $B$  as the columns of  $A$ , in order, and using  $v$  as a column on the right.
- Row reduce  $[A|v]$  to obtain  $\text{RREF}[C|w]$ .



- If there is a row of  $[C|w]$  that contains all zeros on the left and has a nonzero entry on the right,



- If there is a row of  $[C|w]$  that contains all zeros on the left and has a nonzero entry on the right, then  $v \notin \text{span}(B) = V$ ,



- If there is a row of  $[C|w]$  that contains all zeros on the left and has a nonzero entry on the right, then  $v \notin \text{span}(B) = V$ , i.e., coordinatization is not possible.



- If there is a row of  $[C|w]$  that contains all zeros on the left and has a nonzero entry on the right, then  $v \notin \text{span}(B) = V$ , i.e., coordinatization is not possible. Otherwise,  $v \in \text{span}(B) = V$ .



- If there is a row of  $[C|w]$  that contains all zeros on the left and has a nonzero entry on the right, then  $v \notin \text{span}(B) = V$ , i.e., coordinatization is not possible. Otherwise,  $v \in \text{span}(B) = V$ .
- Eliminate all rows consisting entirely of zeros in  $[C|w]$  to obtain  $[I_k|y]$ .





- If there is a row of  $[C|w]$  that contains all zeros on the left and has a nonzero entry on the right, then  $v \notin \text{span}(B) = V$ , i.e., coordinatization is not possible. Otherwise,  $v \in \text{span}(B) = V$ .
- Eliminate all rows consisting entirely of zeros in  $[C|w]$  to obtain  $[I_k|y]$ . Then,  $[v]_B = y$ ,



- If there is a row of  $[C|w]$  that contains all zeros on the left and has a nonzero entry on the right, then  $v \notin \text{span}(B) = V$ , i.e., coordinatization is not possible. Otherwise,  $v \in \text{span}(B) = V$ .
- Eliminate all rows consisting entirely of zeros in  $[C|w]$  to obtain  $[I_k|y]$ . Then,  $[v]_B = y$ , the last column of  $[I_k|y]$ .



### Example 3

**Q:.** Let  $B = \left( \begin{bmatrix} 1 & -2 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 2 & -1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 1 & -1 \\ 3 & 1 \end{bmatrix} \right)$  be an ordered basis of the subspace  $V$  of  $M_{22}$ .



### Example 3

**Q:.** Let  $B = \left( \begin{bmatrix} 1 & -2 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 2 & -1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 1 & -1 \\ 3 & 1 \end{bmatrix} \right)$  be an ordered basis of the subspace  $V$  of  $M_{22}$ . Compute  $[v]_B$  if exists, where  $v = \begin{bmatrix} -3 & -2 \\ 0 & 3 \end{bmatrix}$ .



### Example 3

**Q:.** Let  $B = \left( \begin{bmatrix} 1 & -2 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 2 & -1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 1 & -1 \\ 3 & 1 \end{bmatrix} \right)$  be an ordered basis of the subspace  $V$  of  $M_{22}$ . Compute  $[v]_B$  if exists, where  $v = \begin{bmatrix} -3 & -2 \\ 0 & 3 \end{bmatrix}$ .

**Sol.** Now

$$[A|v] = \left[ \begin{array}{ccc|c} 1 & 2 & 1 & -3 \\ -2 & -1 & -1 & -2 \\ 0 & 1 & 3 & 0 \\ 1 & 0 & 1 & 3 \end{array} \right] \Rightarrow$$



$$\text{RREF}[A|v] = \left[ \begin{array}{ccc|c} 1 & 0 & 0 & 2 \\ 0 & 1 & 0 & -3 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{array} \right]$$



$$\text{RREF}[A|v] = \left[ \begin{array}{ccc|c} 1 & 0 & 0 & 2 \\ 0 & 1 & 0 & -3 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

The row reduced matrix contains no rows with all zero entries on the left and a nonzero entry on the right, so  $[v]_B$  exists,



$$\text{RREF}[A|v] = \left[ \begin{array}{ccc|c} 1 & 0 & 0 & 2 \\ 0 & 1 & 0 & -3 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

The row reduced matrix contains no rows with all zero entries on the left and a nonzero entry on the right, so  $[v]_B$  exists, and  $[v]_B = [2, -3, 1]$ .





# Fundamental properties of Coordinatization



## Fundamental properties of Coordinatization

Let  $B = (v_1, v_2, \dots, v_k)$  be an ordered basis for a vector space  $V$ . Suppose  $w_1, w_2, \dots, w_k \in V$  and  $a_1, a_2, \dots, a_k$  are scalars. Then



## Fundamental properties of Coordinatization

Let  $B = (v_1, v_2, \dots, v_k)$  be an ordered basis for a vector space  $V$ . Suppose  $w_1, w_2, \dots, w_k \in V$  and  $a_1, a_2, \dots, a_k$  are scalars. Then

- $[w_1 + w_2]_B = [w_1]_B + [w_2]_B$



## Fundamental properties of Coordinatization

Let  $B = (v_1, v_2, \dots, v_k)$  be an ordered basis for a vector space  $V$ . Suppose  $w_1, w_2, \dots, w_k \in V$  and  $a_1, a_2, \dots, a_k$  are scalars. Then

- $[w_1 + w_2]_B = [w_1]_B + [w_2]_B$
- $[a_1 w_1]_B = a_1 [w_1]_B$



## Fundamental properties of Coordinatization

Let  $B = (v_1, v_2, \dots, v_k)$  be an ordered basis for a vector space  $V$ . Suppose  $w_1, w_2, \dots, w_k \in V$  and  $a_1, a_2, \dots, a_k$  are scalars. Then

- $[w_1 + w_2]_B = [w_1]_B + [w_2]_B$
- $[a_1 w_1]_B = a_1 [w_1]_B$
- $[a_1 w_1 + a_2 w_2 + \dots + a_k w_k]_B = a_1 [w_1]_B + a_2 [w_2]_B + \dots + a_k [w_k]_B$



## Exercises

**Q:.** Let  $B = (3x^2 - x + 2, x^2 + 2x - 3, 2x^2 + 3x - 1)$  be an ordered basis of the subspace  $V$  of  $P_2$ .



## Exercises

**Q:.** Let  $B = (3x^2 - x + 2, x^2 + 2x - 3, 2x^2 + 3x - 1)$  be an ordered basis of the subspace  $V$  of  $P_2$ . Compute  $[v]_B$  if exists, where  $v = 13x^2 - 5x + 20$ .



## Exercises

**Q:.** Let  $B = (3x^2 - x + 2, x^2 + 2x - 3, 2x^2 + 3x - 1)$  be an ordered basis of the subspace  $V$  of  $P_2$ . Compute  $[v]_B$  if exists, where  $v = 13x^2 - 5x + 20$ .

**Sol.**  $[v]_B = [4, -5, 3]$ .





## Exercises

**Q:.** Let  $B = (3x^2 - x + 2, x^2 + 2x - 3, 2x^2 + 3x - 1)$  be an ordered basis of the subspace  $V$  of  $P_2$ . Compute  $[v]_B$  if exists, where  $v = 13x^2 - 5x + 20$ .

**Sol.**  $[v]_B = [4, -5, 3]$ .

**Q:.** Let  $B = ([-4, 5, -1, 0, -1], [1, -3, 2, 2, 5], [1, -2, 1, 1, 3])$  be an ordered basis of the subspace  $V$  of  $\mathbb{R}^5$ .



## Exercises

**Q:.** Let  $B = (3x^2 - x + 2, x^2 + 2x - 3, 2x^2 + 3x - 1)$  be an ordered basis of the subspace  $V$  of  $P_2$ . Compute  $[v]_B$  if exists, where  $v = 13x^2 - 5x + 20$ .

**Sol.**  $[v]_B = [4, -5, 3]$ .

**Q:.** Let  $B = ([-4, 5, -1, 0, -1], [1, -3, 2, 2, 5], [1, -2, 1, 1, 3])$  be an ordered basis of the subspace  $V$  of  $\mathbb{R}^5$ . Consider  $x = [1, 0, -1, 0, 4], y = [0, 1, -1, 0, 3], z = [0, 0, 0, 1, 5]$ .



## Exercises

**Q:.** Let  $B = (3x^2 - x + 2, x^2 + 2x - 3, 2x^2 + 3x - 1)$  be an ordered basis of the subspace  $V$  of  $P_2$ . Compute  $[v]_B$  if exists, where  $v = 13x^2 - 5x + 20$ .

**Sol.**  $[v]_B = [4, -5, 3]$ .

**Q:.** Let  $B = ([-4, 5, -1, 0, -1], [1, -3, 2, 2, 5], [1, -2, 1, 1, 3])$  be an ordered basis of the subspace  $V$  of  $\mathbb{R}^5$ . Consider  $x = [1, 0, -1, 0, 4], y = [0, 1, -1, 0, 3], z = [0, 0, 0, 1, 5]$ . Compute  $[2x - 7y + 3z]_B$ .



## Exercises

**Q:.** Let  $B = (3x^2 - x + 2, x^2 + 2x - 3, 2x^2 + 3x - 1)$  be an ordered basis of the subspace  $V$  of  $P_2$ . Compute  $[v]_B$  if exists, where  $v = 13x^2 - 5x + 20$ .

**Sol.**  $[v]_B = [4, -5, 3]$ .

**Q:.** Let  $B = ([-4, 5, -1, 0, -1], [1, -3, 2, 2, 5], [1, -2, 1, 1, 3])$  be an ordered basis of the subspace  $V$  of  $\mathbb{R}^5$ . Consider  $x = [1, 0, -1, 0, 4], y = [0, 1, -1, 0, 3], z = [0, 0, 0, 1, 5]$ . Compute  $[2x - 7y + 3z]_B$ .

**Sol.**  $[2x - 7y + 3z]_B = [-2, 9, -15]$ .



## Example 4

**Q:.** Let

$C = ([-4, 5, -1, 0, -1], [1, -3, 2, 2, 5], [1, -2, 1, 1, 3])$  be an ordered basis of the subspace  $V$  of  $\mathbb{R}^5$ .



## Example 4

**Q:.** Let

$C = ([-4, 5, -1, 0, -1], [1, -3, 2, 2, 5], [1, -2, 1, 1, 3])$  be an ordered basis of the subspace  $V$  of  $\mathbb{R}^5$ . Using simplified span method on  $C$ , compute an ordered basis  $B = (x, y, z)$  for  $V$ .



## Example 4

**Q:.** Let

$C = ([-4, 5, -1, 0, -1], [1, -3, 2, 2, 5], [1, -2, 1, 1, 3])$  be an ordered basis of the subspace  $V$  of  $\mathbb{R}^5$ . Using simplified span method on  $C$ , compute an ordered basis  $B = (x, y, z)$  for  $V$ . Also, compute  $[x]_C, [y]_C, [z]_C$ .



## Example 4

**Q:.** Let

$C = ([-4, 5, -1, 0, -1], [1, -3, 2, 2, 5], [1, -2, 1, 1, 3])$  be an ordered basis of the subspace  $V$  of  $\mathbb{R}^5$ . Using simplified span method on  $C$ , compute an ordered basis  $B = (x, y, z)$  for  $V$ . Also, compute  $[x]_C, [y]_C, [z]_C$ .

**Sol.** Using simplified span method, we have  
 $B = ([1, 0, -1, 0, 4], [0, 1, -1, 0, 3], [0, 0, 0, 1, 5]).$





We have the following augmented matrix

$$\left[ A \mid x \ y \ z \right] = \left[ \begin{array}{ccc|ccc} -4 & 1 & 1 & 1 & 0 & 0 \\ 5 & -3 & -2 & 0 & 1 & 0 \\ -1 & 2 & 1 & -1 & -1 & 0 \\ 0 & 2 & 1 & 0 & 0 & 1 \\ -1 & 5 & 3 & 4 & 3 & 5 \end{array} \right]$$



We have the following augmented matrix

$$\left[ A \mid x \ y \ z \right] = \left[ \begin{array}{ccc|ccc} -4 & 1 & 1 & 1 & 0 & 0 \\ 5 & -3 & -2 & 0 & 1 & 0 \\ -1 & 2 & 1 & -1 & -1 & 0 \\ 0 & 2 & 1 & 0 & 0 & 1 \\ -1 & 5 & 3 & 4 & 3 & 5 \end{array} \right]$$

Row reduce above augmented matrix to obtain



We have the following augmented matrix

$$\left[ A \mid x \ y \ z \right] = \left[ \begin{array}{ccc|ccc} -4 & 1 & 1 & 1 & 0 & 0 \\ 5 & -3 & -2 & 0 & 1 & 0 \\ -1 & 2 & 1 & -1 & -1 & 0 \\ 0 & 2 & 1 & 0 & 0 & 1 \\ -1 & 5 & 3 & 4 & 3 & 5 \end{array} \right]$$

Row reduce above augmented matrix to obtain

$$\left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 1 & 1 \\ 0 & 1 & 0 & -5 & -4 & -3 \\ 0 & 0 & 1 & 10 & 8 & 7 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right]$$



Clearly,  $[x]_C = [1, -5, 10]$ ,  $[y]_C = [1, -4, 8]$  and  $[z]_C = [1, -3, 7]$ .



Clearly,  $[x]_C = [1, -5, 10]$ ,  $[y]_C = [1, -4, 8]$  and  $[z]_C = [1, -3, 7]$ .

$$\text{Here, } P = \begin{bmatrix} 1 & 1 & 1 \\ -5 & -4 & -3 \\ 10 & 8 & 7 \end{bmatrix}$$



Clearly,  $[x]_C = [1, -5, 10]$ ,  $[y]_C = [1, -4, 8]$  and  $[z]_C = [1, -3, 7]$ .

Here,  $P = \begin{bmatrix} 1 & 1 & 1 \\ -5 & -4 & -3 \\ 10 & 8 & 7 \end{bmatrix}$  is called **transition matrix** from  **$B$ -coordinates to  $C$ -coordinates**.



# Transition Matrix



## Transition Matrix

Suppose that  $V$  is a nontrivial  $n$ -dimensional vector space with ordered bases  $B$  and  $C$ .





## Transition Matrix

Suppose that  $V$  is a nontrivial  $n$ -dimensional vector space with ordered bases  $B$  and  $C$ . Let  $P$  be the  $n \times n$  matrix whose  $i^{\text{th}}$  column, for  $1 \leq i \leq n$ , equals  $[b_i]_C$ , where  $b_i$  is the  $i^{\text{th}}$  basis vector in  $B$ .



## Transition Matrix

Suppose that  $V$  is a nontrivial  $n$ -dimensional vector space with ordered bases  $B$  and  $C$ . Let  $P$  be the  $n \times n$  matrix whose  $i^{th}$  column, for  $1 \leq i \leq n$ , equals  $[b_i]_C$ , where  $b_i$  is the  $i^{th}$  basis vector in  $B$ . Then  $P$  is called the transition matrix from  $B$ -coordinates to  $C$ -coordinates



## Transition Matrix

Suppose that  $V$  is a nontrivial  $n$ -dimensional vector space with ordered bases  $B$  and  $C$ . Let  $P$  be the  $n \times n$  matrix whose  $i^{th}$  column, for  $1 \leq i \leq n$ , equals  $[b_i]_C$ , where  $b_i$  is the  $i^{th}$  basis vector in  $B$ . Then  $P$  is called the **transition matrix from  $B$ -coordinates to  $C$ -coordinates** (or transition matrix from  $B$  to  $C$ ).



# Transition Matrix Method



# Transition Matrix Method

To find the transition matrix  $P$  from  $B$  to  $C$ ,



## Transition Matrix Method

To find the transition matrix  $P$  from  $B$  to  $C$ , we use row reduction on

$$\left[ \begin{array}{ccc|ccc} 1^{st} & 2^{nd} & & k^{th} & 1^{st} & 2^{nd} & & k^{th} \\ vector & vector & \cdots & vector & vector & vector & \cdots & vector \\ in & in & & in & in & in & & in \\ C & C & & C & B & B & & B \end{array} \right]$$



## Transition Matrix Method

To find the transition matrix  $P$  from  $B$  to  $C$ , we use row reduction on

$$\left[ \begin{array}{ccc|ccc} 1^{st} & 2^{nd} & & k^{th} & 1^{st} & 2^{nd} & & k^{th} \\ vector & vector & \cdots & vector & vector & vector & \cdots & vector \\ in & in & & in & in & in & & in \\ C & C & & C & B & B & & B \end{array} \right]$$

to produce

$$\left[ \begin{array}{c|c} I_k & P \\ \hline rowsof & zeroes \end{array} \right]$$



## Example 5

Q:.. For the ordered bases  $B = \left( \begin{bmatrix} 7 & 3 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}, \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} \right)$





## Example 5

**Q:.** For the ordered bases  $B = \left( \begin{bmatrix} 7 & 3 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}, \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} \right)$   
and  $C = \left( \begin{bmatrix} 22 & 7 \\ 0 & 2 \end{bmatrix}, \begin{bmatrix} 12 & 4 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 33 & 12 \\ 0 & 2 \end{bmatrix} \right)$  of  $U_2$  (set of  $2 \times 2$   
upper triangular matrices),



## Example 5

**Q:.** For the ordered bases  $B = \left( \begin{bmatrix} 7 & 3 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}, \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} \right)$  and  $C = \left( \begin{bmatrix} 22 & 7 \\ 0 & 2 \end{bmatrix}, \begin{bmatrix} 12 & 4 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 33 & 12 \\ 0 & 2 \end{bmatrix} \right)$  of  $U_2$  (set of  $2 \times 2$  upper triangular matrices), find the transition matrix  $P$  from  $B$  to  $C$ .



## Example 5

**Q:.** For the ordered bases  $B = \left( \begin{bmatrix} 7 & 3 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}, \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} \right)$  and  $C = \left( \begin{bmatrix} 22 & 7 \\ 0 & 2 \end{bmatrix}, \begin{bmatrix} 12 & 4 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 33 & 12 \\ 0 & 2 \end{bmatrix} \right)$  of  $U_2$  (set of  $2 \times 2$  upper triangular matrices), find the transition matrix  $P$  from  $B$  to  $C$ .

**Sol.** Row reduce



## Example 5

**Q:.** For the ordered bases  $B = \left( \begin{bmatrix} 7 & 3 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}, \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} \right)$  and  $C = \left( \begin{bmatrix} 22 & 7 \\ 0 & 2 \end{bmatrix}, \begin{bmatrix} 12 & 4 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 33 & 12 \\ 0 & 2 \end{bmatrix} \right)$  of  $U_2$  (set of  $2 \times 2$  upper triangular matrices), find the transition matrix  $P$  from  $B$  to  $C$ .

**Sol.** Row reduce

$$\left[ \begin{array}{ccc|ccc} 22 & 12 & 33 & 7 & 1 & 1 \\ 7 & 4 & 12 & 3 & 2 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 2 & 1 & 2 & 0 & -1 & 1 \end{array} \right]$$



to obtain

$$\left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & -2 & 1 \\ 0 & 1 & 0 & -4 & 1 & 1 \\ 0 & 0 & 1 & 1 & 1 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right]$$



to obtain

$$\left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & -2 & 1 \\ 0 & 1 & 0 & -4 & 1 & 1 \\ 0 & 0 & 1 & 1 & 1 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

The transition matrix  $P$  from  $B$  to  $C$  is



to obtain

$$\left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & -2 & 1 \\ 0 & 1 & 0 & -4 & 1 & 1 \\ 0 & 0 & 1 & 1 & 1 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

The transition matrix  $P$  from  $B$  to  $C$  is

$$\begin{bmatrix} 1 & -2 & 1 \\ -4 & 1 & 1 \\ 1 & 1 & -1 \end{bmatrix}$$



## Example 6

**Q:.** Let  $C = (a, b, c) = ([1, 0, 1], [1, 1, 0], [0, 0, 1])$  and

$B = (x, y, z)$  are ordered bases of  $\mathbb{R}^3$ .





## Example 6

**Q:.** Let  $C = (a, b, c) = ([1, 0, 1], [1, 1, 0], [0, 0, 1])$  and

$B = (x, y, z)$  are ordered bases of  $\mathbb{R}^3$ . Let  $P = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 1 & 1 \\ -1 & -1 & 1 \end{bmatrix}$

be the transition matrix from  $B$  to  $C$ .



## Example 6

**Q:.** Let  $C = (a, b, c) = ([1, 0, 1], [1, 1, 0], [0, 0, 1])$  and

$B = (x, y, z)$  are ordered bases of  $\mathbb{R}^3$ . Let  $P = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 1 & 1 \\ -1 & -1 & 1 \end{bmatrix}$

be the transition matrix from  $B$  to  $C$ . Find the basis  $B$ .



## Example 6

**Q:.** Let  $C = (a, b, c) = ([1, 0, 1], [1, 1, 0], [0, 0, 1])$  and

$B = (x, y, z)$  are ordered bases of  $\mathbb{R}^3$ . Let  $P = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 1 & 1 \\ -1 & -1 & 1 \end{bmatrix}$

be the transition matrix from  $B$  to  $C$ . Find the basis  $B$ .

**Sol.**  $x = 1.a + 2.b - 1.c = [3, 2, 0]$ .



## Example 6

**Q.:** Let  $C = (a, b, c) = ([1, 0, 1], [1, 1, 0], [0, 0, 1])$  and

$B = (x, y, z)$  are ordered bases of  $\mathbb{R}^3$ . Let  $P = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 1 & 1 \\ -1 & -1 & 1 \end{bmatrix}$

be the transition matrix from  $B$  to  $C$ . Find the basis  $B$ .

**Sol.**  $x = 1.a + 2.b - 1.c = [3, 2, 0]$ .

Similarly,  $y = 1.a + 1.b - 1.c = [2, 1, 0]$  and

$z = 2.a + 1.b + 1.c = [3, 1, 3]$ .



## Example 6

**Q:.** Let  $C = (a, b, c) = ([1, 0, 1], [1, 1, 0], [0, 0, 1])$  and

$B = (x, y, z)$  are ordered bases of  $\mathbb{R}^3$ . Let  $P = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 1 & 1 \\ -1 & -1 & 1 \end{bmatrix}$

be the transition matrix from  $B$  to  $C$ . Find the basis  $B$ .

**Sol.**  $x = 1.a + 2.b - 1.c = [3, 2, 0]$ .

Similarly,  $y = 1.a + 1.b - 1.c = [2, 1, 0]$  and

$z = 2.a + 1.b + 1.c = [3, 1, 3]$ .

Hence,  $B = ([3, 2, 0], [2, 1, 0], [3, 1, 3])$ .



Example 6 can be solved by considering the augmented matrix  $[I_3|P]$  and reduce it to  $[C|B]$ ,



Example 6 can be solved by considering the augmented matrix  $[I_3|P]$  and reduce it to  $[C|B]$ , where  $B$  is the required basis.



Example 6 can be solved by considering the augmented matrix  $[I_3|P]$  and reduce it to  $[C|B]$ , where  $B$  is the required basis.

Also, it should be noted that





Example 6 can be solved by considering the augmented matrix  $[I_3|P]$  and reduce it to  $[C|B]$ , where  $B$  is the required basis.

Also, it should be noted that  $CP = B$ ,



Example 6 can be solved by considering the augmented matrix  $[I_3|P]$  and reduce it to  $[C|B]$ , where  $B$  is the required basis.

Also, it should be noted that  $CP = B$ , where matrix  $C$  is obtained by considering vectors of basis  $C$  as columns of the matrix.



# Change of Coordinates Using the Transition Matrix



## Change of Coordinates Using the Transition Matrix

**Theorem:** Suppose that  $V$  is a nontrivial  $n$ -dimensional vector space with ordered bases  $B$  and  $C$ .



## Change of Coordinates Using the Transition Matrix

**Theorem:** Suppose that  $V$  is a nontrivial  $n$ -dimensional vector space with ordered bases  $B$  and  $C$ . Let  $P$  be an  $n \times n$  matrix.



## Change of Coordinates Using the Transition Matrix

**Theorem:** Suppose that  $V$  is a nontrivial  $n$ -dimensional vector space with ordered bases  $B$  and  $C$ . Let  $P$  be an  $n \times n$  matrix. Then  $P$  is the transition matrix from  $B$  to  $C$  if and only if for every  $v \in V$ ,  $P[v]_B = [v]_C$ .



## Example 7

Q: For the ordered bases  $B = \left( \begin{bmatrix} 7 & 3 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}, \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} \right)$



## Example 7

**Q:.** For the ordered bases  $B = \left( \begin{bmatrix} 7 & 3 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}, \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} \right)$  and  $C = \left( \begin{bmatrix} 22 & 7 \\ 0 & 2 \end{bmatrix}, \begin{bmatrix} 12 & 4 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 33 & 12 \\ 0 & 2 \end{bmatrix} \right)$  of  $U_2$  (set of  $2 \times 2$  upper triangular matrices),





## Example 7

**Q.:** For the ordered bases  $B = \left( \begin{bmatrix} 7 & 3 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}, \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} \right)$  and  $C = \left( \begin{bmatrix} 22 & 7 \\ 0 & 2 \end{bmatrix}, \begin{bmatrix} 12 & 4 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 33 & 12 \\ 0 & 2 \end{bmatrix} \right)$  of  $U_2$  (set of  $2 \times 2$  upper triangular matrices), find  $[v]_B$  and  $[v]_C$  where  $v = \begin{bmatrix} 25 & 24 \\ 0 & -9 \end{bmatrix}$ .



**Sol.** Clearly,

$$\begin{bmatrix} 25 & 24 \\ 0 & -9 \end{bmatrix} = 4 \begin{bmatrix} 7 & 3 \\ 0 & 0 \end{bmatrix} + 3 \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix} - 6 \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}$$



**Sol.** Clearly,

$$\begin{bmatrix} 25 & 24 \\ 0 & -9 \end{bmatrix} = 4 \begin{bmatrix} 7 & 3 \\ 0 & 0 \end{bmatrix} + 3 \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix} - 6 \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}$$

Hence,  $[v]_B = [4, 3, -6]^T$ .



**Sol.** Clearly,

$$\begin{bmatrix} 25 & 24 \\ 0 & -9 \end{bmatrix} = 4 \begin{bmatrix} 7 & 3 \\ 0 & 0 \end{bmatrix} + 3 \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix} - 6 \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}$$

Hence,  $[v]_B = [4, 3, -6]^T$ .

$$\text{Now } P = \begin{bmatrix} 1 & -2 & 1 \\ -4 & 1 & 1 \\ 1 & 1 & -1 \end{bmatrix} \Rightarrow$$



**Sol.** Clearly,

$$\begin{bmatrix} 25 & 24 \\ 0 & -9 \end{bmatrix} = 4 \begin{bmatrix} 7 & 3 \\ 0 & 0 \end{bmatrix} + 3 \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix} - 6 \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}$$

Hence,  $[v]_B = [4, 3, -6]^T$ .

$$\text{Now } P = \begin{bmatrix} 1 & -2 & 1 \\ -4 & 1 & 1 \\ 1 & 1 & -1 \end{bmatrix} \Rightarrow$$

$$[v]_C = P[v]_B = [-8, -19, 13]^T.$$



**Sol.** Clearly,

$$\begin{bmatrix} 25 & 24 \\ 0 & -9 \end{bmatrix} = 4 \begin{bmatrix} 7 & 3 \\ 0 & 0 \end{bmatrix} + 3 \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix} - 6 \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}$$

Hence,  $[v]_B = [4, 3, -6]^T$ .

$$\text{Now } P = \begin{bmatrix} 1 & -2 & 1 \\ -4 & 1 & 1 \\ 1 & 1 & -1 \end{bmatrix} \Rightarrow$$

$[v]_C = P[v]_B = [-8, -19, 13]^T$ . Clearly,

$$\begin{bmatrix} 25 & 24 \\ 0 & -9 \end{bmatrix} = -8 \begin{bmatrix} 22 & 7 \\ 0 & 2 \end{bmatrix} - 19 \begin{bmatrix} 12 & 4 \\ 0 & 1 \end{bmatrix} + 13 \begin{bmatrix} 33 & 12 \\ 0 & 2 \end{bmatrix}$$



**Theorem:** Let  $B$  and  $C$  be ordered bases for a nontrivial finite dimensional vector space  $V$ , and let  $P$  be the transition matrix from  $B$  to  $C$ . Then  $P$  is nonsingular, and  $P^{-1}$  is the transition matrix from  $C$  to  $B$ .



## Example 8

**Q:.** For an ordered basis

$B = ([1, -4, 1, 2, 1], [6, -24, 5, 8, 3], [3, -12, 3, 6, 2])$  of a subspace  $V$  of  $\mathbb{R}^5$ ,





## Example 8

**Q:.** For an ordered basis

$B = ([1, -4, 1, 2, 1], [6, -24, 5, 8, 3], [3, -12, 3, 6, 2])$  of a subspace  $V$  of  $\mathbb{R}^5$ , perform the following steps:



## Example 8

**Q:.** For an ordered basis

$B = ([1, -4, 1, 2, 1], [6, -24, 5, 8, 3], [3, -12, 3, 6, 2])$  of a subspace  $V$  of  $\mathbb{R}^5$ , perform the following steps:

- Use the Simplified Span Method to find a second ordered basis  $C$ .



## Example 8

**Q:.** For an ordered basis

$B = ([1, -4, 1, 2, 1], [6, -24, 5, 8, 3], [3, -12, 3, 6, 2])$  of a subspace  $V$  of  $\mathbb{R}^5$ , perform the following steps:

- Use the Simplified Span Method to find a second ordered basis  $C$ .
- Find the transition matrix  $P$  from  $B$  to  $C$ .



## Example 8

**Q:.** For an ordered basis

$B = ([1, -4, 1, 2, 1], [6, -24, 5, 8, 3], [3, -12, 3, 6, 2])$  of a subspace  $V$  of  $\mathbb{R}^5$ , perform the following steps:

- Use the Simplified Span Method to find a second ordered basis  $C$ .
- Find the transition matrix  $P$  from  $B$  to  $C$ .
- Find the transition matrix  $Q$  from  $C$  to  $B$ .



## Example 8

**Q:.** For an ordered basis

$B = ([1, -4, 1, 2, 1], [6, -24, 5, 8, 3], [3, -12, 3, 6, 2])$  of a subspace  $V$  of  $\mathbb{R}^5$ , perform the following steps:

- Use the Simplified Span Method to find a second ordered basis  $C$ .
- Find the transition matrix  $P$  from  $B$  to  $C$ .
- Find the transition matrix  $Q$  from  $C$  to  $B$ .
- For the given vector  $v = [2, -8, -2, -12, 3] \in V$ , calculate  $[v]_B$  and  $[v]_C$ .



- Use the Simplified Span Method to find a second ordered basis  $C$ .



- Use the Simplified Span Method to find a second ordered basis  $C$ .

**Sol.** Here

$$B = \begin{bmatrix} 1 & -4 & 1 & 2 & 1 \\ 6 & -24 & 5 & 8 & 3 \\ 3 & -12 & 3 & 6 & 2 \end{bmatrix} \Rightarrow$$



- Use the Simplified Span Method to find a second ordered basis  $C$ .

**Sol.** Here

$$B = \begin{bmatrix} 1 & -4 & 1 & 2 & 1 \\ 6 & -24 & 5 & 8 & 3 \\ 3 & -12 & 3 & 6 & 2 \end{bmatrix} \Rightarrow$$

$$\text{REF}(B) = \begin{bmatrix} 1 & -4 & 0 & -2 & 0 \\ 0 & 0 & 1 & 4 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix} \Rightarrow$$





- Use the Simplified Span Method to find a second ordered basis  $C$ .

**Sol.** Here

$$B = \begin{bmatrix} 1 & -4 & 1 & 2 & 1 \\ 6 & -24 & 5 & 8 & 3 \\ 3 & -12 & 3 & 6 & 2 \end{bmatrix} \Rightarrow$$

$$\text{REF}(B) = \begin{bmatrix} 1 & -4 & 0 & -2 & 0 \\ 0 & 0 & 1 & 4 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix} \Rightarrow$$

$$C = ([1, -4, 0, -2, 0], [0, 0, 1, 4, 0], [0, 0, 0, 0, 1])$$



- Find the transition matrix  $P$  from  $B$  to  $C$ .



- Find the transition matrix  $P$  from  $B$  to  $C$ .

The transition matrix  $P$  from  $B$  to  $C$  is



- Find the transition matrix  $P$  from  $B$  to  $C$ .

The transition matrix  $P$  from  $B$  to  $C$  is

$$P = \begin{bmatrix} 1 & 6 & 3 \\ 1 & 5 & 3 \\ 1 & 3 & 2 \end{bmatrix}$$



- Find the transition matrix  $Q$  from  $C$  to  $B$ .



- Find the transition matrix  $Q$  from  $C$  to  $B$ .

The transition matrix  $Q$  from  $C$  to  $B$  is



- Find the transition matrix  $Q$  from  $C$  to  $B$ .

The transition matrix  $Q$  from  $C$  to  $B$  is

$$Q = P^{-1} = \begin{bmatrix} 1 & -3 & 3 \\ 1 & -1 & 0 \\ -2 & 3 & -1 \end{bmatrix}$$



- For the given vector  $v = [2, -8, -2, -12, 3] \in V$ , calculate  $[v]_B$  and  $[v]_C$ .





- For the given vector  $v = [2, -8, -2, -12, 3] \in V$ , calculate  $[v]_B$  and  $[v]_C$ .

$$[B|v] = \left[ \begin{array}{ccc|c} 1 & 6 & 3 & 2 \\ -4 & -24 & -12 & -8 \\ 1 & 5 & 3 & -2 \\ 2 & 8 & 6 & -12 \\ 1 & 3 & 2 & 3 \end{array} \right] = \left[ \begin{array}{ccc|c} 1 & 0 & 0 & 17 \\ 0 & 1 & 0 & 4 \\ 0 & 0 & 1 & -13 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \Rightarrow$$



$$[v]_B = [17, 4, -13]$$



$$[v]_B = [17, 4, -13]$$

$$\text{Now } P[v]_B = [v]_C \implies$$



$$[v]_B = [17, 4, -13]$$

$$\text{Now } P[v]_B = [v]_C \implies [v]_C = [2, -2, 3]$$



## Exercises

**Q:.** For the ordered bases

$B = (2x^2 + 3x - 1, 8x^2 + x + 1, x^2 + 6)$  and

$C = (x^2 + 3x + 1, 3x^2 + 4x + 1, 10x^2 + 17x + 5)$  of  $P_2$ , find the transition matrix  $P$  from  $B$  to  $C$ .



## Exercises

**Q.:** For the ordered bases

$B = (2x^2 + 3x - 1, 8x^2 + x + 1, x^2 + 6)$  and

$C = (x^2 + 3x + 1, 3x^2 + 4x + 1, 10x^2 + 17x + 5)$  of  $P_2$ , find the transition matrix  $P$  from  $B$  to  $C$ .

**Sol.**  $P = \begin{bmatrix} 20 & -30 & -69 \\ 24 & -24 & -80 \\ -9 & 11 & 31 \end{bmatrix}$



**Q.:** Let  $P = \begin{bmatrix} 2 & 2 & 1 \\ 1 & -1 & 2 \\ 1 & 1 & 1 \end{bmatrix}$  be the transition matrix from  $B$  to  $C$ . If  $C = \left\{ \begin{pmatrix} 2 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\}$ , find the basis  $B$ .



**Q.:** Let  $P = \begin{bmatrix} 2 & 2 & 1 \\ 1 & -1 & 2 \\ 1 & 1 & 1 \end{bmatrix}$  be the transition matrix from  $B$

to  $C$ . If  $C = \left\{ \begin{pmatrix} 2 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\}$ , find the basis  $B$ .

**Sol.**  $B = \left\{ \begin{pmatrix} 6 \\ 3 \\ 3 \end{pmatrix}, \begin{pmatrix} 4 \\ -1 \\ 3 \end{pmatrix}, \begin{pmatrix} 5 \\ 5 \\ 2 \end{pmatrix} \right\}$





**Q:.** For an ordered basis

$B = ([3, -1, 4, 6], [6, 7, -3, -2], [-4, -3, 3, 4], [-2, 0, 1, 2])$   
of a subspace  $V$  of  $\mathbb{R}^4$ , perform the following steps:

- Use the Simplified Span Method to find a second ordered basis  $C$ .
- Find the transition matrix  $P$  from  $B$  to  $C$ .
- Find the transition matrix  $Q$  from  $C$  to  $B$ .
- For the given vector  $v = [10, 14, 3, 12] \in V$ , calculate  $[v]_B$  and  $[v]_C$ .

