

1: Path Integrals and Miscellaneous Topics

1.1 Preliminaries

Here are some things that need to be kept in mind :-

$$\begin{aligned}
 X|x\rangle &= x|x\rangle \\
 \langle x|x'\rangle &= \delta(x-x') \\
 \int dx |x\rangle\langle x| &= 1 \\
 P|p\rangle &= p|p\rangle \\
 \langle p|x\rangle &= \frac{1}{\sqrt{2\pi\hbar}} \int dp e^{-ipx/\hbar} = \langle x|p\rangle^* \\
 f(x) = \langle x|f\rangle &= \int dp \langle x|p\rangle \langle p|f\rangle = \frac{1}{\sqrt{2\pi\hbar}} \int dp e^{-ipx/\hbar} \langle p|f\rangle
 \end{aligned}$$

1.2 Ordering and Conventions

In the Hmailtonian way of looking at things, we just promote the classical observables to operators while making the classical \rightarrow quantum transition. But there is a slight problem - $[x_{cl}, p_{cl}] = 0$, but $[x_{op}, p_{op}] \neq 0$. Hence, we need to worry about the order in which we write the p 's and x 's.

There is no unique well-defined principle to do this. Two conventions are defined below :-

- **Normal Ordering** - One just puts all p 's on the left of all the x 's. For example.

$$px^2 \xrightarrow{NO} px^2$$

$$xpx \xrightarrow{NO} px^2$$

$$x^2p \xrightarrow{NO} px^2$$

- **Weyl Ordering** - One symmetrizes the product, and weights them equally.

$$px \xrightarrow{WO} \frac{px + xp}{2}$$

$$px^2 \xrightarrow{WO} \frac{px^2 + xpx + x^2p}{3}$$

$$x^m p^n \xrightarrow{WO} \frac{(\dots)}{\binom{m+n}{m}}$$

How is the normal ordered Hamiltonian related to the classical Hamiltonian? We consider the matrix elements of H^{NO} , $\langle x'|H^{NO}|x\rangle$.

$$\langle x'|H^{NO}|x\rangle = \int \langle x'|p\rangle \langle p|H^{NO}|x\rangle dp = \int dp e^{-\frac{ip(x-x')}{\hbar}} H(p,x)$$