

Running head: Burn severity and ecosystem transformation

Title: Fuel connectivity, burn severity, and seedbank survivorship drive ecosystem transformation in a semi-arid shrubland.

Adam L. Mahood<sup>1,2,3,\*</sup>, Michael J. Koontz<sup>2</sup>, Jennifer K. Balch<sup>1,2,4</sup>

<sup>1</sup> Department of Geography, University of Colorado Boulder, Boulder, CO, USA

<sup>2</sup> Earth Lab, University of Colorado, Boulder, CO, USA

<sup>3</sup> Water Resources, Agricultural Research Service, United States Department of Agriculture, Fort Collins, CO, USA

<sup>4</sup> Environmental Data Science Innovation and Inclusion Lab, University of Colorado, Boulder, CO

\* Corresponding author: [admahood@gmail.com](mailto:admahood@gmail.com)

Open Research Statement: Data and code to recreate the analysis are freely available at  
<https://doi.org/10.5281/zenodo.5293996>.

<sup>1</sup> **Abstract**

<sup>2</sup> A key challenge in ecology is understanding how multiple drivers interact to precipitate  
<sup>3</sup> persistent vegetation state changes. These state changes may be both precipitated and  
<sup>4</sup> maintained by disturbances, but predicting whether the state change is fleeting or persistent  
<sup>5</sup> requires an understanding of the mechanisms by which disturbance affects the alternative  
<sup>6</sup> communities. In the sagebrush shrublands of the western United States, widespread annual  
<sup>7</sup> grass invasion has increased fuel connectivity, which increases the size and spatial contiguity  
<sup>8</sup> of fires, leading to post-fire monocultures of introduced annual grasses (IAG). The novel  
<sup>9</sup> grassland state can be persistent, and more likely to promote large fires than the shrubland  
<sup>10</sup> it replaced. But the mechanisms by which pre-fire invasion and fire occurrence are linked to  
<sup>11</sup> higher post-fire flammability are not fully understood. A natural experiment to explore these  
<sup>12</sup> interactions presented itself when we arrived in northern Nevada immediately after a 50,000  
<sup>13</sup> ha wildfire was extinguished.

<sup>14</sup> We hypothesized that the novel grassland state is maintained via a reinforcing feedback  
<sup>15</sup> where higher fuel connectivity increases burn severity, which subsequently increases post-fire  
<sup>16</sup> IAG dispersal, seed survivorship, and fuel connectivity. We used a Bayesian joint species  
<sup>17</sup> distribution model and structural equation model framework to assess the strength of the  
<sup>18</sup> support for each element in this feedback pathway. We found that pre-fire fuel connectivity  
<sup>19</sup> increased burn severity and that higher burn severity had mostly positive effects on the  
<sup>20</sup> occurrence of IAG and another non-native species, and mostly negative or neutral relationships  
<sup>21</sup> with all other species. Finally, we found that the abundance of IAG seeds in the seedbank  
<sup>22</sup> immediately post-fire had a positive effect on the fuel connectivity 3 years after fire, completing  
<sup>23</sup> a positive feedback promoting IAG. These results demonstrate that the strength of the positive  
<sup>24</sup> feedback is controlled by measurable characteristics of ecosystem structure, composition  
<sup>25</sup> and disturbance. Further, each node in the loop is affected independently by multiple  
<sup>26</sup> global change drivers. It is possible that these characteristics can be modeled to predict

27 threshold behavior and inform management actions to mitigate or slow the establishment of  
28 the grass-fire cycle, perhaps via targeted restoration applications or pre-fire fuel treatments.

29 *Keywords:* *Artemisia tridentata*, alternative stable states, *Bromus tectorum*, burn severity,  
30 cheatgrass, fuel connectivity, grass-fire cycle, joint species distribution model, resilience,  
31 sagebrush

## 32 1. Introduction

33 Ecosystems around the world are being affected simultaneously by multiple facets of global  
34 change. For example, changes in land use can facilitate exotic plant invasions ([Allan et al.](#)  
35 [2015](#)), which can alter ecosystem structure ([Davies and Nafus 2013](#)). Altered structure can  
36 change the likelihood of a disturbance, the properties of a disturbance and the capacity  
37 of the system to recover after a disturbance ([Brooks et al. 2004](#)). Global climate change  
38 can also directly affect the magnitude of disturbances ([S. A. Parks and Abatzoglou 2020](#)),  
39 and act as a demographic filter that influences how ecosystems recover after disturbances  
40 ([Rother, Veblen, and Furman 2015](#); [Davis et al. 2019](#)) via impacts on adult plant survival  
41 and seed dispersal ([Davis, Higuera, and Sala 2018](#); [Eskelinen et al. 2020](#)). The combined  
42 effects of global change forces on structure, function and disturbance can cascade and interact.  
43 For example, while burn severity (or the proportion of biomass burned ([Keeley 2009](#))) is  
44 influenced by vegetation structure ([Koontz et al. 2020](#); [Sean A. Parks et al. 2018](#)), it also  
45 increases with temperature and aridity ([S. A. Parks and Abatzoglou 2020](#)). These forces  
46 can ultimately lead to permanent compositional change, biodiversity losses and the loss of  
47 ecosystem services ([Ratajczak et al. 2018](#); [Mahood and Balch 2019](#); [Mahood et al. 2022](#))  
48 due to internal, self-reinforcing mechanisms that arise from those structural and functional  
49 changes which then maintain an alternative stable state ([Marten Scheffer and Carpenter](#)  
50 [2003](#); [Ratajczak et al. 2018](#)).

51 There is a long history of univariate time series observations that show sudden state changes

52 (Marten Scheffer and Carpenter 2003), and these have informed the development of theories  
53 that help us understand how systems of any type can change state suddenly, and  
54 exist in persistent alternative stable states (Marten Scheffer et al. 2015; Ratajczak et al.  
55 2018). These theories typically represent the system's state with a single variable, of which  
56 the mean is observed to abruptly change in time or space (Marten Scheffer et al. 2015).  
57 Descriptive evidence of alternative stable states has been documented at broad scales in  
58 tropical ecosystems, where forests, savannas and grasslands are considered alternative stable  
59 states because they are floristically distinct (Aleman et al. 2020) and cluster around static  
60 values of woody cover (80, 30 and 0 percent) while occurring along overlapping ranges of  
61 precipitation (Hirota et al. 2011; Staver, Archibald, and Levin 2011). The forested state has  
62 a self-reinforcing, positive feedback between evapotranspiration and tree cover (Staal et al.  
63 2020), while the grassland and savanna states are maintained by feedbacks between grass  
64 flammability and fire occurrence (D'Antonio and Vitousek 1992; Staver, Archibald, and Levin  
65 2011). Alternative stable states are believed to be widespread (M. Scheffer et al. 2001), but  
66 their existence is rarely proven at broader scales, with most demonstrative studies having  
67 been conducted in greenhouse and laboratory microcosm experiments (Schröder, Persson,  
68 and De Roos 2005). One of the reasons for this is that ecological systems are much more  
69 complex than a simple bivariate system with a single driver and a single response. There  
70 may be multiple drivers, and the state is the product of interactions between organisms and  
71 their immediate environment, as well as countless inter- and intra-specific interactions.

72 A central challenge in ecology in the 21st century is to move from describing how plant  
73 communities are affected by global change to the capacity to predict how species pools will  
74 assemble and persist in response to global change (Davis, Higuera, and Sala 2018; Keddy  
75 and Laughlin 2021). Prediction of community response to multi-faceted global change  
76 drivers is enhanced with a better understanding of the mechanisms that underlie community  
77 stability in the face of disturbances. A classic example of an ecosystem that appears to have  
78 disturbance-mediated alternative stable states (but see Morris and Leger (2016)), but whose

79 stability mechanisms aren't well understood is the invasion of *Bromus tectorum* L. and other  
80 introduced annual grasses in the Great Basin of the western United States. Here, it is well  
81 documented how the interaction of annual grass invasion, fire ([Balch et al. 2013](#)) and grazing  
82 ([Williamson et al. 2019](#)) are associated with the degradation or loss of over half of Wyoming  
83 big sagebrush (*Artemisia tridentata* ssp. *wyomingensis* Beetle & Young) ecosystems ([Davies  
et al. 2011](#)). These systems had a precolonial fire regime of infrequent, patchy fires ([Bukowski  
and Baker 2013](#)). In uninvaded areas, the space between shrubs is typically composed of bare  
86 ground covered in biological soil crust and caespitose perennial plants (Figure 1). Because  
87 fire does not spread readily below a threshold of approximately 60% cover of flammable  
88 vegetation ([Archibald, Staver, and Levin 2012](#)), the low fuel connectivity in these areas limits  
89 fire spread. Annual grass invasion increases fuel connectivity while decreasing fuel moisture  
90 ([Brooks et al. 2004; Davies and Nafus 2013](#)), leading to increased fire size and frequency  
91 ([Balch et al. 2013](#)). Sagebrush stands with high native perennial cover might need only a  
92 small amount of additional annual grass cover to alter ecosystem structure enough to alter the  
93 fire regime (Figure 2). After fire, the landscape is typically dominated by introduced annual  
94 grasses. But in order to understand how fire drives the persistence of the grassland state, we  
95 need to understand the demographic mechanisms by which fire impacts propagule dispersal  
96 and benefits the alternative state ([Davis, Higuera, and Sala 2018](#)). As with forested systems,  
97 propagule dispersal is a key filter through which species must pass in order to establish and  
98 persist in a post-fire landscape ([Gill et al. 2022](#)).

99 Petraitis and Latham ([1999](#)) posited that the maintenance of alternate species assemblages  
100 requires first a disturbance that removes the species from the initial assemblage and second  
101 the arrival of the species of the alternate assemblage. One understudied mechanism that  
102 may explain both for the *Artemisia/Bromus* system is the interaction between the species  
103 composition of the soil seed bank and burn severity. Because the invading species are annual,  
104 and many of the key native plant species are seed obligates, the seed is the key life history stage  
105 that fire must act upon to benefit the invading plants. Seeds and seedlings are particularly

<sup>106</sup> vulnerable to climate, competition and disturbance (Enright et al. 2015). Warmer and drier  
<sup>107</sup> conditions simultaneously reduce recruitment, growth, and survival of seeds and seedlings  
<sup>108</sup> (Enright et al. 2015; Schlaepfer, Lauenroth, and Bradford 2014), while also increasing burn  
<sup>109</sup> severity (S. A. Parks and Abatzoglou 2020). In fire prone ecosystems, seed obligate species  
<sup>110</sup> typically have life history strategies to cope with fires that burn at different severities (Maia  
<sup>111</sup> et al. 2012; Wright, Latz, and Zuur 2016; Palmer, Denham, and Ooi 2018). Soil heating from  
<sup>112</sup> fire affects the response of vegetation to fire (Gagnon et al. 2015), including the capacity of  
<sup>113</sup> seeds to remain viable after fire (Humphrey and Schupp 2001). High severity fire can affect  
<sup>114</sup> species that use the seedbank positively (Kimura and Tsuyuzaki 2011), negatively (Heydari  
<sup>115</sup> et al. 2017), or have no effect (Lipoma, Funes, and Díaz 2018), depending on species-specific  
<sup>116</sup> adaptations. Both the depth of the burn and fire temperature can affect subsequent recovery  
<sup>117</sup> by seed germination (Morgan and Neuenschwander 1988; Schimmel and Granström 1996), as  
<sup>118</sup> well as seed mortality and physical seed dormancy mechanisms (Liyanage and Ooi 2017).

<sup>119</sup> In addition to size and frequency, exotic plant invasions can alter fire temperature (Brooks et  
<sup>120</sup> al. 2004; R. O. Jones et al. 2015) and burn severity. While in many cases fires that burn  
<sup>121</sup> at higher temperatures will also consume more biomass (i.e. burn at higher severity), grass  
<sup>122</sup> fires may not always have such a relationship. Direct measurements have shown that *B.*  
<sup>123</sup> *tectorum* burns at low temperatures (Beckstead et al. 2011; Germino, Chambers, and Brown  
<sup>124</sup> 2016), but because it also increases horizontal fuel connectivity (Davies and Nafus 2013), it  
<sup>125</sup> leads to more contiguously burned areas and therefore higher burn severity, despite lower fire  
<sup>126</sup> temperatures. To benefit from fire, *B. tectorum* would need to gain a fitness benefit relative  
<sup>127</sup> to other species

<sup>128</sup> One way to achieve this is to disperse more viable seeds into the post-fire landscape than  
<sup>129</sup> the other species and become well-represented in the post-fire plant assemblage (Bond and  
<sup>130</sup> Midgley 1995). If the fire is patchy, this can happen through post-fire seed dispersal (Monty,  
<sup>131</sup> Brown, and Johnston 2013). Without unburned patches, seeds must survive the fire. If  
<sup>132</sup> the increase in fuel connectivity caused by *B. tectorum* increases the severity of fire, one

way burn severity might then influence the community composition of the post-fire seed bank to facilitate the post-fire dominance of *B. tectorum* would be to burn a contiguous area at a temperature high enough to kill fire-intolerant native seeds, but low enough that *B. tectorum* seeds survive and germinate more readily from fire-induced germination cues (Naghipour et al. 2016; Fenesi et al. 2016). In other words, an area with high burn severity should have a lower relative occurrence of viable seeds of native species, and a higher relative occurrence of the seeds of fire-tolerant introduced annual plants. This would allow for the often-observed dominance of introduced annual grasses after a few years and would result in higher fuel connectivity, closing the positive feedback loop. Plants that are not adapted to frequent fire would be less likely to produce seeds that are adapted to surviving fire, or dispersal mechanisms to take advantage of the resources available immediately after fire (Keeley et al. 2011). To our knowledge, despite several studies on the relationship between fire occurrence and the seed bank in this system (Hassan and West 1986; Humphrey and Schupp 2001; Boudell, Link, and Johansen 2002), no studies to date have examined the effect of burn severity on the seed bank. Burn severity is more ecologically meaningful than fire occurrence, and is more useful for understanding threshold effects and stable states than a binary variable.

Here, we collected soil cores from 14 locations along the perimeter of a large fire (the Hot Pot fire, ~50,000 ha) immediately after it was extinguished, in northern Nevada in July 2016. Each location had paired burned and unburned samples. Because it burned a large area in only three days, we could sample a broad area while being reasonably certain that the weather conditions during the fire were similar at all sites. Because we collected our samples immediately after the fire was extinguished, we felt confident that the seed bank samples did not contain seeds deposited by post-fire dispersal. We put the samples in cold storage and germinated the seeds from those cores in a greenhouse the following spring. In spring 2017 and fall 2019 we collected information on vegetation structure and diversity at each location. We tested three hypotheses in this study that are depicted in Figure 3: (H1)

160 Pre-fire fuel connectivity would be positively related to burn severity; (H2) burn severity  
161 would increase the occurrence probability of introduced annual species in the seed bank and  
162 reduce the occurrence probability of native species; and (H3) the abundance of post-fire  
163 *B. tectorum* seeds in the seedbank would be positively related to post-fire fuel connectivity.  
164 We examined two alternatives to H2: (H2a), increased fuel connectivity brought on by the  
165 invasion of annual grasses may have already depleted the diversity of the soil seed bank  
166 before the fire occurred; and (H2b) prefire fuel connectivity is solely reflective of annual grass  
167 cover, which drives post-fire annual grass seed abundance. In addition, because in our study  
168 system post-fire sites are floristically distinct from the pre-fire state ([Mahood and Balch](#)  
169 [2019](#)), typically with near monocultures of *B. tectorum*, we hypothesized that (H4) high  
170 post-fire fuel connectivity of those near-monocultures would result in lower aboveground  
171 species diversity due to competitive exclusion of native plants.

## 172 2. Methods

### 173 2.1 Study Area

174 The study was conducted in north-central Nevada the day after a large fire (the Hot Pot Fire)  
175 was extinguished (Appendix S1, Fig. S1). The Hot Pot Fire burned just over 50,000 hectares  
176 in less than a week. The pre-fire landcover was predominantly *B. tectorum* and Wyoming big  
177 sagebrush plant communities. The fire occurred after the early season plants, including *B.*  
178 *tectorum* and *Poa secunda* J. Presl, the most abundant native understory species, had gone  
179 to seed, and before the late season species, including Wyoming big sagebrush, had produced  
180 flowers. Thus we were able to isolate the effect of the fire without any confounding effects of  
181 post-fire seed dispersal, while achieving a broad spatial extent. The sites we sampled ranged  
182 from 1,397 to 1,607 meters in elevation.

### 183 2.2 Seed Bank Sampling

184 In early July 2016, we collected samples of the soil seed bank at fourteen locations the day

after the Hot Pot fire was contained. Each site was located at the perimeter of the fire where it was clearly delineated by a bulldozer line or in one case a narrow dirt road. We were confident paired sites were of the same pre-fire composition because we had been working in these areas all summer collecting data for another study. Eleven sites were mature sagebrush communities with no history of fire since at least 1984. Three sites had previously burned in 1984 according to the Monitoring Trends in Burn Severity (MTBS) fire history (Eidenshink et al. 2007) and had high cover of *B. tectorum*, but still had scattered sagebrush cover. We used a metal stake to mark paired burned and unburned sampling locations on each side of the perimeter, 10 m from the nearest evidence of anthropogenic disturbance (i.e. bulldozer effects, footprints) associated with active fire suppression along the perimeter. Within 3 m of each marker, we extracted twelve, 6 cm deep, 5 cm diameter, soil cores. Seeds of sagebrush generally do not fall far (<30 m) from their parent plants in this system (Shinneman and McIlroy 2016), and so they are not uniformly distributed (Boudell, Link, and Johansen 2002). In addition, seeds from *B. tectorum* and *Artemisia* have different germination rates based on the micro-site they find themselves in (i.e. under a shrub or in the bare ground between shrubs, Eckert et al. 1986). To account for these potentially confounding effects, we placed half of the core locations under shrubs, half in shrub interspaces, and aggregated the cores for each site. In the burned areas, it was obvious where shrubs had been located. Even when they were completely incinerated, their imprint remained on the soil surface (Bechtold and Inouye 2007). To examine the effect of seed depth, we divided each soil core into 0-2 cm and 2-6 cm depths. Litter was aggregated with the 0-2 cm samples. Samples were then placed in cold storage (~2 deg C) for 3 months (Meyer, Monsen, and Mcarthur 2013). At all sites, to be sure that we were at a site where sagebrush germination could occur we checked for first year germinants on the unburned side (we found them at all sites), and to ensure that there were no confounding effects of post-fire seed dispersal, we determined whether or not the sagebrush were flowering (they were not flowering at all sites), and recorded species occupancy for all aboveground plant species.

212 We followed the methodology of Ter Heert et al. (1996) to germinate the seeds. Each sample  
213 was run through 0.2 mm sieve, and spread in a 3-5 mm layer over the top of 1 - 4 pots.  
214 These pots were filled 3 cm deep with potting soil, topped by a thin layer of sand. Pots were  
215 watered as needed to stay at field capacity. Every week emerging germinants were identified,  
216 counted and removed. Most of the germination occurred within 6 weeks, and after 8 weeks  
217 we ended the germination assay.

218 *2.3 Post-Fire Vegetation Sampling*

219 We sampled the aboveground fuel structure and plant diversity in May 2017, the growing  
220 season immediately after the fire and again in September 2019. At each location, we established  
221 50m transects starting at the boundary of the burned and unburned sides of the perimeter,  
222 running perpendicular to the fire perimeter, and marked the transect ends with rebar. In  
223 order to characterize aboveground plant diversity, we measured the occupancy and abundance  
224 of all plant species by measuring cover of every species in 0.1 m<sup>2</sup> quadrats spaced every 5  
225 m along each transect. We measured shrub cover (coarse fuels) and herbaceous plant cover  
226 (fine fuels) using the line intercept method along the transect, a commonly-used approach for  
227 characterizing fuel structure (Elzinga, Salzer, and Willoughby 1998). We calculated total  
228 vegetation cover (TVC) as the sum of the fine and coarse fuel measurements. Both live and  
229 dead plants were included in these measurements.

230 *2.4 Remotely-Sensed Burn Severity*

231 We downloaded the “fire bundle” of the Hot Pot fire from www.mtbs.gov. This included  
232 cloud-free Landsat 8 scenes collected before the Hot Pot fire, and already calculated layers of  
233 the Differenced Normalized Burn Ratio (dNBR, Equations 1 & 2, J. D. Miller et al. 2009).  
234 Because our sites were generally within 10 meters of the burn perimeter, The pixels directly  
235 intersecting the site locations were likely to be mixed pixels (i.e. containing burned and  
236 unburned ground). To minimize this effect, we extracted all the dNBR values within a 120  
237 meter buffer of each seed bank site for pixels whose centroids fell inside of the fire perimeter

238 and calculated the mean.

239 **Equation 1:**  $NBR = (NIR - SWIR_1) / (NIR + SWIR_1)$

240 **Equation 2:**  $dNBR = (NBR_{prefire} - NBR_{postfire}) * 1000$

241 *2.5 Statistical Analysis*

242 Our statistical analysis centered around trying to understand each component of the positive  
243 feedback loop posited by the 4 hypotheses described above. In order to understand how  
244 pre-fire fuel connectivity influenced burn severity (H1), we used total vegetation cover (TVC)  
245 from two separate data sources as a proxy for fuel connectivity, and created separate linear  
246 models with TVC as the predictor variable and burn severity (dNBR, [J. D. Miller et al.](#)  
247 [2009](#)) as the response variable. With the field data we collected, we created an ordinary  
248 least squares (OLS) linear model with burn severity as the dependent variable and TVC  
249 (defined as shrub cover plus herbaceous plant cover from the unburned side of the paired  
250 sites), elevation and aspect as independent variables.

251 We were concerned that because our data were collected at the edge of the fire, the burn  
252 severity calculated at each point may have included partially burned pixels. So, as a  
253 supplement, we examined the same relationship by creating a model of TVC using Landsat  
254 Thematic Mapper (TM) surface reflectance data using field measurements of TVC from the  
255 Bureau of Land Management's Assessment, Inventory and Monitoring dataset (AIM, [U.S.](#)  
256 [Department of Interior 2018](#)). The AIM dataset contained 813 sampling locations within  
257 the Central Basin and Range ecoregion ([Commission for Environmental Cooperation 2006](#))  
258 that were visited by BLM field crews between 2011 and 2015. They were mostly sampled  
259 once but there were some repeats, for 1,117 total measurements. For each of these points,  
260 we extracted the surface reflectance values of each Landsat band for the sampling year  
261 near peak biomass using a cloud-free scene from May or early June. Then, we used those  
262 surface reflectance values to calculate various vegetation indexes (Appendix S1: Table S1),  
263 including the Green Normalized Difference Vegetation Index (Green NDVI, Equation 3), and

264 Normalized Difference Senesced Vegetation Index (NDSVI, Equation 4). We used these two  
265 indexes and their interactions as predictors in a generalized linear model of TVC with a  
266 beta distribution. We used the model to create a layer of estimated pre-fire TVC for the  
267 study area, and extracted both our predictions of TVC and dNBR of the fire from 1000  
268 regularly-spaced points within the fire perimeter. Finally, to quantify the effect of TVC on  
269 burn severity, we created an OLS linear model with our modeled TVC and its second-order  
270 polynomial as predictor variables and burn severity as the response variable.

271 **Equation 3:**  $Green\ NDVI = \frac{NIR-Green}{NIR+Green}$

272 **Equation 4:**  $NDSVI = \frac{SWIR_1-Red}{SWIR_1+Red}$

273 To examine how burn severity affected the community composition of the seed bank (H2),  
274 we created a joint species distribution model (JSDM) in a Bayesian framework ([Tikhonov et](#)  
275 [al. 2020](#)) for the occurrence of all species germinated from the seed bank that were found  
276 at more than one location. We created four Markov Chain Monte Carlo (MCMC) chains,  
277 each consisting of 150,000 iterations. We discarded the first 50,000 iterations for each chain  
278 and then recorded every 100th for a total of 1,000 posterior samples per chain, and 4,000  
279 total. We assessed model convergence using the effective sample size and the potential  
280 scale reduction factor ([Gelman, Rubin, et al. 1992](#)). We used the model to predict the  
281 probability of occurrence of germinable seeds of a given species along a gradient of burn  
282 severity. We included burn severity, elevation, aspect, pre-fire seedbank diversity and soil  
283 depth as independent variables.

284 To account for the possibility that increased fuel connectivity brought on by the invasion  
285 of annual grasses may have already depleted the diversity of the soil seed bank before the  
286 fire occurred (H2a) as a confounding factor, we included the Shannon-Weaver diversity  
287 index ([Shannon and Weaver 1949](#)) in the paired, unburned seed bank samples as one of the  
288 predictor variables in our JSDM. We also created OLS models with the unburned species  
289 richness and Shannon-Weaver diversity index predicted by prefire fuel connectivity, with the

expectation that pre-fire fuel connectivity would have had a negative effect on the prefire seedbank diversity. To examine how community composition and burn severity then affected subsequent fuel connectivity (H3), we created OLS models with fuel connectivity three years post-fire as the dependent variable, and burn severity, seed counts for *B. tectorum*, *P. secunda* and other species, elevation, aspect, depth, and alpha diversity as independent variables. To examine how the resulting fuel connectivity was related to biodiversity (H4), we used the aboveground diversity data and connectivity data that we collected in 2019 to create a Poisson GLM with number of species encountered at each site as the dependent variable, as well as an OLS linear model with the Shannon-Weaver index for the plant species as a dependent variable. We used fuel connectivity, elevation, and aspect as independent variables.

In order to examine hypotheses 1-3 in a single framework we constructed a path model (Rosseel 2012). We had paths leading from pre-fire connectivity, through burn severity to the log of the post-fire count of *B. tectorum* seeds in the seedbank, and finally to post-fire connectivity. Pre-fire cover of *B. tectorum*, elevation, pre-fire seed bank diversity and pre-fire aboveground diversity were also accounted for.

All analyses were done in R (R Core Team 2020). Data and code to recreate the analysis are freely available at <https://doi.org/10.5281/zenodo.5293996>.

### 3. Results

We found support for each hypothesized component of the positive feedback loop independently and when combined in the path model ( $\chi^2 = 3.17$ ,  $p = 0.39$ , Figure 4a, Appendix S1, Tables S4 & S5). For H1, TVC had a weak positive relationship with burn severity ( $\beta = 2.4$ ,  $p = 0.083$ ,  $R^2 = 0.27$ , Figure 4b, Appendix S1: Table S2). For our remotely sensed analysis, Green NDVI, NDSVI and their interaction explained 35% of the variation in pre-fire TVC (Appendix S1: Table S2). This predicted TVC had a positive relationship with burn severity ( $p \ll 0.01$ ,  $R^2 = .42$ , Figure 4b, Appendix S1: Table S2).

315 The majority of seeds that germinated in the greenhouse were the two most common grass  
316 species, *P. secunda* and *B. tectorum* (Appendix S1: Table S3, Fig. S2). Eight dicot species  
317 were found in more than one location, and these 10 prevalent species are those that were  
318 used in our JSDM. Burned sites had an average of  $34 \pm 32$  total seeds in the top 2 cm, and  
319  $12 \pm 14$  in the bottom 4 cm. Unburned sites had an average of  $299 \pm 170$  in the top 2 cm  
320 and  $59 \pm 29$  in the bottom 4 cm (Appendix S1: Fig. S3). For H2, the JSDM converged  
321 well (Appendix S1: Fig S4). Gelman diagnostics were all very close to 1 and the effective  
322 sample size centered on 4,000, which indicated good model convergence. Elevation had the  
323 strongest effects on individual species occurrence and explained the most variance on average  
324 (36%). Burn severity explained 23% of the variance on average and was supported at the 95%  
325 level for 5 species (Appendix S1: Fig S2b). For the introduced species, the predictions along  
326 a gradient of burn severity were positive for *B. tectorum*, *Sisymbrium altissimum* L. and  
327 *Lepidium perfoliatum* L., and negative for *Ceratocephala testiculata* and *Alyssum desertorum*  
328 Stapf (Figure 4e). For native species, the effect of burn severity on occurrence was positive  
329 for *A. tridentata*, but the mean predictions were still low, never rising above 50%. It was  
330 neutral for *P. secunda* and negative for the remaining species. Testing H2a revealed a positive  
331 relationship between pre-fire aboveground species diversity and pre-fire fuel connectivity in  
332 the single model, and neutral relationships in the path model, and so we felt it was reasonable  
333 to rule out pre-fire fuel connectivity as a confounding factor for H2. Testing H2b showed a  
334 negative relationship, allowing us to rule out the idea that both pre-fire connectivity and  
335 post-fire seed bank composition were simply a function of pre-fire annual grass cover.

336 For H3, we found that, after accounting for elevation, pre-fire aboveground richness, and  
337 the number of *P. secunda* seeds, the number of *B. tectorum* seeds in the post-fire seedbank  
338 was positively associated with the fuel connectivity in 2019 ( $\beta = 0.54$ ,  $p = 0.01$ , Adj  $R^2 =$   
339 0.75, Figure 4c, Appendix S1: Table S2). For H4 the most parsimonious model (Adj  $R^2 =$   
340 0.89, Appendix S1: Table S2) had elevation, aspect, fuel connectivity and an interaction  
341 between elevation and fuel connectivity as predictors of aboveground Shannon-Weaver alpha

<sup>342</sup> diversity. Fuel connectivity was negatively associated with Shannon-Weaver diversity ( $\beta =$   
<sup>343</sup> -0.28,  $p=0.004$ , Figure 4d).

## <sup>344</sup> 4. Discussion

<sup>345</sup> Here we document how changes in ecosystem structure brought on by invasion can lead  
<sup>346</sup> to cascading effects on ecosystem function and composition via changes in the disturbance  
<sup>347</sup> regime. It has already been shown that *B. tectorum* invasion increases fire frequency (Balch  
<sup>348</sup> et al. 2013), and is indicative of a grass-fire cycle. However, an understanding of the positive  
<sup>349</sup> feedback mechanisms that link *B. tectorum* invasion success to fire occurrence is required  
<sup>350</sup> to infer the long-term persistence of such a cycle. The interaction between burn severity  
<sup>351</sup> and seed bank composition documented here may explain that link. Prior work has shown  
<sup>352</sup> that annual grass invasion increases fuel connectivity by filling in shrub interspaces with a  
<sup>353</sup> contiguous bed of fine fuels (Davies and Nafus 2013). This change in the spatial distribution  
<sup>354</sup> of fine fuels has been associated with larger and more frequent fires (Balch et al. 2013).

<sup>355</sup> Here, we found higher fuel connectivity (via TVC) increased burn severity (H1, Figure 4b).

<sup>356</sup> Higher burn severity was associated with an increased occurrence of introduced annuals in  
<sup>357</sup> the post-fire seedbank and a decreased occurrence of native plants with the exception of  
<sup>358</sup> *A. tridentata* (H2, Figure 4e), but the gains of *A. tridentata* would likely not be enough to  
<sup>359</sup> counter the gains of *B. tectorum*, especially after a few years of annual grass reproduction  
<sup>360</sup> and population growth without similar gains for the shrubs (Shriver et al. 2019). Finally,  
<sup>361</sup> greater abundance of *B. tectorum* seeds in the post-fire seedbank resulted in higher post-fire  
<sup>362</sup> fuel connectivity (H3, Figure 4c). In addition, we found evidence that high post-fire fuel  
<sup>363</sup> connectivity was associated with lower aboveground diversity (H4, Figure 4d). This suggests  
<sup>364</sup> that during inter-fire intervals, there may be additional mechanisms (e.g. competition, altered  
<sup>365</sup> ecohydrology) maintaining the post-fire, annual grass-dominated species assemblage.

<sup>366</sup> The difference in species composition before and after fire explains an apparent contradiction

367 in results between H2a (positive to neutral relationship between pre-fire fuel connectivity and  
368 diversity) and H4 (negative relationship between post-fire fuel connectivity and diversity).  
369 Most site locations had mature canopies of native shrubs with the inter-shrub space occupied  
370 mostly by native bunchgrasses and forbs, with no fire occurrence since 1984. Even in  
371 locations with high annual grass cover between shrubs, shrubs provide ecosystem structural  
372 heterogeneity and islands of fertility ([Doescher, Miller, and Winward 1984](#); [Bechtold and](#)  
373 [Inouye 2007](#)), and perennial natives that may have been established before invasion have deep  
374 roots established that allow for the avoidance of competition for water with shallow-rooted  
375 annuals ([Gibbens and Lenz 2001](#); [Ottaviani et al. 2020](#)). This may provide enough niche  
376 compartmentalization to allow native plants to persist in spite of the invasion prior to fire  
377 occurrence. Three years after fire, almost all of the sites were dominated by introduced  
378 annuals, and lacked any structural heterogeneity (Appendix S1, Fig. S6c). Thus native  
379 plants may have been able to persist via niche compartmentalization after the initial invasion,  
380 but fire burned away most of the seeds (Appendix S1, Fig. S3, S7) and removed all of the  
381 structural benefits, and microclimatic refugia that shrub cover provides. In this clean slate  
382 post-fire environment, the altered species composition of the seedbank and superior post-fire  
383 dispersal of *B. tectorum* ([Monty, Brown, and Johnston 2013](#)) allow the process of interspecific  
384 competition to be dominant ([Schlaepfer, Lauenroth, and Bradford 2014](#)).

385 ***Contrasts among forests and shrublands as it pertains to remote sensing***

386 Burn severity metrics like dNBR were conceived of in the context of forested ecosystems, and  
387 calibrated using the composite burn index ([Key and Benson 1999](#)), tree mortality, and percent  
388 change in tree canopy cover ([J. D. Miller et al. 2009](#)). It is unclear how well these metrics  
389 carry over to shrubland systems. We recorded qualitative observations of burn severity while  
390 we were sampling, mainly to ensure that we sampled a range of severities, and the dNBR  
391 we used appears to correspond with our observations. In areas where the space between  
392 shrubs was well-connected by fine fuels (Figure 1 a-c) the burn severity was higher, and the  
393 shrubs had completely burned throughout the root system, leaving only a hole in the ground

394 filled with ashes as evidence of their prior presence. In these areas the entirety of the soil  
395 surface—underneath shrub canopy and in canopy interspaces—was consumed by fire, and  
396 there was little evidence of remaining litter or biological soil crust. Areas with lower fuel  
397 connectivity had lower burn severity (Figure 1 d-f). Here, shrubs were usually consumed  
398 only to the stumps, and sometimes left standing and charred, destined for mortality. In  
399 these areas the soil surface often still had biological soil crust, partially consumed litter  
400 ([R. O. Jones et al. 2015](#)) and unconsumed annual and perennial grass bases. The manual  
401 severity classification provided by MTBS had exclusively low and medium severity, but our  
402 observations of essentially complete consumption of plant and litter tissues and very few  
403 unburned patches suggested that these should have been mostly medium and high severity.  
404 This discrepancy was not unexpected, as the ordinal burn severity classifications produced by  
405 MTBS are known to be flawed for research use ([Kolden, Smith, and Abatzoglou 2015](#)).

406 Spectral reflectance has long been used to characterize ecosystem structure, including wildfire  
407 fuels. Unique signatures of remotely-sensed spectral reflectance are typically matched to  
408 categorical fuel classifications (CFCs), which describe the physiognomy of vegetation and  
409 its potential to support various fire behavior ([Ottmar et al. 2007](#)). While different CFCs  
410 can provide a general understanding of fuel amount and connectivity, recent efforts using  
411 data with finer spatial and spectral resolution may improve fuel classification with more  
412 continuous, multi-dimensional measurements ([Stavros et al. 2018](#)). The continuous measure  
413 of NDVI in western U.S. coniferous forests is a proxy for live fuel biomass, which likely  
414 explains its positive association with wildfire severity ([Sean A. Parks et al. 2018; Koontz et al.](#)  
415 [2020](#)). NDVI also correlates with vegetation cover in these forested systems, and so greater  
416 crown connectivity may also explain the NDVI/severity relationship at local scales. When  
417 using a more direct NDVI-derived measure of vegetation connectivity in Sierra Nevada yellow  
418 pine/mixed-conifer, Koontz et al. ([2020](#)) found that greater variability in forest structure,  
419 decreased the probability of high-severity fire, likely due to decreased fuel connectivity (i.e.,  
420 live tree canopies in the yellow pine/mixed-conifer forest). Here, we arrived at a combination

421 of NDVI and NDSVI to describe the fuel connectivity of the annual grass invaded Great  
422 Basin sagebrush community to better reflect key differences in the physiognomies of forest  
423 and arid shrublands. In sagebrush shrublands, the fuel that contributes to large wildfires is a  
424 mixture of evergreen shrubs interspersed with herbaceous plants that remain green for only a  
425 portion of the growing season, and then become dry and straw-colored. Thus, both the live  
426 and dead fuel need to be taken into account in remote measurements of fuel connectivity for  
427 this system.

428 ***Management implications***

429 These results demonstrate that the strength of the grass-fire cycle in this system is controlled  
430 by measurable fire properties and ecosystem structural components. We found that annual  
431 grass cover was not the single variable that explained burn severity and fuel connectivity  
432 (Appendix S1, Fig S5). Rather, it was the contribution of annual grass cover to the total  
433 connectivity of the system (Figure 2). The most important areas to prioritize for management  
434 interventions could paradoxically be areas with relatively low levels of annual grass cover that  
435 join previously disconnected vegetation. Land managers may be able to increase their chances  
436 of restoration success by using existing methods or developing novel ones that manipulate  
437 these components to weaken or even break the positive feedback cycle. This work provides  
438 further evidence that the post-fire annual grassland is a system where the degraded state  
439 represents an alternative species assemblage from that of the restoration target. Because the  
440 propagules of the original assemblage are no longer present, methods that rely on natural  
441 succession may not be sufficient (Suding, Gross, and Houseman 2004). One-off seeding  
442 treatments have a low probability of success (Pyke et al. 2020; Arkle et al. 2022), and more  
443 labor-intensive methods involving site preparation (Farrell, Fehmi, and Gornish 2021), seed  
444 coating and priming (Pedrini et al. 2020), as well as planting live plants (Pyke et al. 2020)  
445 may improve the probability of success, as will prioritizing efforts in cooler, wetter years  
446 (Bradford et al. 2018; Hardegree et al. 2018; Shriner et al. 2018). Estimating burn severity  
447 using satellite imagery may be used in conjunction with site suitability and climate forecasts

448 to help land managers identify areas with a greater likelihood of successful seeding. Our  
449 results highlight the importance of prioritizing the preservation of existing native shrub cover  
450 and in particular policies that encourage land managers to maximize the preservation of  
451 unburned patches within the fire perimeter during the suppression of wildfires in this system  
452 (Steenvoorden et al. 2019), as these are the primary sources of native propagules and act as  
453 nurse plants (Arkle et al. 2022). In many areas, conditions are now or will in the near future  
454 be unsuitable for sagebrush due to annual grass dominance and increases in aridity (Shriver  
455 et al. 2019). In these areas it may still be feasible to restore the system's ability to sequester  
456 carbon by planting other native woody species that are more drought tolerant and resilient  
457 against fire.

458 Livestock grazing can reduce fuel connectivity in uninvaded sagebrush (Davies et al. 2010).  
459 At the same time, livestock grazing can decrease the resistance to invasion by *B. tectorum* via  
460 negative effects on biological soil crust (BSC) (Condon and Pyke 2018), and can reduce the  
461 survival of *Artemisia* seedlings that are not protected by shrub canopies (Owens and Norton  
462 1992). Targeted spring grazing in annual grass monocultures may reduce fuel connectivity  
463 and alleviate fire risk. Post-fire grazing may help reduce *B. tectorum* cover, but it may  
464 also exacerbate the problem by introducing *B. tectorum* in uninvaded sites (Williamson et  
465 al. 2019) or increasing the already superior post-fire dispersal of *B. tectorum* seeds (Monty,  
466 Brown, and Johnston 2013). Management interventions should be specifically tailored each  
467 year to the conditions of a given site, and focused on native plant restoration.

468 Herbaceous cover in these dryland systems has high interannual variability (Mahood et al.  
469 2022). Because the components of ecosystem structure and disturbance severity in positive  
470 feedback cycle described here are continuous mechanistic variables, it may be possible to  
471 develop theoretical models (*sensu* (Archibald, Staver, and Levin 2012)) to estimate the  
472 threshold of vegetation cover that will lead to high burn severity. These can then be applied  
473 in conjunction with near real time fuel loading forecasts (M. O. Jones et al. 2021) to identify  
474 areas that are vulnerable to high severity fire, which can be used by land managers to take

475 preemptive measures in high value areas.

476 ***Global environmental change implications***

477 Understanding how different facets of global environmental change create multiple mechanisms  
478 that act in concert to drive ecosystem transformation will provide important insights about  
479 ecosystem change from regional to global scales. The system studied here has at least four  
480 external processes that may influence the positive feedback we documented. First, land use  
481 change via livestock grazing facilitates invasion ([Ponzetti, Mccune, and Pyke 2007](#); [Williamson  
et al. 2019](#)). Second, the introduction of exotic grasses increases fuel connectivity ([Davies  
and Nafus 2013](#)), affects burn severity. Third, increasing temperatures due to climate change  
484 increase burn severity in forests ([S. A. Parks and Abatzoglou 2020](#)). We expect this to be  
485 true for shrublands, and is an important area for future research. Increasing temperatures  
486 simultaneously decrease seed viability and seedling survival ([Schlaepfer, Lauenroth, and  
Bradford 2014](#); [Enright et al. 2015](#)). Fourth, CO<sub>2</sub> enrichment may preferentially enhance  
488 biomass (i.e. higher fuel connectivity) and seed production of annual grass species ([Smith  
et al. 2000](#); [Nagel et al. 2004](#)). All four of these external drivers are globally ubiquitous  
490 consequences of global change.

491 An ecosystem “state” is the product of countless endogenous interactions. The grass-fire  
492 cycle studied here is strengthened through providing fitness benefits to the introduced annual  
493 grasses via at least three reinforcing processes. First, we document how it changes the  
494 composition of the seedbank. Second, introduced annual grasses competitively exclude native  
495 plants. Third, the dominance of introduced annual grasses initiates ecohydrological feedbacks  
496 to create a warmer, drier microclimate ([Turnbull et al. 2012](#)). It is possible that some  
497 of these feedbacks are idiosyncratic to the system being studied, while others may reflect  
498 fundamental properties of ecosystem function that change when a system is converted from  
499 being dominated by deep-rooted woody plants to being dominated by annual herbaceous  
500 plants ([Kitzberger et al. 2016](#)). At least 13 grass species initiate self-reinforcing feedbacks

501 with fire in the U.S. alone (Fusco et al. 2019; Tortorelli, Krawchuk, and Kerns 2020). There  
502 are many more fire-inducing grass invasions worldwide, with documented cases in Australia  
503 (G. Miller et al. 2010), Brazil (Rossi et al. 2014) and South Africa (Milton 2004). The  
504 conversion of forests and shrublands to grasslands may have consequences relevant to the  
505 global carbon cycle, especially when ecosystems dominated by deep-rooted plants that store  
506 carbon belowground are replaced by shallow-rooted ecosystems that lose carbon to grazing  
507 and fire (Kerns et al. 2020; Mahood et al. 2022).

## 508 Acknowledgements

509 We thank Abdelhakim Farid, Julia Lopez, Dylan Murphy and C. Nick Whittemore for help  
510 in the field and in the greenhouse, and Lindsay P. Chiquoine, Thomas T. Veblen and two  
511 anonymous reviewers for constructive feedback that greatly improved the manuscript. We  
512 are grateful to everyone in the Winnemucca office of the Bureau of Land Management, the  
513 Central Nevada Interagency Dispatch Center, and CU Boulder's Ecology Evolution and  
514 Biology Greenhouse. This project was funded in part by the CU Boulder Undergraduate  
515 Research Opportunities Program and the Geography department's Adam Kolff Memorial  
516 Graduate Research Grant.

## 517 References

- 518 Aleman, J. C., A. Fayolle, C. Favier, A. C. Staver, K. G. Dexter, C. M. Ryan, A. F.  
519 Azihou, et al. 2020. "Floristic Evidence for Alternative Biome States in Tropical  
520 Africa." *Proceedings of the National Academy of Sciences* 117 (45): 28183–90. <https://doi.org/10.1073/pnas.2011515117>.
- 522 Allan, Eric, Pete Manning, Fabian Alt, Julia Binkenstein, Stefan Blaser, Nico Blüthgen,  
523 Stefan Böhm, et al. 2015. "Land Use Intensification Alters Ecosystem Multifunctionality  
524 via Loss of Biodiversity and Changes to Functional Composition." *Ecology Letters* 18 (8):

- 525 834–43. <https://doi.org/10.1111/ele.12469>.
- 526 Archibald, Sally, A. Carla Staver, and Simon A. Levin. 2012. “Evolution of Human-Driven  
527 Fire Regimes in Africa.” *Proceedings of the National Academy of Sciences* 109 (3): 847–52.  
528 <https://doi.org/10.1073/pnas.1118648109>.
- 529 Arkle, Robert S., David S. Pilliod, Matthew J. Germino, Michelle I. Jeffries, and Justin  
530 L. Welty. 2022. “Reestablishing a Foundational Species: Limitations on Post-wildfire  
531 Sagebrush Seedling Establishment.” *Ecosphere* 13 (8). <https://doi.org/10.1002/ecs2.4195>.
- 532 Balch, Jennifer K., Bethany A. Bradley, Carla M. D’Antonio, and José Gómez-Dans. 2013.  
533 “Introduced annual grass increases regional fire activity across the arid western USA  
534 (1980-2009).” *Global Change Biology* 19 (1): 173–83. <https://doi.org/10.1111/gcb.12046>.
- 535 Bechtold, H. A., and R. S. Inouye. 2007. “Distribution of carbon and nitrogen in sage-  
536 brush steppe after six years of nitrogen addition and shrub removal.” *Journal of Arid  
537 Environments* 71 (1): 122–32. <https://doi.org/10.1016/j.jaridenv.2007.02.004>.
- 538 Beckstead, Julie, Laura E. Street, Susan E. Meyer, and Phil S. Allen. 2011. “Fire effects on  
539 the cheatgrass seed bank pathogen Pyrenophora semeniperda.” *Rangeland Ecology and  
540 Management* 64 (2): 148–57. <https://doi.org/10.2111/REM-D-10-00052.1>.
- 541 Bond, William J., and Jeremy J. Midgley. 1995. “Kill Thy Neighbour: An Individualistic  
542 Argument for the Evolution of Flammability.” *Oikos* 73 (1): 79. <https://doi.org/10.2307/3545728>.
- 543
- 544 Boudell, JA, SO Link, and JR Johansen. 2002. “Effect of soil microtopography on seed  
545 bank distribution in the shrub-steppe.” *Western North American Naturalist* 62 (1): 14–24.  
546 <https://doi.org/10.2307/41717153>.
- 547 Bradford, John B, Julio L Betancourt, Bradley J Butterfield, Seth M Munson, and Troy E  
548 Wood. 2018. “Anticipatory Natural Resource Science and Management for a Changing  
549 Future.” *Frontiers in Ecology and the Environment* 16 (5): 295–303. <https://doi.org/10.1002/fee.1806>.
- 550
- 551 Brooks, Matthew L., Carla M. D’Antonio, David M. Richardson, James B. Grace, Jon E.

- 552 Keeley, Joseph M. DiTomaso, Richard J. Hobbs, Mike Pellant, and David Pyke. 2004.
- 553 “Effects of Invasive Alien Plants on Fire Regimes.” *BioScience* 54 (7): 677–88.
- 554 Bukowski, Beth, and William L. Baker. 2013. “Historical fire regimes, reconstructed from
- 555 land-survey data, led to complexity and fluctuation in sagebrush landscapes.” *Ecological*
- 556 *Applications* 23 (3): 546–64.
- 557 Commission for Environmental Cooperation. 2006. “Ecological regions of North America –
- 558 Levels I, II, and III: Montreal, Quebec, Canada, Commission for Environmental Cooper-
- 559 ation, scale 1:10,000,000.” <https://www.epa.gov/eco-research/ecoregions-north-america>.
- 560 Condon, Lea A., and David A. Pyke. 2018. “Fire and Grazing Influence Site Resistance to
- 561 Bromus tectorum Through Their Effects on Shrub, Bunchgrass and Biocrust Communities
- 562 in the Great Basin (USA).” *Ecosystems* 21 (7): 1416–31. <https://doi.org/10.1007/s10021-018-0230-8>.
- 563
- 564 D’Antonio, Carla M., and Peter M. Vitousek. 1992. “Biological invasions by exotic grasses,
- 565 the grass/fire cycle, and global change.” *Annual Review of Ecological Systems* 23: 63–87.
- 566 Davies, Kirk W., Jonathan D. Bates, Tony J. Svejcar, and Chad S. Boyd. 2010. “Effects
- 567 of long-term livestock grazing on fuel characteristics in rangelands: An example from
- 568 the sagebrush steppe.” *Rangeland Ecology and Management* 63 (6): 662–69. <https://doi.org/10.2111/REM-D-10-00006.1>.
- 569
- 570 Davies, Kirk W., Chad S. Boyd, Jeffrey L. Beck, Jon D. Bates, Tony J. Svejcar, and
- 571 Michael A. Gregg. 2011. “Saving the sagebrush sea: An ecosystem conservation plan
- 572 for big sagebrush plant communities.” *Biological Conservation* 144 (11): 2573–84. <https://doi.org/10.1016/j.biocon.2011.07.016>.
- 573
- 574 Davies, Kirk W., and Aleta M. Nafus. 2013. “Exotic annual grass invasion alters fuel amounts,
- 575 continuity and moisture content.” *International Journal of Wildland Fire* 22 (3): 353–58.
- 576 <https://doi.org/10.1071/WF11161>.
- 577
- 578 Davis, Kimberley T., Solomon Z. Dobrowski, Philip E. Higuera, Zachary A. Holden, Thomas
- T. Veblen, Monica T. Rother, Sean A. Parks, Anna Sala, and Marco P. Maneta. 2019.

- 579        “Wildfires and Climate Change Push Low-Elevation Forests Across a Critical Climate  
580        Threshold for Tree Regeneration.” *Proceedings of the National Academy of Sciences*,  
581        201815107. <https://doi.org/10.1073/pnas.1815107116>.
- 582        Davis, Kimberley T., Philip E. Higuera, and Anna Sala. 2018. “Anticipating Fire-mediated  
583        Impacts of Climate Change Using a Demographic Framework.” Edited by Charles Fox.  
584        *Functional Ecology* 32 (7): 1729–45. <https://doi.org/10.1111/1365-2435.13132>.
- 585        Doescher, Paul S., Richard F. Miller, and Alma H. Winward. 1984. “Soil Chemical Patterns  
586        under Eastern Oregon Plant Communities Dominated by Big Sagebrush.” <https://doi.org/10.2136/sssaj1984.03615995004800030038x>.
- 588        Eckert, Richard E., Frederick F. Peterson, Michael S. Meurisse, and L. Stephens. 1986.  
589        “Effects of Soil-Surface Morphology on Emergence and Survival of Seedlings in Big  
590        Sagebrush Communities.” *Journal of Range Management* 39 (5): 414–20. <http://www.jstor.org/stable/3899441>.
- 592        Eidenshink, Jeff, Brian Schwind, Ken Brewer, Zhi-liang Zhu, Brad Quayle, and Stephen  
593        Howard. 2007. “A Project for Monitoring Trends in Burn Severity.” *Fire Ecology* 3 (1):  
594        3–21. <https://doi.org/10.4996/firecology.0301003>.
- 595        Elzinga, Caryl L, Daniel W Salzer, and John W Willoughby. 1998. “Measuring & Monitoring  
596        Plant Populations.” BLM Technical Reference 1730-1. Bureau of Land Management.
- 597        Enright, Neal J., Joseph B. Fontaine, David M. J. S. Bowman, Ross A. Bradstock, and  
598        Richard J. Williams. 2015. “Interval Squeeze: Altered Fire Regimes and Demographic  
599        Responses Interact to Threaten Woody Species Persistence as Climate Changes.” *Frontiers  
600        in Ecology and the Environment* 13 (5): 265–72. <https://doi.org/10.1890/140231>.
- 601        Eskelinen, Anu, Kelly Gravuer, W Stanley Harpole, Susan Harrison, Risto Virtanen, and  
602        Yann Hautier. 2020. “Resource-enhancing Global Changes Drive a Whole-ecosystem Shift  
603        to Faster Cycling but Decrease Diversity.” *Ecology* 101 (12). <https://doi.org/10.1002/ecy.3178>.
- 605        Farrell, Hannah L., Jeffrey S. Fehmi, and Elise S. Gornish. 2021. “The Effects of Site

- 606 Preparation Equal Those of Seeding at a Dryland Restoration Site: 6 Years of Plant  
607 Community Development." *Restoration Ecology* 29 (6). <https://doi.org/10.1111/rec.13482>.
- 608 Fenesi, Annamária, Sandra Saura-Mas, Robert R. Blank, Anita Kozma, Beáta-Magdolna  
609 Lózer, and Eszter Ruprecht. 2016. "Enhanced Fire-Related Traits May Contribute to  
610 the Invasiveness of Downy Brome (*Bromus Tectorum*)."*Invasive Plant Science and*  
611 *Management* 9 (3): 182–94. <https://doi.org/10.1614/IPSM-D-16-00006.1>.
- 612 Fusco, Emily J., John T. Finn, Jennifer K. Balch, R. Chelsea Nagy, and Bethany A. Bradley.  
613 2019. "Invasive Grasses Increase Fire Occurrence and Frequency Across US Ecoregions."  
614 *Proceedings of the National Academy of Sciences* 116 (47): 23594–99. <https://doi.org/10.1073/pnas.1908253116>.
- 615
- 616 Gagnon, Paul R., Heather A. Passmore, Matthew Slocum, Jonathan A. Myers, Kyle E.  
617 Harms, William J. Platt, and C. E. Timothy Paine. 2015. "Fuels and fires influence  
618 vegetation via above- and belowground pathways in a high-diversity plant community."  
619 *Journal of Ecology* 103 (4): 1009–19. <https://doi.org/10.1111/1365-2745.12421>.
- 620 Gelman, Andrew, Donald B Rubin, et al. 1992. "Inference from Iterative Simulation Using  
621 Multiple Sequences." *Statistical Science* 7 (4): 457–72.
- 622 Germino, Matthew J., Jeanne C. Chambers, and Cynthia S. Brown. 2016. *Exotic Brome-  
623 Grasses in Arid and Semiarid Ecosystems of the Western US Causes, Consequences, and  
624 Management Implications*. <http://www.springer.com/series/412>.
- 625 Gibbens, Robert P., and James M. Lenz. 2001. "Root systems of some Chihuahuan Desert  
626 plants." *Journal of Arid Environments* 49: 221–63.
- 627 Gill, Nathan S, Monica G Turner, Carissa D Brown, Sydney I Glassman, Sandra L Haire,  
628 Winslow D Hansen, Elizabeth R Pansing, Samuel B St Clair, and Diana F Tomback. 2022.  
629 "Limitations to Propagule Dispersal Will Constrain Postfire Recovery of Plants and Fungi  
630 in Western Coniferous Forests." *BioScience*, January, biab139. <https://doi.org/10.1093/biosci/biab139>.
- 631
- 632 Hardegree, Stuart P., John T. Abatzoglou, Mark W. Brunson, Matthew J. Germino, Katherine

- 633 C. Hegewisch, Corey A. Moffet, David S. Pilliod, Bruce A. Roundy, Alex R. Boehm,  
634 and Gwendwr R. Meredith. 2018. "Weather-Centric Rangeland Revegetation Planning."  
635 *Rangeland Ecology & Management* 71 (1): 1–11. <https://doi.org/10.1016/j.rama.2017.07.003>.
- 636
- 637 Hassan, M. A., and N. E. West. 1986. "Dynamics of Soil Seed Pools in Burned and Unburned  
638 Sagebrush Semi-Deserts." *Ecology* 67 (1): 269–72.
- 639 Heerdt, G. N. J. Ter, G. L. Verweij, R. M. Bekker, and J. P. Bakker. 1996. "An Improved  
640 Method for Seed-Bank Analysis: Seedling Emergence After Removing the Soil by Sieving."  
641 *Functional Ecology* 10 (1): 144. <https://doi.org/10.2307/2390273>.
- 642 Heydari, Mehdi, Reza Omidipour, Mehdi Abedi, and Carol Baskin. 2017. "Effects of  
643 fire disturbance on alpha and beta diversity and on beta diversity components of soil  
644 seed banks and aboveground vegetation." *Plant Ecology and Evolution* 150 (3): 247–56.  
645 <https://doi.org/10.5091/plecevo.2017.1344>.
- 646 Hirota, Marina, Milena Holmgren, Egbert H. Van Nes, and Marten Scheffer. 2011. "Global  
647 resilience of tropical forest and savanna to critical transitions." *Science* 334 (6053): 232–35.  
648 <https://doi.org/10.1126/science.1210657>.
- 649 Humphrey, L David, and Eugene W Schupp. 2001. "Seed banks of Bromus tectorum-  
650 dominated communities in the Great Basin." *Western North American Naturalist* 61 (1):  
651 85–92. <https://doi.org/10.2307/41717080>.
- 652 Jones, Matthew O., Nathaniel P. Robinson, David E. Naugle, Jeremy D. Maestas, Matthew  
653 C. Reeves, Robert W. Lankston, and Brady W. Allred. 2021. "Annual and 16-Day  
654 Rangeland Production Estimates for the Western United States." *Rangeland Ecology &*  
655 *Management* 77 (July): 112–17. <https://doi.org/10.1016/j.rama.2021.04.003>.
- 656 Jones, Rachel O., Jeanne C. Chambers, David I. Board, Dale W. Johnson, and Robert R.  
657 Blank. 2015. "The role of resource limitation in restoration of sagebrush ecosystems  
658 dominated by cheatgrass (Bromus tectorum)." *Ecosphere* 6 (7): 1–21.
- 659 Keddy, Paul A, and Daniel C Laughlin. 2021. *A Framework for Community Ecology: Species*

- 660        *Pools, Filters and Traits*. Cambridge University Press.
- 661    Keeley, Jon E. 2009. "Fire intensity, fire severity and burn severity: A brief review and  
662        suggested usage." *International Journal of Wildland Fire* 18 (1): 116–26. <https://doi.org/10.1071/WF07049>.
- 663
- 664    Keeley, Jon E., Juli G. Pausas, Philip W. Rundel, William J. Bond, and Ross A. Bradstock.  
665        2011. "Fire as an evolutionary pressure shaping plant traits." *Trends in Plant Science* 16  
666        (8): 406–11. <https://doi.org/10.1016/j.tplants.2011.04.002>.
- 667    Kerns, Becky K., Claire Tortorelli, Michelle A. Day, Ty Nietupski, Ana M. G. Barros, John  
668        B. Kim, and Meg A. Krawchuk. 2020. "Invasive grasses: A new perfect storm for  
669        forested ecosystems?" *Forest Ecology and Management* 463 (November 2019): 117985.  
670        <https://doi.org/10.1016/j.foreco.2020.117985>.
- 671    Key, Carl H, and Nathan C Benson. 1999. "The Composite Burn Index (CBI): Field Rating  
672        of Burn Severity." *USGS, NRMSC Research,[online]* Available: [Http://Nrmsc. Usgs. Gov/Research/Cbi. Htm \[3/14/2006\]](Http://Nrmsc. Usgs. Gov/Research/Cbi. Htm [3/14/2006]).
- 673
- 674    Kimura, Hideo, and Shiro Tsuyuzaki. 2011. "Fire severity affects vegetation and seed bank  
675        in a wetland." *Applied Vegetation Science* 14 (3): 350–57. <https://doi.org/10.1111/j.1654-109X.2011.01126.x>.
- 676
- 677    Kitzberger, Thomas, G. L. W. Perry, J. Paritsis, J. H. Gowda, A. J. Tepley, A. Holz, and T.  
678        T. Veblen. 2016. "Fire–vegetation feedbacks and alternative states: common mechanisms  
679        of temperate forest vulnerability to fire in southern South America and New Zealand."  
680        *New Zealand Journal of Botany* 54 (2): 247–72. <https://doi.org/10.1080/0028825X.2016.1151903>.
- 681
- 682    Kolden, Crystal A, Alistair M S Smith, and John T. Abatzoglou. 2015. "Limitations and  
683        utilisation of Monitoring Trends in Burn Severity products for assessing wildfire severity  
684        in the USA." *International Journal of Wildland Fire* 24: 1023–28.
- 685
- 686    Koontz, Michael J, Malcolm P North, Chhaya M Werner, Stephen E Fick, and Andrew M  
Latimer. 2020. "Local Forest Structure Variability Increases Resilience to Wildfire in Dry

- 687 Western US Coniferous Forests." *Ecology Letters* 23 (3): 483–94.
- 688 Lipoma, M. Lucrecia, Guillermo Funes, and Sandra Díaz. 2018. "Fire effects on the soil  
689 seed bank and post-fire resilience of a semi-arid shrubland in central Argentina." *Austral  
690 Ecology* 43 (1): 46–55. <https://doi.org/10.1111/aec.12533>.
- 691 Liyanage, Ganesha S., and Mark K. J. Ooi. 2017. "Do dormancy-breaking temperature  
692 thresholds change as seeds age in the soil seed bank?" *Seed Science Research* 27 (1): 1–11.  
693 <https://doi.org/10.1017/S0960258516000271>.
- 694 Mahood, Adam L., and Jennifer K. Balch. 2019. "Repeated fires reduce plant diversity  
695 in low-elevation Wyoming big sagebrush ecosystems (1984 – 2014)." *Ecosphere* 10 (2):  
696 e02591. <https://doi.org/10.1002/ecs2.2591>.
- 697 Mahood, Adam L., Rachel O. Jones, David I. Board, Jennifer K. Balch, and Jeanne C.  
698 Chambers. 2022. "Interannual Climate Variability Mediates Changes in Carbon and  
699 Nitrogen Pools Caused by Annual Grass Invasion in a Semiarid Shrubland." *Global Change  
700 Biology* 28 (1): 267–84. <https://doi.org/10.1111/gcb.15921>.
- 701 Maia, P., J. G. Pausas, V. Arcenegui, C. Guerrero, A. Pérez-Bejarano, J. Mataix-Solera, M.  
702 E. T. Varela, I. Fernandes, E. T. Pedrosa, and J. J. Keizer. 2012. "Wildfire effects on the  
703 soil seed bank of a maritime pine stand - The importance of fire severity." *Geoderma* 191:  
704 80–88. <https://doi.org/10.1016/j.geoderma.2012.02.001>.
- 705 Meyer, Susan E, Stephen B Monsen, and E Durant Mcarthur. 2013. "Germination Response  
706 of Artemisia tridentata (Asteraceae) to Light and Chill: Patterns of Between-Population  
707 Variation." *Botanical Gazette* 151 (2): 176–83.
- 708 Miller, Georgia, Margaret Friedel, Paul Adam, and Vanessa Chewings. 2010. "Ecological  
709 impacts of buffel grass (*Cenchrus ciliaris* L.) invasion in central Australia does field  
710 evidence support a fire-invasion feedback?" *The Rangeland Journal* 32 (4): 353–65.  
711 <https://doi.org/10.1071/RJ09076>.
- 712 Miller, Jay D., Eric E. Knapp, Carl H. Key, Carl N. Skinner, Clint J. Isbell, R. Max Creasy,  
713 and Joseph W. Sherlock. 2009. "Calibration and validation of the relative differenced

- 714      Normalized Burn Ratio (RdNBR) to three measures of fire severity in the Sierra Nevada  
715      and Klamath Mountains, California, USA.” *Remote Sensing of Environment* 113 (3):  
716      645–56. <https://doi.org/10.1016/j.rse.2008.11.009>.
- 717      Milton, Sue J. 2004. “Grasses as invasive alien plants in South Africa.” *South African Journal*  
718      *of Science* 100 (1-2): 69–75.
- 719      Monty, Arnaud, Cynthia S. Brown, and Danielle B. Johnston. 2013. “Fire promotes downy  
720      brome (*Bromus tectorum* L.) seed dispersal.” *Biological Invasions* 15 (5): 1113–23.  
721      <https://doi.org/10.1007/s10530-012-0355-1>.
- 722      Morgan, P., and L. F. Neuenschwander. 1988. “Seed-Bank Contributions to Regeneration of  
723      Shrub Species After Clear-Cutting and Burning.” *Canadian Journal of Botany* 66 (1):  
724      169–72. <https://doi.org/10.1139/b88-026>.
- 725      Morris, Lesley R., and Elizabeth A. Leger. 2016. “Secondary Succession in the Sagebrush  
726      Semidesert 66 Years After Fire in the Great Basin, USA.” *Natural Areas Journal* 36 (2):  
727      187–93. <https://doi.org/10.3375/043.036.0211>.
- 728      Nagel, Jennifer M., Travis E. Huxman, Kevin L. Griffin, and Stanley D. Smith. 2004. “CO<sub>2</sub>  
729      enrichment reduces the energetic cost of biomass construction in an invasive desert grass.”  
730      *Ecology* 85 (1): 100–106. <https://doi.org/10.1890/02-3005>.
- 731      Naghipour, Ali Asghar, Hossein Bashari, Seyed Jamaleddin Khajeddin, Pejman Tahmasebi,  
732      and Majid Iravani. 2016. “Effects of Smoke, Ash and Heat Shock on Seed Germination of  
733      Seven Species from Central Zagros Rangelands in the Semi-Arid Region of Iran.” *African*  
734      *Journal of Range & Forage Science* 33 (1): 67–71. <https://doi.org/10.2989/10220119.2015.1119194>.
- 736      Ottaviani, Gianluigi, Rafael Molina-Venegas, Tristan Charles-Dominique, Stefano Chelli,  
737      Giandiego Campetella, Roberto Canullo, and Jitka Klimešová. 2020. “The Neglected  
738      Belowground Dimension of Plant Dominance.” *Trends in Ecology and Evolution* 35 (9):  
739      763–66. <https://doi.org/10.1016/j.tree.2020.06.006>.
- 740      Ottmar, Roger D, David V Sandberg, Cynthia L Riccardi, and Susan J Prichard. 2007. “An

- 741 Overview of the Fuel Characteristic Classification System—Quantifying, Classifying, and  
742 Creating Fuelbeds for Resource Planning.” *Canadian Journal of Forest Research* 37 (12):  
743 2383–93.
- 744 Owens, M. K., and B. E. Norton. 1992. “Interactions of Grazing and Plant Protection on  
745 Basin Big Sagebrush (*Artemisia tridentata* ssp . *tridentata*) Seedling Survival.” *Journal*  
746 *of Range Management* 45 (3): 257–62. <http://www.jstor.org/stable/4002974>.
- 747 Palmer, Harrison D., Andrew J. Denham, and Mark K. J. Ooi. 2018. “Fire severity drives  
748 variation in post-fire recruitment and residual seed bank size of Acacia species.” *Plant*  
749 *Ecology* 219 (5): 527–37. <https://doi.org/10.1007/s11258-018-0815-5>.
- 750 Parks, S. A., and J. T. Abatzoglou. 2020. “Warmer and Drier Fire Seasons Contribute to  
751 Increases in Area Burned at High Severity in Western US Forests From 1985 to 2017.”  
752 *Geophysical Research Letters* 47 (22). <https://doi.org/10.1029/2020GL089858>.
- 753 Parks, Sean A., Lisa M. Holsinger, Morgan A. Voss, Rachel A. Loehman, and Nathaniel  
754 P. Robinson. 2018. “Mean composite fire severity metrics computed with google earth  
755 engine offer improved accuracy and expanded mapping potential.” *Remote Sensing* 10  
756 (6): 1–15. <https://doi.org/10.3390/rs10060879>.
- 757 Pedrini, Simone, Alma Balestrazzi, Matthew D. Madsen, Khiraj Bhalsing, Stuart P. Hardegree,  
758 Kingsley W. Dixon, and Olga A. Kildisheva. 2020. “Seed Enhancement: Getting Seeds  
759 Restoration-ready.” *Restoration Ecology* 28 (S3). <https://doi.org/10.1111/rec.13184>.
- 760 Petraitis, Peter S., and Roger Earl Latham. 1999. “The importance of scale in testing the  
761 origins of alternative community states.” *Ecology* 80 (2): 429–42. [https://doi.org/10.1890/0012-9658\(1999\)080%5B0429:TIOSIT%5D2.0.CO;2](https://doi.org/10.1890/0012-9658(1999)080%5B0429:TIOSIT%5D2.0.CO;2).
- 763 Ponzetti, J. M., B. Mccune, and David A. Pyke. 2007. “Biotic Soil Crusts in Relation to  
764 Topography, Cheatgrass and Fire in the Columbia Basin, Washington.” *The Bryologist* 110  
765 (4): 706–22. [https://doi.org/10.1639/0007-2745\(2007\)110%5B706:BSCIRT%5D2.0.CO;2](https://doi.org/10.1639/0007-2745(2007)110%5B706:BSCIRT%5D2.0.CO;2).
- 766 Pyke, David A., Robert K. Shriver, Robert S. Arkle, David S. Pilliod, Cameron L. Aldridge,  
767 Peter S. Coates, Matthew J. Germino, Julie A. Heinrichs, Mark A. Ricca, and Scott E.

- 768 Shaff. 2020. "Postfire Growth of Seeded and Planted Big Sagebrush—Strategic Designs for  
769 Restoring Greater Sage-grouse Nesting Habitat." *Restoration Ecology* 28 (6): 1495–1504.  
770 <https://doi.org/10.1111/rec.13264>.
- 771 R Core Team. 2020. *R: A Language and Environment for Statistical Computing*. Vienna,  
772 Austria: R Foundation for Statistical Computing. <https://www.R-project.org/>.
- 773 Ratajczak, Zak, Stephen R. Carpenter, Anthony R. Ives, Christopher J. Kucharik, Tanjona  
774 Ramiadantsoa, M. Allison Stegner, John W. Williams, Jien Zhang, and Monica G. Turner.  
775 2018. "Abrupt Change in Ecological Systems: Inference and Diagnosis." *Trends in Ecology  
776 and Evolution* 33 (7): 513–26. <https://doi.org/10.1016/j.tree.2018.04.013>.
- 777 Rosseel, Yves. 2012. "lavaan: An R Package for Structural Equation Modeling." *Journal of  
778 Statistical Software* 48 (2): 1–36. <https://doi.org/10.18637/jss.v048.i02>.
- 779 Rossi, Rafael Drumond, Carlos Romero Martins, Pedro Lage Viana, Evandro Luís Rodrigues,  
780 and José Eugênio Côrtes Figueira. 2014. "Impact of invasion by molasses grass (*Melinis  
781 minutiflora* P. Beauv.) on native species and on fires in areas of campo-cerrado in Brazil."  
782 *Acta Botanica Brasilica* 28 (4): 631–37. <https://doi.org/10.1590/0102-33062014abb3390>.
- 783 Rother, Monica T., Thomas T. Veblen, and Luke G. Furman. 2015. "A Field Experiment  
784 Informs Expected Patterns of Conifer Regeneration After Disturbance Under Changing  
785 Climate Conditions." *Canadian Journal of Forest Research* 45 (11): 1607–16. <https://doi.org/10.1139/cjfr-2015-0033>.
- 787 Scheffer, Marten, and Stephen R. Carpenter. 2003. "Catastrophic Regime Shifts in Ecosys-  
788 tems: Linking Theory to Observation." *Trends in Ecology & Evolution* 18 (12): 648–56.  
789 <https://doi.org/10.1016/j.tree.2003.09.002>.
- 790 Scheffer, Marten, Stephen R. Carpenter, Vasilis Dakos, and Egbert H. van Nes. 2015. "Generic  
791 Indicators of Ecological Resilience: Inferring the Chance of a Critical Transition." *Annual  
792 Review of Ecology, Evolution, and Systematics* 46 (1): 145–67. <https://doi.org/10.1146/annurev-ecolsys-112414-054242>.
- 794 Scheffer, M, S Carpenter, JA Foley, C Folke, and B Walker. 2001. "Catastrophic Shifts in

- 795 Ecosystems." *Nature* 413: 591–96.
- 796 Schimmel, Johnny, and Anders Granström. 1996. "Fire Severity and Vegetation Response in  
797 the Boreal Swedish Forest." *Ecology* 77 (5): 1436–50.
- 798 Schlaepfer, Daniel R., William K. Lauenroth, and John B. Bradford. 2014. "Natural  
799 Regeneration Processes in Big Sagebrush (*Artemisia tridentata*)."*Rangeland Ecology &*  
800 *Management* 67 (4): 344–57. <https://doi.org/10.2111/REM-D-13-00079.1>.
- 801 Schröder, Arne, Lennart Persson, and André M. De Roos. 2005. "Direct Experimental  
802 Evidence for Alternative Stable States: A Review." *Oikos* 110 (1): 3–19. <https://doi.org/10.1111/j.0030-1299.2005.13962.x>.
- 803
- 804 Shannon, CE, and W Weaver. 1949. "The Mathematical Theory of Communication. University  
805 of Illinois Press, Urbana-Champaign, Illinois, USA, 117 p."
- 806 Shinneman, Douglas J., and Susan K. McIlroy. 2016. "Identifying key climate and environ-  
807 mental factors affecting rates of post-fire big sagebrush (*Artemisia tridentata*) recovery in  
808 the northern Columbia Basin, USA." *International Journal of Wildland Fire* 25: 933–45.  
809 <https://doi.org/10.1071/WF16013>.
- 810 Shriner, Robert K., Caitlin M. Andrews, Robert S. Arkle, David M. Barnard, Michael C.  
811 Duniway, Matthew J. Germino, David S. Pilliod, David A. Pyke, Justin L. Welty, and  
812 John B. Bradford. 2019. "Transient Population Dynamics Impede Restoration and May  
813 Promote Ecosystem Transformation After Disturbance." Edited by Shaopeng Wang.  
814 *Ecology Letters* 22 (9): 1357–66. <https://doi.org/10.1111/ele.13291>.
- 815 Shriner, Robert K., Caitlin M. Andrews, David S. Pilliod, Robert S. Arkle, Justin L. Welty,  
816 Matthew J. Germino, Michael C. Duniway, David A. Pyke, and John B. Bradford.  
817 2018. "Adapting Management to a Changing World: Warm Temperatures, Dry Soil,  
818 and Interannual Variability Limit Restoration Success of a Dominant Woody Shrub in  
819 Temperate Drylands." *Global Change Biology* 24 (10): 4972–82. <https://doi.org/10.1111/gcb.14374>.
- 820
- 821 Smith, Stanley D., Travis E. Huxman, Stephen F. Zitzer, Therese N. Charlet, David C.

- 822 Housman, James S. Coleman, Lynn K. Fenstermaker, Jeffrey R. Seemann, and Robert S.  
823 Nowak. 2000. “Elevated CO<sub>2</sub> increases productivity and invasive species success in an  
824 arid ecosystem.” *Nature* 408 (6808): 79–82. <https://doi.org/10.1038/35040544>.
- 825 Staal, Arie, Ingo Fetzer, Lan Wang-Erlandsson, Joyce H. C. Bosmans, Stefan C. Dekker,  
826 Egbert H. van Nes, Johan Rockström, and Obbe A. Tuinenburg. 2020. “Hysteresis  
827 of tropical forests in the 21st century.” *Nature Communications* 11 (1): 1–8. <https://doi.org/10.1038/s41467-020-18728-7>.
- 829 Staver, A. Carla, Sally Archibald, and Simon A. Levin. 2011. “The global extent and  
830 determinants of savanna and forest as alternative biome states.” *Science* 334 (6053):  
831 230–32. <https://doi.org/10.1126/science.1210465>.
- 832 Stavros, E. Natasha, Janice Coen, Birgit Peterson, Harshvardhan Singh, Kama Kennedy,  
833 Carlos Ramirez, and David Schimel. 2018. “Use of Imaging Spectroscopy and LIDAR to  
834 Characterize Fuels for Fire Behavior Prediction.” *Remote Sensing Applications: Society*  
835 *and Environment* 11: 41–50. <https://doi.org/10.1016/j.rse.2018.04.010>.
- 836 Steenvoorden, Jasper, Arjan J. H. Meddens, Anthony J. Martinez, Lee J. Foster, and W.  
837 Daniel Kissling. 2019. “The potential importance of unburned islands as refugia for the  
838 persistence of wildlife species in fire-prone ecosystems.” *Ecology and Evolution* 9 (15):  
839 8800–8812. <https://doi.org/10.1002/ece3.5432>.
- 840 Suding, Katharine N., Katherine L. Gross, and Gregory R. Houseman. 2004. “Alternative  
841 states and positive feedbacks in restoration ecology.” *Trends in Ecology & Evolution* 19  
842 (1): 46–53. <https://doi.org/10.1016/j.tree.2003.10.005>.
- 843 Tikhonov, Gleb, Otso Ovaskainen, Jari Oksanen, Melinda de Jonge, Oystein Opedal, and  
844 Tad Dallas. 2020. *Hmsc: Hierarchical Model of Species Communities*. <https://CRAN.R-project.org/package=Hmsc>.
- 846 Tortorelli, Claire M., Meg A. Krawchuk, and Becky K. Kerns. 2020. “Expanding the  
847 invasion footprint: Ventenata dubia and relationships to wildfire, environment, and plant  
848 communities in the Blue Mountains of the Inland Northwest, USA.” *Applied Vegetation*

- 849        *Science*, no. May: 1–13. <https://doi.org/10.1111/avsc.12511>.
- 850    Turnbull, Laura, Bradford P. Wilcox, J. Benlap, S. Ravi, P. D'Odorico, D. Childers, W.  
851        Gwenzi, et al. 2012. “Understanding the role of ecohydrological feedbacks in ecosystem  
852        state change in drylands.” *Ecohydrology* 5: 174–83. <https://doi.org/10.1002/eco>.
- 853    U.S. Department of Interior, Bureau of Land Management (BLM). 2018. “BLM AIM  
854        TerrADat TerrestrialAIM point.” BLM National Operations Center: BLM. [https://gis.blm.gov/AIMdownload/layerpackages/BLM\\_AIM\\_Terrestrial.lpk](https://gis.blm.gov/AIMdownload/layerpackages/BLM_AIM_Terrestrial.lpk).
- 855        Williamson, Matthew A., Erica Fleishman, Ralph C. Mac Nally, Jeanne C. Chambers,  
856        Bethany A. Bradley, David S. Dobkin, David I. Board, et al. 2019. “Fire, livestock  
857        grazing, topography, and precipitation affect occurrence and prevalence of cheatgrass  
858        (*Bromus tectorum*) in the central Great Basin, USA.” *Biological Invasions* 22 (2): 663–80.  
859        <https://doi.org/10.1007/s10530-019-02120-8>.
- 860        Wright, Boyd R., Peter K. Latz, and A. F. Zuur. 2016. “Fire severity mediates seedling  
861        recruitment patterns in slender mulga (*Acacia aptaneura*), a fire-sensitive Australian  
862        desert shrub with heat-stimulated germination.” *Plant Ecology* 217 (6): 789–800. <https://doi.org/10.1007/s11258-015-0550-0>.

865 **Figure Captions**

866 **Figure 1.** Visual illustration of the relationship between fuel connectivity and burn severity.

867 On the left, panel a shows the inter-shrub space invaded by annual grasses. The photo in  
868 panel b was taken in the exact same place two weeks later, days after all of the biomass was  
869 consumed by the fire. Panel C is a closeup of the soil surface, showing in more detail how the  
870 litter was also almost completely consumed by the fire. On the right, the photos in panels d  
871 and e were on opposite sides of a fire line in an area that had minimal annual grass invasion  
872 over a broad area, and thus lower fuel connectivity. Note the remaining plants and stumps in  
873 panel e and the presence of only partially consumed litter in panel f.

874 **Figure 2.** Sites with little to no shrub cover require high IAG cover to meet the threshold  
875 necessary to carry a fire, while sites with higher shrub cover may reach that threshold  
876 with much lower IAG cover. Therefore, annual grass cover alone may not be sufficient for  
877 quantifying fire risk. Panel a illustrates this point using publicly available data from the  
878 Bureau of Land Management's Assessment, Inventory and Monitoring dataset. Panels b and  
879 c show quadrats at a site with high, pre-fire native perennial cover weeks before and days  
880 after the Hot Pot fire, which burned at high severity at that site.

881 **Figure 3.** Conceptual diagram of the hypotheses tested in this study.

882 **Figure 4.** Panel a is a path model showing support for the various hypotheses depicted in  
883 Figure 3. Red arrows are negative relationships, blue arrows are positive relationships, and  
884 grey arrows are not significant ( $p > 0.05$ ) but still accounted for in the model. Abbreviations:  
885 pre = pre-fire; post = post-fire; cv = cover; elv = elevation; ag = aboveground; sb = seed bank;  
886 sev = severity; div = diversity. On the left side of (b), burn severity (dNBR) as predicted by  
887 total vegetation cover (TVC; the sum of live and dead, shrub and herbaceous cover). On  
888 the right, burn severity is predicted by modelled TVC. In (c), fuel connectivity three years  
889 post-fire is modelled by seedbank composition, elevation and pre-fire aboveground species  
890 richness. In (d) Shannon-Weaver diversity index of the aboveground, post-fire community

composition, was negatively affected by fuel connectivity after accounting for elevation. For  
a, c and d, lines are the fitted partial effects, points are the partial residuals, and dotted lines  
are the 95% confidence intervals.  $p < 0.05$  for black lines,  $p > 0.05$  for grey lines. Panel e  
shows the modeled occurrence of germinable seeds for all species found at more than one  
location along a gradient of burn severity, after accounting for soil depth, aspect, elevation  
and pre-fire diversity. Black line is the mean prediction, each colored line represents one  
posterior sample.




Figure 1: .




Figure 2: .




Figure 3: .




Figure 4: .